

APPROXIMATION AND
CONVERGENCE OF THE
FIRST INTRINSIC VOLUME

RYNARTICE 2014

HERBERT JDELSBRUNNER

IST AUSTRIA

- I AN EXAMPLE
- II INTRINSIC VOLUME
- III PERSISTENT HOMOLOGY
- IV CONVERGENCE


I AN EXAMPLE

II INTRINSIC VOLUME

III PERSISTENT HOMOLOGY


IV CONVERGENCE

I.1 VOLUME OF UNIT BALL




$$B^3 : \|x\| \leq 1.$$

I.1 VOLUME OF UNIT BALL




$$B^3 : \|x\| \leq 1.$$




B_t^3 is resolution- t approximation

I.1 VOLUME OF UNIT BALL




$B^3 : \|x\| \leq 1.$




B_t^3 is resolution- t approximation

I.1 VOLUME OF UNIT BALL




$B^3: \|x\| \leq 1.$




B_t^3 is resolution- t approximation

$$\lim_{t \rightarrow 0} \text{Vol}(B_t^3) = \lim_{t \rightarrow 0} t^3 \#(B^3 \cap t\mathbb{Z}^3) = \text{Vol}(B^3) = \frac{4}{3}\pi$$

I.2 AREA OF UNIT BALL




$\mathbb{B}^3 : \|x\| \leq 1.$




\mathbb{B}_t^3 is resolution- t approximation

I.2 AREA OF UNIT BALL




$B^3 : \|x\| \leq 1.$




B_t^3 is resolution- t approximation

I.2 AREA OF UNIT BALL




$B^3: \|x\| \leq 1.$




B_t^3 is resolution- t approximation

$$\lim_{t \rightarrow 0} \text{Area}(B_t^3) = \lim_{t \rightarrow 0} 6t^2 \#(B^2 \cap t\mathbb{Z}^2) = 6 \text{Area}(B^2) = 6\pi$$

I.3 MEAN CURVATURE OF UNIT BALL




$B^3 : \|x\| \leq 1.$




B_t^3 is resolution- t approximation

I.3 MEAN CURVATURE OF UNIT BALL




$B^3 : \|x\| \leq 1.$




B_t^3 is resolution- t approximation

I.3 MEAN CURVATURE OF UNIT BALL



$B^3 : \|x\| \leq 1.$



B_t^3 is resolution- t approximation

$$\lim_{t \rightarrow 0} \text{Mean}(B_t^3) = \lim_{t \rightarrow 0} 3\pi t \#(B_t^1 \cap \mathbb{Z}^1) = 3\pi \text{Length}(B^1) = 6\pi$$

I AN EXAMPLE

II INTRINSIC VOLUME

III PERSISTENT HOMOLOGY

IV CONVERGENCE

II.1 STEINER POLYNOMIAL

(1796-1863)

$$M \subseteq \mathbb{R}^n.$$

II.1 STEINER POLYNOMIAL

(1796-1863)

$$M \subseteq \mathbb{R}^n.$$

The Steiner polynomial of M is

$$\text{Vol}(M+rB^n) = \sum_{k=0}^n b_k V_{n-k}(M) r^k$$

II.1 STEINER POLYNOMIAL

(1796-1863)

$$M \subseteq \mathbb{R}^n.$$

The Steiner polynomial of M is

$$\text{Vol}(M+rB^n) = \sum_{k=0}^n b_k V_{n-k}(M) r^k$$

/ /
volume (n-k)-th
of B^k intrinsic volume

II.1 STEINER POLYNOMIAL

(1796-1863)

$$M \subseteq \mathbb{R}^n.$$

The Steiner polynomial of M is

$$\text{Vol}(M+rB^n) = \sum_{k=0}^n b_k V_{n-k}(M) r^k$$

/ /
volume (n-k)-th
of B^k intrinsic volume

in \mathbb{R}^3 : $V_3 = \text{volume}$, $V_1 = \frac{1}{\pi}$ mean curvature
 $V_2 = \frac{1}{2}$ area, $V_0 = \frac{1}{4\pi}$ Gaussian curvature

II.1 STEINER POLYNOMIAL

(1796-1863)

$$M \subseteq \mathbb{R}^n.$$

The Steiner polynomial of M is

$$\text{Vol}(M+r\mathbb{B}^n) = \sum_{k=0}^n b_k V_{n-k}(M) r^k$$

Hadwiger (1951): characterization theorem.

Weyl (1939): generalization to tubes.

Federer (1969): to positive reach.

II.2 GRASSMANIAN

(1809-1877)

$\mathcal{L}_k^n \subseteq \mathcal{E}_k^n$ spaces of k -planes in \mathbb{R}^n

II.2 GRASSMANIAN

(1809-1877)

$$\begin{array}{ccc} \mathcal{L}_k^n & \subseteq & \mathcal{E}_k^n \\ / & & | \\ \text{linear} & & \text{affine} \end{array} \quad \text{spaces of } k\text{-planes in } \mathbb{R}^n$$

II.2 GRASSMANIAN

(1809-1877)

$$\begin{array}{ccc} \mathcal{L}_k^n \subseteq \mathcal{E}_k^n & \text{spaces of } k\text{-planes in } \mathbb{R}^n \\ / & | \\ \text{linear} & \text{affine} \end{array}$$

Haar measure normalized s.t.

$$\nu(\mathcal{L}_k^n) = 1,$$

II.2 GRASSMANIAN

(1809-1877)

$$\begin{array}{ccc} \mathcal{L}_k^n \subseteq \mathcal{E}_k^n & \text{spaces of } k\text{-planes in } \mathbb{R}^n \\ \downarrow & \downarrow \\ \text{linear} & \text{affine} \end{array}$$

Haar measure normalized s.t.

$$\nu(\mathcal{L}_k^n) = 1,$$

$$\mu(E \in \mathcal{E}_k^n \mid E \cap B^n \neq \emptyset) = b_{n-k}.$$

II.3 CROFTON FORMULA

(1826-1915)

$$V_{n-k}(M) = c_{k,n} \cdot \int_{E \in \mathcal{E}_k^n} \chi(M \cap E) dE$$

II.3 CROFTON FORMULA

(1826-1915)

$$V_{n-k}(M) = \frac{C_{k,n}}{\binom{n}{k} \frac{b_n}{b_k \cdot b_{n-k}}} \int_{E \in \mathcal{E}_k^n} \chi(M \cap E) dE$$

II.3 CROFTON FORMULA

(1826-1915)

$$V_{n-k}(M) = c_{k,n} \cdot \int_{E \in \mathcal{E}_k^n} \chi(M \cap E) dE$$

$$\frac{\binom{n}{k} b_n}{b_k \cdot b_{n-k}}$$

$$V_{n-k}(\mathbb{B}^n) = \binom{n}{k} \frac{b_n}{b_k}.$$


II.3 CROFTON FORMULA

(1826-1915)

$$V_{n-k}(M) = c_{k,n} \cdot \int_{E \in \mathcal{E}_k^n} \chi(M \cap E) dE$$

$$\binom{n}{k} \frac{b_n}{b_k \cdot b_{n-k}}$$

$$V_{n-k}(\mathbb{B}^n) = \binom{n}{k} \frac{b_n}{b_k}$$



I AN EXAMPLE


II INTRINSIC VOLUME

III PERSISTENT HOMOLOGY


IV CONVERGENCE

III.1 HEIGHT FUNCTION



III.1 HEIGHT FUNCTION



III.1 HEIGHT FUNCTION



III.1 HEIGHT FUNCTION




level set

$$f^{-1}(x)$$

sublevel set

$$f^{-1}(-\infty, x]$$


III.1 HEIGHT FUNCTION



superlevel set $f^{-1}[x, \infty)$
level set $f^{-1}(x)$
sublevel set $f^{-1}(-\infty, x]$


III.2 LEVEL SETS

level set
 $f^{-1}(x)$




III.2 LEVEL SETS

level set
 $f^{-1}(x)$




III.2 LEVEL SETS

sublevel set $f^{-1}(-\infty, x]$ level set $f^{-1}(x)$




III.2 LEVEL SETS

sublevel set $f^{-1}(-\infty, x]$ level set $f^{-1}(x)$ superlevel set $f^{-1}[x, \infty)$




1	1	0	0	0	0	1	1	0
1	1	0	1	0	0	0	1	0
1	0	0	2	0	0	0	1	0
2	0	0	3	0	0	0	1	0
2	0	0	2	0	0	0	0	0
1	0	0	1	0	0	0	0	0
0	0	0	0	0	0	0	0	0


III.3 DIAGRAM




III.3 DIAGRAM




III.3 DIAGRAM




III.3 DIAGRAM




III.3 DIAGRAM




III.3 DIAGRAM



III.3 DIAGRAM STABILITY



III.3 DIAGRAM STABILITY



THM. $W_{\infty}(\text{Dgm}(f), \text{Dgm}(g)) \leq \|f - g\|_{\infty}.$

III.4

MOMENT

$$\chi(f^{-1}(x_1)) = \sum_{k=0}^n (-1)^k [\#U_{p_k}(f, x) - \#D_{n_k}(f, x)].$$

III.4

MOMENT

$$\chi(f^{-1}(x)) = \sum_{k=0}^n (-1)^k [\#U_{p_k}(f, x) - \#D_{n_k}(f, x)].$$

The χ -moment of $D_{gm}(f)$ is

$$X(f) = \sum_{k=0}^n (-1)^k \sum_{A \in D_{gm}^k} (x_d - x_b) = \int_{x=-\infty}^{\infty} \chi(f^{-1}(x)) dx$$

III.4 MODIFIED MOMENT

$$\chi(f^{-1}(x)) = \sum_{k=0}^n (-1)^k [\#U_{p_k}(f, x) - \#D_{n_k}(f, x)].$$

The modified χ -moment of $D_{gm}(f)$ and $\epsilon > 0$ is

$$X(f, \epsilon) = \sum_{k=0}^n (-1)^k \sum_{\substack{A \in D_{gm}^k \\ |x_d - x_b| > \epsilon}} (x_d - x_b) = \int_{x=-\infty}^{\infty} \chi(f^{-1}(x)) dx$$

III.5

INTRINSIC VOLUME

The 1st intrinsic volume
of $M \subseteq \mathbb{R}^n$ is

$$V_1(M) = c_{n-1,n} \cdot \int_{E \in \mathcal{E}_{n-1}^n} \chi(M \cap E) dE$$

III.5

INTRINSIC VOLUME

The 1st intrinsic volume
of $M \subseteq \mathbb{R}^n$ is

$$\begin{aligned} V_1(M) &= c_{n-1,n} \cdot \int_{E \in \mathcal{E}_{n-1}^n} \chi(M \cap E) \, dE \\ &= c_{n-1,n} \cdot \int_{L \in \mathcal{L}_{n-1}^n} \int_{x=-\infty}^{\infty} \chi(f_L^{-1}(x)) \, dx \, dL \end{aligned}$$

III.5

INTRINSIC VOLUME

The 1st intrinsic volume
of $M \subseteq \mathbb{R}^n$ is

$$\begin{aligned}
 V_1(M) &= c_{n-1,n} \cdot \int_{E \in \mathcal{E}_{n-1}^n} \chi(M \cap E) \, dE \\
 &= c_{n-1,n} \cdot \int_{L \in \mathcal{L}_{n-1}^n} \int_{x=-\infty}^{\infty} \chi(f_L^{-1}(x)) \, dx \, dL \\
 &= c_{n-1,n} \cdot \int_{L \in \mathcal{L}_{n-1}^n} X(f_L) \, dL.
 \end{aligned}$$

III.5 MODIFIED INTRINSIC VOLUME

DEF. The modified 1st intrinsic volume of $M \subseteq \mathbb{R}^n$ and $\varepsilon > 0$ is

$$\begin{aligned} V_1(M, \varepsilon) &= c_{n-1, n} \cdot \int_{E \in \mathcal{E}_{n-1}^n} \chi(M \cap E) dE \\ &= c_{n-1, n} \cdot \int_{L \in \mathcal{L}_{n-1}^n} \int_{x=-\infty}^{\infty} \chi(f_L^{-1}(x)) dx dL \\ &= c_{n-1, n} \cdot \int_{L \in \mathcal{L}_{n-1}^n} X(f_L, \varepsilon) dL. \end{aligned}$$

I AN EXAMPLE

II INTRINSIC VOLUME

III PERSISTENT HOMOLOGY


IV CONVERGENCE

IV.1 UNIT BALL

THM. $|V_1(\mathbb{B}^n) - V_1(\mathbb{B}_{t,1}^n, t\sqrt{n})| \leq c_{n-1,n} \cdot t\sqrt{n}.$


IV.1 UNIT BALL

THM. $|V_1(\mathbb{B}^n) - V_1(\mathbb{B}_{\frac{1}{2}}^n, t\sqrt{n})| \leq c_{n-1,n} \cdot t\sqrt{n}.$




IV.1 UNIT BALL

THM. $|V_1(B^n) - V_1(B_{\frac{1}{t}}^n, t\sqrt{n})| \leq c_{n-1,n} \cdot t\sqrt{n}.$




IV.1 UNIT BALL

THM. $|V_1(\mathbb{B}^n) - V_1(\mathbb{B}_t^n, t\sqrt{n})| \leq c_{n-1,n} \cdot t\sqrt{n}.$



IV.1 UNIT BALL


THM. $|V_1(\mathbb{B}^n) - V_1(\mathbb{B}_t^n, t\sqrt{n})| \leq c_{n-1,n} \cdot t\sqrt{n}.$



$h_t : \mathbb{B}^n \rightarrow \mathbb{B}_t^n$ with distortion $\sup_{x \in \mathbb{B}^n} \|x - h_t(x)\| \leq \frac{1}{2} t\sqrt{n}.$


IV.1 UNIT BALL

THM. $|V_1(\mathbb{B}^n) - V_1(\mathbb{B}_{t\sqrt{n}}^n)| \leq c_{n-1,n} \cdot t\sqrt{n}.$



IV.1 UNIT BALL

THM. $|V_1(B^n) - V_1(B_{t\sqrt{n}}^n)| \leq c_{n-1,n} \cdot t\sqrt{n}.$




IV.1 UNIT BALL

THM. $|V_1(B^n) - V_1(B_{t/\sqrt{n}}^n)| \leq c_{n-1,n} \cdot t\sqrt{n}.$

$(-a, a)$ with
 $1 - \frac{1}{2}t\sqrt{n} \leq a \leq 1 + \frac{1}{2}t\sqrt{n}$

$$2a - 2 \leq t\sqrt{n}$$



IV.2 SOLID BODIES

DEF. A solid body is a compact set $M \subseteq \mathbb{R}^n$ s.t.

IV.2 SOLID BODIES

DEF. A solid body is a compact set $M \subseteq \mathbb{R}^n$ s.t.

(i) ∂M is a smoothly embedded $(n-1)$ -manifold,

IV.2 SOLID BODIES


DEF. A solid body is a compact set $M \subseteq \mathbb{R}^n$ s.t.

- (i) ∂M is a smoothly embedded $(n-1)$ -manifold,
- (ii) $\#Dgm(f) \leq C$ for every height function.

IV.2 SOLID BODIES

DEF. A solid body is a compact set $M \subseteq \mathbb{R}^n$ s.t.


- (i) ∂M is a smoothly embedded $(n-1)$ -manifold,
- (ii) $\#Dgm(f) \leq C$ for every height function.



IV.2 SOLID BODIES

DEF. A **solid body** is a compact set $M \subseteq \mathbb{R}^n$ s.t.

- (i) ∂M is a smoothly embedded $(n-1)$ -manifold,
- (ii) $\#D_{gm}(f) \leq C$ for every height function.



$$\text{reach}(\partial M) \cdot \kappa_{\max}(\partial M) \leq 1.$$

IV.2 SOLID BODIES


DEF. A **solid body** is a compact set $M \subseteq \mathbb{R}^n$ s.t.

- (i) ∂M is a smoothly embedded $(n-1)$ -manifold,
- (ii) $\#D_{gm}(f) \leq C$ for every height function.

THM. Solid body $M \subseteq \mathbb{R}^n$, $\forall k_{max} \geq 3tn$, $reach > 2t\sqrt{n}$, C finite.


$$\text{Then } |V_1(M) - V_1(M_t, 4t\sqrt{n})| \leq c_{n-1,n} \cdot 4t\sqrt{n} \cdot C.$$

IV.3 DISTORTED NORMAL BUNDLE




IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$



IV.3 DISTORTED NORMAL BUNDLE


$$Q^n = [-1, +1]^n / (2\sqrt{n})$$



IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$


$$Q_n: S^{n-1} \rightarrow S^{n-1}$$



IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$

$$Q_n: S^{n-1} \rightarrow S^{n-1}$$



$$\sin \varphi_n = \sqrt{\frac{1}{4n}}$$

$$\cos \varphi_n = \sqrt{\frac{3n+1}{4n}}$$


IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$

$$Q_n: S^{n-1} \rightarrow S^{n-1}$$

$$\langle u, e_i \rangle = \sin \varphi_n \quad \text{iff}$$

$$\langle Q_n(u), e_i \rangle = 0$$



$$\sin \varphi_n = \sqrt{\frac{1}{4n}}$$

$$\cos \varphi_n = \sqrt{\frac{3n+1}{4n}}$$


IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$

$$Q_n: S^{n-1} \rightarrow S^{n-1}$$

$$\langle u, e_i \rangle \leq \sin \psi_n \quad \text{iff}$$

$$\langle Q_n(u), e_i \rangle = 0$$



$$\langle u, Q_n(u) \rangle \geq \cos \psi_n$$

$$\sin \psi_n = \sqrt{\frac{1}{4n}}$$

$$\cos \psi_n = \sqrt{\frac{3n+1}{4n}}$$


IV.3 DISTORTED NORMAL BUNDLE

$$Q^n = [-1, +1]^n / (2\sqrt{n})$$

$$Q_n: S^{n-1} \rightarrow S^{n-1}$$

$$\langle u, e_i \rangle \leq \sin \psi_n \iff$$

$$\langle Q_n(u), e_i \rangle = 0$$



$$\langle u, Q_n(u) \rangle \geq \cos \psi_n$$


$$\arccos \langle Q_n(u), Q_n(v) \rangle$$

$$\leq 2 \arccos \langle u, v \rangle.$$


$$\sin \psi_n = \sqrt{\frac{1}{4n}}$$

$$\cos \psi_n = \sqrt{\frac{3n+1}{4n}}$$


IV.4 SYMMETRIC DIFFERENCE



IV.4 SYMMETRIC DIFFERENCE




IV.4 SYMMETRIC DIFFERENCE



Coord.-aligned normal bundle

maps $x \in \partial M \mapsto Q_n(N(x))$.

IV.4 SYMMETRIC DIFFERENCE




Coord.-aligned normal bundle

maps $x \in \partial M \mapsto Q_n(N(x))$.

$$L_x = \{x + \lambda Q_n(N(x)) \mid -t\sqrt{n} \leq \lambda \leq t\sqrt{n}\}$$

$$F_x = L_x \cap [(M \setminus M_t) \cup (M_t \setminus M)].$$

IV.4 SYMMETRIC DIFFERENCE



Coord.-aligned normal bundle

maps $x \in \partial M \mapsto Q_n(N(x))$.


$$L_x = \{x + \lambda Q_n(N(x)) \mid -t\sqrt{n} \leq \lambda \leq t\sqrt{n}\}$$

$$F_x = L_x \cap [(M \setminus M_t) \cup (M_t \setminus M)].$$


If $\partial M \subseteq \mathbb{R}^n$ s.t. $\forall k_{\max} \geq 3t\sqrt{n}$, $\text{reach} > 2t\sqrt{n}$,

then $\{F_x \mid x \in \partial M\}$ is a fibration of symm. diff.

IV.5 DEFORMATION




IV.5 DEFORMATION



$$U_t = M \cup M_t$$

$$F: U_t \times [0, 1] \rightarrow U_t \text{ (to } M_t)$$

IV.5 DEFORMATION




$$U_t = M \cup M_t$$

$$F: U_t \times [0,1] \rightarrow U_t \text{ (to } M_t)$$

$$G: U_t \times [0,1] \rightarrow U_t \text{ (to } M)$$

IV.5 DEFORMATION




$$U_t = M \cup M_t$$
$$M \cong U_t \cong M_t$$


$$F: U_t \times [0,1] \rightarrow U_t \text{ (to } M_t)$$

$$G: U_t \times [0,1] \rightarrow U_t \text{ (to } M)$$

IV.5 DEFORMATION



IV.5 DEFORMATION




height functions

$$f_u: M \rightarrow \mathbb{R}$$

$$h_u: U_t \rightarrow \mathbb{R}$$

$$g_u: M_t \rightarrow \mathbb{R}$$

IV.5 DEFORMATION



height functions

$$f_u: M \rightarrow \mathbb{R}$$

$$h_u: U_t \rightarrow \mathbb{R}$$

$$g_u: M_t \rightarrow \mathbb{R}$$

$$F_r = H(f_u(-\infty, r])$$

$$H_r = H(h_u(-\infty, r])$$

$$G_r = H(g_u(-\infty, r])$$

IV.6 TOWERS

$$\mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+s} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+s} \rightarrow \dots \rightarrow 0$$

$$\mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+s} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+s} \rightarrow \dots \rightarrow 0$$

$$\mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+s} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+s} \rightarrow \dots \rightarrow 0$$


IV.6 TOWERS

$$\begin{array}{l} \mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+s} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+s} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+s} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+s} \rightarrow \dots \rightarrow 0 \end{array}$$

IV.6 TOWERS

$$\begin{array}{l} \mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+s} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+s} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+s} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+s} \rightarrow \dots \rightarrow 0 \end{array}$$

IV.6 TOWERS

$$\begin{array}{l} \mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+s} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+s} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+s} \rightarrow \dots \rightarrow 0 \\ \mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+s} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+s} \rightarrow \dots \rightarrow 0 \end{array}$$


\mathcal{F}, \mathcal{H} are s -interleaving $\Rightarrow W_\infty(\text{Dgm}(\mathcal{F}), \text{Dgm}(\mathcal{H})) \leq s$

IV.6 TOWERS

$$\begin{array}{l} \mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+g} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+g} \rightarrow \dots \rightarrow 0 \\ \mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+g} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+g} \rightarrow \dots \rightarrow 0 \\ \mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+g} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+g} \rightarrow \dots \rightarrow 0 \end{array}$$

$$\begin{aligned} \mathcal{F}, \mathcal{H} \text{ are } g\text{-interleaving} &\Rightarrow W_\infty(\mathrm{Dgm}(\mathcal{F}), \mathrm{Dgm}(\mathcal{H})) \leq g \\ &\dots \Rightarrow W_\infty(\mathrm{Dgm}(\mathcal{F}), \mathrm{Dgm}(\mathcal{G})) \leq 2g. \end{aligned}$$

IV.6 TOWERS

$$\begin{array}{l}
 \mathcal{F}: \quad 0 \rightarrow \dots \rightarrow F_r \rightarrow F_{r+g} \rightarrow \dots \rightarrow F^r \rightarrow F^{r+g} \rightarrow \dots \rightarrow 0 \\
 \mathcal{H}: \quad 0 \rightarrow \dots \rightarrow H_r \rightarrow H_{r+g} \rightarrow \dots \rightarrow H^r \rightarrow H^{r+g} \rightarrow \dots \rightarrow 0 \\
 \mathcal{G}: \quad 0 \rightarrow \dots \rightarrow G_r \rightarrow G_{r+g} \rightarrow \dots \rightarrow G^r \rightarrow G^{r+g} \rightarrow \dots \rightarrow 0
 \end{array}$$

\mathcal{F}, \mathcal{H} are g -interleaving $\Rightarrow W_\infty(\text{Dgm}(\mathcal{F}), \text{Dgm}(\mathcal{H})) \leq g$

$\dots \Rightarrow W_\infty(\text{Dgm}(\mathcal{F}), \text{Dgm}(\mathcal{G})) \leq 2g.$

$$g = t\sqrt{n}$$

THM. Solid body $M \subseteq \mathbb{R}^n$, $1/k_{\max} \geq 3t\sqrt{n}$, reach $> 2t\sqrt{n}$, C finite.

Then $|V_1(M) - V_1(M_t, 4t\sqrt{n})| \leq c_{n-1,n} \cdot 4t\sqrt{n} \cdot C.$

THANK YOU