

MATHEMATICAL ANALYSIS 3 (NMAI056)

summer term 2022/23

lecturer: Martin Klazar

**LECTURE 4 (March 8, 2023) PROOF OF FTAlg.  
COMPLETE SPACES. BAIRE'S THEOREM**

• *n*-th complex roots. First, we realize that when proving the existence of *n*-th roots of complex numbers, it suffices to restrict to odd *n* and to numbers with modulus 1, i.e., lying on the complex unit circle *S*.

**Exercise 1.** *Using the last two exercises given in the pervious lecture, prove that if for every  $u \in S$  and for every odd  $n \in \mathbb{N}$  there exists a  $v \in S$  such that  $v^n = u$ , then the following theorem holds.*

**Theorem 2 (*n*-th roots in  $\mathbb{C}$ ).** *Complex numbers contain all *n*-th roots, that is*

$$\forall u \in \mathbb{C} \forall n \in \mathbb{N} \exists v \in \mathbb{C} (v^n = u) .$$

**Proof.** By the previous exercise, we can assume that  $u \in S$  and that  $n \in \mathbb{N}$  is odd. We need to prove that the map

$$f(z) = z^n : S \rightarrow S ,$$

which is clearly continuous, is onto. We assume for contradiction that there is a number

$$w \in S \setminus f[S]$$

(i.e.,  $w$  has no *n*-th root). Since  $n$  is odd, also  $-w \in S \setminus f[S]$  (always  $f(-z) = -f(z)$ ). We consider the line  $\ell \subset \mathbb{C}$  going through  $w$

and  $-w$ . Then we have the partition

$$\mathbb{C} = A \cup \ell \cup B,$$

where  $A$  and  $B$  are open half-planes determined by the line  $\ell$ . By Exercise 3 below,  $A$  and  $B$  are disjoint open sets. By Exercise 4 below,  $(A \cup B) \cap S = S \setminus \{w, -w\}$ ,  $\{1, -1\} \subset f[S] \cap (A \cup B)$  and  $|A \cap \{1, -1\}| = 1$ . Thus, the sets  $A$  and  $B$  cut the set  $f[S]$  and it is disconnected. This contradicts Theorem 21 in the last lecture, because  $f[S]$  is the image of the connected set  $S$  (we proved its connectedness last time) by the continuous function  $f$  and is therefore connected.  $\square$

**Exercise 3.** *Prove that for every line  $\ell \subset \mathbb{C}$ ,  $\mathbb{C} \setminus \ell$  is the disjoint union of two open sets.*

**Exercise 4.** *Let  $\ell \subset \mathbb{C}$  be a line passing through the origin,  $\ell \cap S = \{w, -w\}$  and  $A$  and  $B$  are the open half-planes determined by it. Prove that  $(A \cup B) \cap S = S \setminus \{w, -w\}$  and that for every  $u \in S \setminus \{w, -w\}$ , the points  $u$  and  $-u$  lie in different half-planes  $A$  and  $B$ .*

We move on to the second step of the proof of FTAlg. Using compact subsets in  $\mathbb{C}$ , we deduce the FTAlg from the existence of  $n$ -th roots. Recall that the complex numbers  $\mathbb{C}$  are the MS  $(\mathbb{C}, |u - v|)$  which is isometric to the Euclidean space  $(\mathbb{R}^2, e_2)$ .

**Exercise 5.** *Prove that for every real numbers  $\alpha \leq \alpha'$  and  $\beta \leq \beta'$ , the rectangle*

$$R := \{a + bi \mid \alpha \leq a \leq \alpha' \wedge \beta \leq b \leq \beta'\}$$

*is a compact set.*

**Proposition 6 (reduction to  $n$ -th roots).** *If  $\mathbb{C}$  contains all  $n$ -th roots, then FTAlg holds – every non-constant complex polynomial has a root.*

**Proof.** Let

$$p(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$$

be a non-constant complex polynomial, that is,  $n \in \mathbb{N}$ ,  $a_j \in \mathbb{C}$  and  $a_n \neq 0$ . The function

$$f(z) := |p(z)|: \mathbb{C} \rightarrow [0, +\infty) \subset \mathbb{C}$$

is continuous. We prove that  $f(u) = 0$  for some  $u \in \mathbb{C}$ . Then also  $p(u) = 0$  and  $u$  is a root of the polynomial  $p(z)$ .

First we prove that  $f$  attains on its definition domain  $\mathbb{C}$  a minimum value  $f(u)$ , and then that  $f(u) = 0$ . Let the real number  $K > 0$  be so large that

$$\frac{K^n |a_n|}{2} > |a_0| \quad \text{and} \quad \sum_{j=0}^{n-1} |a_j| K^{j-n} < \frac{|a_n|}{2}.$$

Then for  $z \in \mathbb{C}$  we have the estimate that

$$\begin{aligned} |z| > K \Rightarrow f(z) = |p(z)| &\geq |z|^n \left( |a_n| - \sum_{j=0}^{n-1} |a_j| \cdot |z|^{j-n} \right) \\ &> |a_0| = |p(0)| = f(0). \end{aligned}$$

We define a rectangle

$$R := \{a + bi \mid -K \leq a, b \leq K\} \subset \mathbb{C}.$$

Clearly,  $z \in \mathbb{C} \setminus R \Rightarrow |z| > K$ . By Theorem 15 in the second lecture (maximum principle) and Exercise 5 there exists  $u \in R$  such that

$f(u) \leq f(v)$  for every  $v \in R$ . Since  $0 \in R$ ,  $f(u) \leq f(0)$ . By the above estimate we have that

$$\forall v \in \mathbb{C} \left( f(u) \leq f(v) \right) .$$

Thus  $f$  attains at  $u$  its smallest value on the whole  $\mathbb{C}$ .

We prove that  $f(u) = 0$ . For this purpose we express the polynomial  $p(z)$  by Exercise 7 in the form

$$p(z) = \sum_{j=0}^n b_j (z - u)^j ,$$

where  $b_j \in \mathbb{C}$  and  $b_n = a_n$ . Thus, in this expression,  $f(u) = |p(u)| = |b_0|$ . We assume for contrary that  $f(u) = |b_0| > 0$ . We find the first non-zero non-constant coefficient in the polynomial  $p(z)$  and write  $p(z)$  as

$$p(z) = b_0 + b_k (z - u)^k + \underbrace{b_{k+1} (z - u)^{k+1} + \dots + b_n (z - u)^n}_{q(z)} ,$$

where  $q \in \mathbb{C}[z]$ ,  $k \in \mathbb{N}$ ,  $b_0 \neq 0$  and  $b_k \neq 0$ . We use the assumption about  $n$ -th roots and take an  $\alpha \in \mathbb{C}$  such that

$$\alpha^k = -\frac{b_0}{b_k} .$$

It is clear that  $q(z) = o((z - u)^k)$  (for  $z \rightarrow u$ ), so that

$$\lim_{z \rightarrow u} q(z) (z - u)^{-k} = 0 .$$

So we can take a  $\delta \in (0, 1)$  such that for

$$v := u + \delta \alpha$$

one has that

$$|q(v)| < \delta^k \cdot \frac{|b_0|}{2}.$$

Then we get the contradiction that  $f(v) < f(u)$ :

$$\begin{aligned} f(v) = |p(v)| &= |b_0 + b_k \alpha^k \delta^k + q(v)| \\ &\stackrel{\text{def. of } \alpha}{=} |b_0(1 - \delta^k) + q(v)| \\ &\stackrel{\Delta\text{'s ineq. and mult. } |\cdot|}{\leq} |b_0|(1 - \delta^k) + |q(v)| \\ &\stackrel{|q(v)| < \dots}{<} |b_0|(1 - \delta^k/2) \\ &\stackrel{\delta \in (0, 1)}{<} |b_0| = f(u). \end{aligned}$$

So  $f(u) = 0$  and  $p(u) = 0$ . □

**Exercise 7.** Prove that for every  $n \in \mathbb{N}_0$  and any complex numbers  $a_0, a_1, \dots, a_n$  and  $u$  there exist complex numbers  $b_0, b_1, \dots, b_n$  such that  $b_n = a_n$  and the polynomial equality

$$\sum_{j=0}^n a_j z^j = \sum_{j=0}^n b_j (z - u)^j$$

holds.

• *Complete sets and complete MSs.* A MS  $(M, d)$  is *complete* if every Cauchy sequence  $(a_n) \subset M$  is convergent. A *Cauchy sequence*  $(a_n)$  is one such that

$$\forall \varepsilon \exists n_0 (m, n \geq n_0 \Rightarrow d(a_m, a_n) < \varepsilon).$$

A set  $X \subset M$  is *complete* if the subspace  $(X, d)$  is complete.

**Exercise 8.** Let  $(M, d)$  be MS and  $X \subset Y \subset M$ . Prove that a set  $X$  is complete in the MS  $(Y, d)$  if and only if it is complete in the MS  $(M, d)$ .

**Exercise 9.** Prove that the Cartesian product

$$(M \times N, d \times e)$$

of complete MSs  $(M, d)$  and  $(N, e)$  is a complete MS.

A basic example of a complete MS is the Euclidean space

$$(\mathbb{R}, e_1) = (\mathbb{R}, |x - y|),$$

which is complete due to the fact that every sequence  $(a_n) \subset \mathbb{R}$  is convergent if and only if it is Cauchy. By Exercise 9 all Euclidean spaces  $(\mathbb{R}^n, e_n)$ ,  $n \in \mathbb{N}$ , are complete. We can construct many complete MSs by means of the following simple result.

**Proposition 10 (closed subspaces).** *In every complete MS  $(M, d)$  every closed subset  $X \subset M$  is complete.*

**Proof.** Let  $(a_n) \subset X$  be a Cauchy sequence in the closed set  $X \subset M$  in the complete MS  $(M, d)$ . There exists  $a := \lim a_n \in M$ . Since  $X$  is a closed set,  $a \in X$  (closed sets are closed also to limits). So the set  $X$  is complete.  $\square$

**Exercise 11.** Let  $X \subset M$  be a compact set in a MS  $(M, d)$ . Prove that  $X$  is complete.

**Exercise 12.** Give an example of a complete and non-compact set  $X \subset \mathbb{R}$  in the Euclidean MS  $(\mathbb{R}, e_1)$ .

**Exercise 13.** Which of the following implications holds in a MS  $(M, d)$ ?

1.  $X \subset M$  is a complete set  $\Rightarrow X$  is closed.
2.  $X \subset M$  and  $Y \subset M$  are complete sets  $\Rightarrow X \cup Y$  is a complete set.
3.  $X \subset M$  and  $Y \subset M$  are complete sets  $\Rightarrow X \cap Y$  is a complete set.
4.  $X \subset M$  is a complete set  $\Rightarrow X$  is bounded.
5.  $X \subset M$  is finite  $\Rightarrow X$  is complete.

• *Baire's theorem.* The main result about complete MSs is, besides completeness of particular MSs, Baire's theorem: no complete MS is a countable union of sparse sets. A set  $X \subset M$  in a MS  $(M, d)$  is *sparse (in  $M$ )* if

$$\forall a \in M \forall r > 0 \exists b \in M \exists s > 0 \\ (B(b, s) \subset B(a, r) \wedge B(b, s) \cap X = \emptyset) .$$

In words, every ball in the MS  $(M, d)$  contains a subball disjoint to  $X$ .

Similarly, a set  $X \subset M$  in a MS  $(M, d)$  is *dense (in  $M$ )* if

$$\forall a \in M \forall r > 0 (B(a, r) \cap X \neq \emptyset) .$$

In words, every ball in the MS  $(M, d)$  contains an element of the set  $X$ .

**Exercise 14.** Let  $(M, d)$  be a MS and  $X \subset M$  be a subset. Prove the equivalence that

$$X \text{ is dense} \iff \forall a \in M \exists (a_n) \subset X (\lim a_n = a) .$$

**Proposition 15 (density and continuity).** *Let  $(M, d)$  and  $(N, e)$  be MSs,  $X \subset M$  be dense in  $M$  and let*

$$f, g: M \rightarrow N$$

*be continuous mappings such that  $f|X = g|X$  (their restrictions to the set  $X$  coincide). Then  $f = g$ .*

**Proof.** Let  $a \in M$  be an arbitrary point. Since  $X$  is dense, by the previous exercise there exists a sequence  $(a_n) \subset X$  such that  $\lim a_n = a$ . Using Heine's definition of continuity of functions and the assumption about  $f$  and  $g$ , we have that

$$f(a) = \lim f(a_n) = \lim g(a_n) = g(a).$$

So  $f = g$ . □

**Exercise 16.** *Prove that any finite union of sparse sets is a sparse set. Show by an example that this is not generally true for countable unions.*

**Exercise 17.** *Prove that the intersection of two dense sets, one of which is open, is a dense set. Show that this is not in general true if we omit the assumption of openness.*

For  $a \in M$  and real  $r > 0$ , the *closed ball*  $\overline{B}(a, r)$  in a MS  $(M, d)$  is the set

$$\overline{B}(a, r) := \{x \in M \mid d(a, x) \leq r\}.$$

**Exercise 18.** *Every closed ball  $\overline{B}(a, r)$  is a closed set. For every  $a \in M$  and  $r, s \in \mathbb{R}$  with  $0 < r < s$ ,*

$$\overline{B}(a, r) \subset B(a, s).$$



**Theorem 19 (Baire's).** *Let  $(M, d)$  be a complete MS and*

$$M = \bigcup_{n=1}^{\infty} X_n .$$

*Then for some  $n$ , the set  $X_n$  is not sparse. In other words, no complete metric space is a countable union of sparse sets.*

**Proof.** We assume that all sets  $X_n$  are sparse and deduce a contradiction. We construct a nested sequence  $(\overline{B}_n)$  of closed balls with centers converging to a point  $a \in M$  outside any  $X_n$ , which is clearly a contradiction.

Let  $B(b, 1) \subset M$  be an arbitrary ball. Since  $X_1$  is sparse, there exists an  $a_1 \in M$  and an  $s_1 > 0$  such that  $B(a_1, s_1) \subset B(b, 1)$  and  $B(a_1, s_1) \cap X_1 = \emptyset$ . We set

$$\overline{B}(a_1, r_1) := \overline{B}(a_1, \min(s_1/2, 1/2)) .$$

Then  $\overline{B}(a_1, r_1) \subset B(a_1, s_1)$ , thus  $\overline{B}(a_1, r_1) \cap X_1 = \emptyset$ , and  $r_1 \leq 1/2$ .

Suppose that we already defined the closed balls

$$\overline{B}(a_1, r_1) \supset \overline{B}(a_2, r_2) \supset \cdots \supset \overline{B}(a_n, r_n)$$

such that for  $i = 1, 2, \dots, n$ ,  $\overline{B}(a_i, r_i) \cap X_i = \emptyset$  and  $r_i \leq 2^{-i}$ . Since  $X_{n+1}$  is sparse, there exist  $a_{n+1} \in M$  and  $s_{n+1} > 0$  such that  $B(a_{n+1}, s_{n+1}) \subset B(a_n, r_n)$  and  $B(a_{n+1}, s_{n+1}) \cap X_{n+1} = \emptyset$ . We set

$$\overline{B}(a_{n+1}, r_{n+1}) := \overline{B}(a_{n+1}, \min(s_{n+1}/2, 2^{-n-1})) .$$

Then

$$\overline{B}(a_{n+1}, r_{n+1}) \subset \overline{B}(a_n, r_n) \cap B(a_{n+1}, s_{n+1}) ,$$

hence also  $\overline{B}(a_{n+1}, r_{n+1}) \cap X_{n+1} = \emptyset$ , and  $r_{n+1} \leq 2^{-n-1}$ .

The sequence  $(a_n) \subset M$  of the centers of the closed balls defined above is Cauchy, since

$$m \geq n \Rightarrow \overline{B}(a_m, r_m) \subset \overline{B}(a_n, r_n), \text{ so } d(a_m, a_n) \leq r_n \leq \frac{1}{2^n}.$$

We use completeness of the MS  $(M, d)$  and take the limit

$$a := \lim a_n \in M.$$

Since  $m \geq n \Rightarrow a_m \in \overline{B}(a_n, r_n)$  and since by Exercise 18 every  $\overline{B}(a_n, r_n)$  is a closed set, the limit  $a$  lies in every closed ball  $\overline{B}(a_n, r_n)$  and therefore in none of the sets  $X_n$ , which is a contradiction.  $\square$

Baire's theorem has many applications, of which we now mention only one. A point  $a \in M$  in a MS  $(M, d)$  is *isolated* if

$$\exists r > 0 (B(a, r) = \{a\}).$$

**Exercise 20.** *Prove that in any MS  $(M, d)$ ,*

*$a \in M$  is not an isolated point  $\iff \{a\} \subset M$  is a sparse set.*

**Corollary 21 (getting uncountability).** *Any complete MS  $(M, d)$  without isolated points is uncountable.*

**Proof.** Suppose for the contrary that  $M$  is countable. Then

$$M = \bigcup_{a \in M} \{a\}$$

is a countable union. Since each set  $\{a\}$  is sparse (by the previous exercise), we have a contradiction with Baire's theorem.  $\square$

THANK YOU FOR YOUR ATTENTION

Homework Exercises. Please send to Mgr. J. Rondoš, Ph.D. by the end of the coming Monday by e-mail (jakub.rondos@gmail.com) solutions to the Exercises 7, 11, 13, 16 and 20.