

# Intervals of Sign Regular Matrices

<sup>1</sup>Mohammad Adm, <sup>2</sup>Jürgen Garloff, and <sup>1</sup>Jihad Titi

<sup>1</sup> Department of Mathematics and Statistics, University of Konstanz,  
D-78464 Konstanz, Germany  
mjamathe@yahoo.com, jihadtiti@yahoo.com

<sup>2</sup> Faculty of Computer Sciences, University of Applied Sciences/ HTWG Konstanz,  
D-78462 Konstanz, Germany  
garloff@htwg-konstanz.de

**Keywords:** Interval matrix, checkerboard ordering, totally nonnegative matrix, sign regular matrix.

We say that a class  $\mathcal{C}$  of  $n$ -by- $n$  matrices possesses the *interval property* if for any  $n$ -by- $n$  interval matrix  $[A] = [\underline{A}, \overline{A}] = ([a_{ij}, \bar{a}_{ij}])_{i,j=1,\dots,n}$  the membership  $[A] \subseteq \mathcal{C}$  can be inferred from the membership to  $\mathcal{C}$  of a specified set of its vertex matrices; here a *vertex matrix* of  $[A]$  is a real matrix  $B = (b_{ij})_{i,j=1,\dots,n}$  with  $b_{ij} \in \{a_{ij}, \bar{a}_{ij}\}$  for all  $i, j = 1, \dots, n$ . Examples of such classes include the

- $M$ -matrices or, more generally, inverse-nonnegative matrices [9], where only the bound matrices  $\underline{A}$  and  $\overline{A}$  are required to be in the class;
- inverse  $M$ -matrices [8], where all vertex matrices are needed;
- positive definite matrices [3], [12], where a subset of cardinality  $2^{n-1}$  is required (here only symmetric matrices in  $[A]$  are considered).

A class of matrices which in the nonsingular case are somewhat related to the inverse nonnegative matrices are the totally nonnegative matrices. A real matrix is called *totally nonnegative* if all its minors are nonnegative. Such matrices arise in a variety of ways in mathematics and its applications, e.g., in differential and integral equations,

numerical mathematics, combinatorics, statistics, and computer aided geometric design. For background information we refer to the recently published monographs [4], [11]. The second author posed in 1982 the conjecture that the set of the nonsingular totally nonnegative matrices possesses the interval property, where only two vertex matrices are involved [5], see also [4, Section 3.2] and [11, Section 3.2]. The two vertex matrices are the bound matrices with respect to the checkerboard ordering which is obtained from the usual entry-wise ordering in the set of the square matrices of fixed order by reversing the inequality sign for each entry in a checkerboard fashion. In our talk we apply the Cauchon algorithm [2] (also called deleting derivation algorithm [7] and Cauchon reduction algorithm [10]) to settle the conjecture. We also obtain the result that a fixed zero-nonzero pattern of the minors stays unchanged through an interval of nonsingular totally nonnegative matrices.

As a generalization of the totally nonnegative matrices we further consider *sign regular* matrices, i.e., matrices with the property that all their minors of fixed order have one specified sign or are allowed also to vanish. We identify some subclasses of the sign regular matrices which exhibit the interval property. The subclasses which require to check only two vertex matrices include the following sets (here it is understood that the two bound matrices have the same signature of their minors):

- the strictly sign regular matrices, i.e., the matrices with the property that all their minors of fixed order have one (strict) specified sign;
- the nonsingular almost strictly sign regular matrices, a class in between the nonsingular sign regular matrices and the strictly sign regular matrices;
- the tridiagonal nonsingular sign regular matrices;
- the nonsingular totally nonpositive matrices, i.e., the matrices with

the property that all their minors are nonpositive.

In some instances, the assumption of nonsingularity can be somewhat relaxed. These results lead us to the following new conjecture: Assume that the two bound matrices with respect to the checkerboard ordering are nonsingular and sign regular; then all matrices lying between the two bound matrices are nonsingular and sign regular, too. It was shown in [6] that the conclusion is true if we consider instead of the two bound matrices a set of vertex matrices with the cardinality of at most  $2^{2n-1}$  ( $n$  being the order of the matrices).

## References and Literature for Further Reading

- [1] M. ADM AND J. GARLOFF, Intervals of totally nonnegative matrices, *Linear Algebra Appl.*, 439:3796–3806, 2013.
- [2] M. ADM AND J. GARLOFF, Improved tests and characterizations of totally nonnegative matrices, *Electron. Linear Algebra*, 27:588–610, 2014.
- [3] S. BIALAS AND J. GARLOFF, Intervals of P-matrices and related matrices, *Linear Algebra Appl.*, 58:33–41, 1984.
- [4] S. M. FALLAT AND C. R. JOHNSON, *Totally Nonnegative Matrices*, Princeton Series in Applied Mathematics, Princeton University Press, Princeton and Oxford, 2011.
- [5] J. GARLOFF, Criteria for sign regularity of sets of matrices, *Linear Algebra Appl.*, 44:153–160, 1982.
- [6] J. GARLOFF, Vertex Implications for Totally Nonnegative Matrices, in *Total Positivity and its Applications*, M. Gasca and C.A. Micchelli, Eds., Kluwer Acad. Publ., Dordrecht, Boston, London, 103–107, 1996.
- [7] K. R. GOODEARL, S. LAUNOIS AND T. H. LENAGAN, Totally nonnegative cells and matrix Poisson varieties, *Adv. Math.*, 226:779–826, 2011.

- [8] C. R. JOHNSON AND R. S. SMITH, Intervals of inverse M-matrices, *Reliab. Comput.*, 8:239–243, 2002.
- [9] J. R. KUTTLER, A fourth-order finite-difference approximation for the fixed membrane eigenproblem, *Math. Comp.*, 25:237–256, 1971.
- [10] S. LAUNOIS AND T. H. LENAGAN, Efficient recognition of totally nonnegative matrix cells, *Found. Comput. Math.*, 14(2):371–387, 2014.
- [11] A. PINKUS, *Totally Positive Matrices*, Cambridge Tracts in Mathematics 181, Cambridge Univ. Press, Cambridge, UK, 2010.
- [12] J. ROHN, Positive definiteness and stability of interval matrices, *SIAM J. Matrix Anal. Appl.*, 15(1):175–184, 1994.