

# Simultaneous partitions of measures by $k$ -fans

IMRE BÁRÁNY\*

Rényi Institute of Mathematics  
Hungarian Academy of Sciences  
POBox 127, 1364 Budapest  
Hungary

and  
Department of Mathematics  
University College London  
Gower Street, London WC1E 6BT  
England

JIŘÍ MATOUŠEK\*\*

Department of Applied Mathematics  
Charles University  
Malostranské nám. 25, 118 00 Praha 1  
Czech Republic

## Abstract

A  $k$ -fan is a point in the plane and  $k$  semilines emanating from it. Motivated by a neat question of A. Kaneko and M. Kano, we study equipartitions by  $k$ -fans of two or more probability measures in the plane, as well as partitions in other prescribed ratios. One of our results: for any two measures there is a 4-fan such that one of its sectors contains  $\frac{2}{5}$  of both measures, and each of the the remaining three sectors contains  $\frac{1}{5}$  of both measures.

## 1 Introduction

For an integer  $k \geq 2$ , we define a  $k$ -fan as a point  $x$  (the *center*) in the plane and  $k$  semilines emanating from  $x$  (the *rays*); Fig. 1(a) shows an example of a 3-fan. A  $k$ -tuple of parallel lines, as in Fig. 1(b), is also considered to be a  $k$ -fan (this is a limit case for  $x$  receding to infinity). In this case, it is even possible that some of the parallel lines are also at the infinity. The  $k$  rays emanating from  $x$  are numbered as  $\ell_1, \ell_2, \dots, \ell_k$  in a cyclic order (clockwise or counterclockwise) around  $x$ . Each  $k$ -fan has an *orientation* (clockwise or counterclockwise) associated to it; for  $k \geq 3$ , the orientation is given by the labeling of the lines, and for  $k = 2$ , it is an extra information attached to the 2-fan.

The open angular sector between  $\ell_i$  and  $\ell_{i+1}$  is denoted by  $\sigma_i$  (for  $k = 2$ ,  $\sigma_1$  is the sector following after  $\ell_1$  in the given orientation of the 2-fan). For  $x$  at

---

\*Supported by Hungarian National Foundation Grants T 016391 and T 020914

\*\*Supported by Charles University grants No. 193 and 194.




Figure 1: Examples of 3-fans.

the infinity, the two unbounded regions together form one sector; see Fig. 1(b) (unless, of course, one of the rays is at the infinity). We also allow for  $\sigma_i = \emptyset$ ; in this case,  $\ell_i$  and  $\ell_{i+1}$  coincide. A  $k$ -fan is called *convex* if all of its sectors are convex.

Let  $\mu_1, \mu_2, \dots, \mu_m$  be Borel probability measures in the plane,  $m \geq 2$ , and let  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$  be a vector of nonnegative real numbers whose components sum up to 1. We say that a  $k$ -fan with sectors  $\sigma_1, \sigma_2, \dots, \sigma_k$   $\alpha$ -partitions the measure  $\mu_j$  if the following holds: For any  $i_1, i_2 \in \{1, 2, \dots, k\}$ , the open angular sector between  $\ell_{i_1}$  and  $\ell_{i_2}$  (in the sense given by the orientation of the  $k$ -fan) has  $\mu_j$ -measure at most  $\alpha_{i_1} + \alpha_{i_1+1} + \dots + \alpha_{i_2}$  (where the indices are taken in the cyclic order, with 1 following  $k$ ). If  $\mu_j$  is such that any line has  $\mu_j$ -measure 0, then this definition can of course be simplified to  $\mu_j(\sigma_i) = \alpha_i$ ,  $i = 1, 2, \dots, k$ . For measures partially concentrated on some of the  $\ell_i$ , the part of  $\mu_j$  on  $\ell_i$  can be arbitrarily divided between the adjacent sectors; this is captured by the rather complicated general definition above (which also covers the case of several  $\ell_i$ 's coinciding). If  $\alpha_1 = \alpha_2 = \dots = \alpha_k = \frac{1}{k}$  we speak of an *equipartition* of  $\mu_j$ .

In this paper, we investigate the following problem: for what combinations of  $m$ ,  $k$ , and  $\alpha$  can any  $\mu_1, \dots, \mu_m$  be simultaneously  $\alpha$ -partitioned by some  $k$ -fan?

This problem takes its origin from a very nice question of A. Kaneko and M. Kano [KK98]. Given an integer  $n \geq 2$  and two measures  $\mu_1$  and  $\mu_2$  in the plane (finite point sets, in fact) with  $\mu_1(\mathbf{R}^2) = \mu_2(\mathbf{R}^2) = n$ , does there exist a convex partition  $C_1, \dots, C_n$  of  $\mathbf{R}^2$  such that  $\mu_1(C_i) = \mu_2(C_i) = 1$  for all  $i$ . (As expected,  $C_1, \dots, C_n$  form a convex partition of  $\mathbf{R}^2$  by definition if the  $C_i$  are convex sets that are pairwise internally disjoint and their union is  $\mathbf{R}^2$ .) The case  $n = 2$  is easy: it is the planar ham-sandwich theorem. The case  $n = 3$  leads immediately to the problem whether a convex 3-fan equipartition exists for any two measures.

As we learned during the preparation of this paper, results answering Kaneko and Kano's original question, in various levels of generality, were proved, independent of our work, in several very recent papers. Akiyama et al. [ARNU98] prove the special case when the two measures are the surface area and perimeter of a plane convex body. Ito et al. [IUY98] show the case  $n = 3$  of the conjecture.

Finally Bespamyatnikh et al. [BKS98] and Sakai [Sa98], independently of each other, prove the conjecture in full generality. The argument of Bespamyatnikh et al. [BKS98] and Sakai [Sa98] is similar to a proof of the 2-dimensional Brouwer fixed-point theorem via Sperner’s lemma, but it is considerably more complicated. The method doesn’t seem to extend to  $\alpha$ -partitions with  $\alpha \neq (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ .

The following theorem summarizes our results.

**Theorem 1.1** *Let  $m \geq 2$ ,  $k \geq 2$ , and let  $\alpha = (\alpha_1, \dots, \alpha_k)$  have no zero component.*

- (i) (a) *For any  $k \geq 2$  and any  $\alpha$ , there are 4 measures that cannot be simultaneously  $\alpha$ -partitioned by a  $k$ -fan.*
- (b) *For any  $k \geq 3$  and any  $\alpha$ , there are 3 measures that cannot be simultaneously  $\alpha$ -partitioned by a  $k$ -fan.*
- (c) *For any  $k \geq 5$  and any  $\alpha$ , there are 2 measures that cannot be simultaneously  $\alpha$ -partitioned by a  $k$ -fan.*
- (d) *For  $k = 4$  and any  $\alpha$ , there are 2 measures that cannot be simultaneously  $\alpha$ -partitioned by a convex 4-fan.*
- (ii) (a) *Any 2 measures can be simultaneously  $\alpha$ -partitioned by a 2-fan, for all  $\alpha$ . The center of the 2-fan can be prescribed arbitrarily.*
- (b) *Any 3 measures can be simultaneously  $\alpha$ -partitioned by a 2-fan for  $\alpha = (\frac{1}{2}, \frac{1}{2})$  and for  $\alpha = (\frac{2}{3}, \frac{1}{3})$ .*
- (c) *Any 2 measures can be simultaneously equipartitioned by a 3-fan, or  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ -partitioned by a 3-fan. They can also be  $(\frac{2}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})$ -partitioned by a 4-fan.*

Note that the last statement in (ii)(c) implies that any two measures can be simultaneously  $(\frac{3}{5}, \frac{1}{5}, \frac{1}{5})$ -partitioned and  $(\frac{2}{5}, \frac{2}{5}, \frac{1}{5})$ -partitioned by a 3-fan. The results are summarized in Fig. 2.

The negative results in (i) are proved, using simple counterexamples, in Section 2. In parts (b) and (c), the reason for the non-existence of partitions is that  $k$ -fans do not have enough “degrees of freedom,” while in (a) and (d), there is a geometric constraint (although the degrees of freedom of 2-fans appear sufficient to partition 4 measures).

Part (ii)(a) follows by a rather simple averaging argument. Also the  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ -partition is very simple. The other positive results in (ii) are proved using equivariant topology.

**Previous work.** Equipartitions of measures in  $\mathbf{R}^d$  by simple geometric configuration have been studied in many papers. The primary example is the well-known ham-sandwich theorem, stating that any  $d$  measures in  $\mathbf{R}^d$  can be simultaneously bisected by a hyperplane. In the plane, any two measures can be simultaneously




Figure 2: Results on  $\alpha$ -partitioning  $m$  measures by  $k$ -fans ( $\frac{*}{5}$  means all combinations with denominators 5).

bisected by a line. Note that no  $\alpha$ -partition by a line with  $\alpha \neq (\frac{1}{2}, \frac{1}{2})$  is possible in general. This is shown by the example in Fig. 3, where one measure is uniformly distributed on the circle and the other one is concentrated in its center.




Figure 3: Two measures that cannot be  $\alpha$ -partitioned by a line for  $\alpha \neq (\frac{1}{2}, \frac{1}{2})$ .

An easy consequence of the 2-dimensional ham-sandwich theorem is the possibility of partitioning a measure in the plane into 4 equal parts by 2 lines. Various generalizations of this fact, most notably partitions of  $m$  measures in  $\mathbf{R}^d$  into  $2^k$  equal parts by  $k$  hyperplanes, have been considered by several researchers; see Ramos [Ram96] for recent results and references, and the survey by Živaljević [Živ] for a description of still newer results of Petrović et al. The most challenging problem in this area is probably partitioning a measure in  $\mathbf{R}^4$  into 16 equal parts by 4 hyperplanes, which remains still unsolved.

Another interesting equipartition result, namely equipartitioning a measure into 8 parts by a “cobweb” (two lines and a convex quadrilateral with vertices on the lines and surrounding the intersection of the lines), is due to Schulman [Sch93]. Makeev [Mak88] established the existence of 6-partitions by suitable cones in  $\mathbf{R}^3$ ; for example, for any measure, there is a cube  $C$  such that the six

cones with apex in the center of  $C$  and with the facets of  $C$  as bases form an equipartition.

In the literature we are aware of, all measure partition results of this type only concern equipartitions. For  $k$ -fan partitions, partitions other than equipartitions are sometimes possible, and this, in our opinion, makes the problem of partitioning by  $k$ -fans quite interesting.

## 2 Counterexamples

**Four measures.** To prove part (a) of Theorem 1.1(i), it is sufficient to consider 2-fans and arbitrary  $\alpha$ , since  $\alpha$ -partitions by  $k$ -fans imply  $\alpha'$ -partitions by  $(k-1)$ -fans by “omitting a ray.”

We consider 4 points  $p_1, \dots, p_4$  such that one of them is in the convex hull of the other 3 (Fig. 4). Let  $\mu_j$  be concentrated in  $p_j$ ,  $j = 1, \dots, 4$ . Let  $\alpha = (\alpha_1, \alpha_2)$




Figure 4: Four measures that cannot be partitioned by a 2-fan.

with  $\alpha_1 > 0$  and  $\alpha_2 > 0$ . If a 2-fan  $\alpha$ -partitions  $\mu_j$ , then  $p_j$  must lie on a ray of the 2-fan. But it is not possible that all of  $p_1, \dots, p_4$  simultaneously lie on the rays of a 2-fan.

**Three measures.** In part (b), it is enough to consider 3-fans. We choose a set  $P = \{p_{11}, p_{12}, p_{21}, p_{22}, p_{31}, p_{32}\}$  of 6 points in *strongly general position*, meaning that no 3 of them are collinear and no three lines determined by disjoint pairs of points of  $P$  have a common intersection (or are parallel). Note that such  $P$  cannot be simultaneously covered by the rays of a 3-fan (for at most 2 rays can cover 2 points each).

Next, given  $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ , we choose weights  $w_1, w_2 > 0$  with  $w_1 + w_2 = 1$  and  $w_1, w_2 \notin \{\alpha_1, \alpha_2, \alpha_3\}$ . The measure  $\mu_j$  is concentrated on  $p_{j1}$  and  $p_{j2}$ , and we put  $\mu_j(p_{ji}) = w_i$ ,  $i = 1, 2$ . We claim that if a 3-fan  $\alpha$ -partitions  $\mu_j$ , then *either one of  $p_{j1}, p_{j2}$  is the center of the 3-fan, or both  $p_{j1}$  and  $p_{j2}$  lie on rays of the 3-fan*. Indeed, supposing that the center is not one of  $p_{j1}, p_{j2}$ , it is clear that at least one of these points, say  $p_{j1}$ , must lie on a ray (see Fig. 5). If  $p_{j2}$  is inside a sector, then this sector must be adjacent to the ray containing  $p_{j1}$ , otherwise the sector would have the wrong measure  $w_2$ . But then there is a sector of measure 0. Thus, the claim holds, and from the strongly general position of  $P$ , it follows that a simultaneous  $\alpha$ -partition of  $\mu_1, \mu_2, \mu_3$  by a 3-fan is impossible.




Figure 5: A 2-point measure and a 3-fan.

**Two measures by 5-fans.** In part (c) of Theorem 1.1(i), it is enough to consider the case  $k = 5$  and  $\alpha = (\alpha_1, \dots, \alpha_5)$  with all  $\alpha_j > 0$ .

There is a very simple construction showing the impossibility of equipartitioning. Namely, choose a set  $P$  of 8 points in strongly general position. Let  $\mu_1$  be uniformly distributed on some 4 of the points of  $P$ , and let  $\mu_2$  be uniformly distributed on the remaining 4 points. Since the rays of a 5-fan can cover at most 7 points of  $P$ , there is a point of  $P$  lying inside some sector, and so the measure of that sector is at least  $\frac{1}{4}$ . Hence a 5-fan cannot equipartition these measures.

For  $\alpha$ -partitioning by a 5-fan with arbitrary  $\alpha$ , we use the following construction. Let  $\mu_1$  be the uniform measure whose support  $S_1$  is the segment  $[(-1, 1), (1, 1)]$ , and let  $\mu_2$  be the measure whose support  $S_2$  is the  $x$ -axis and whose distribution function  $F(t)$  is continuous with  $0 < F(t) < 1$  and, most importantly, no line intersects the graph of  $F$  graph in more than three points.

We claim that there is no  $\alpha$ -partition by a 5-fan for these two measures. Assume there is one. Then four consecutive rays must intersect each support as otherwise there are no five pieces. So the center is not between  $S_2$  and the line of  $S_1$ : it is either below  $S_2$  or above  $S_1$ . In both cases the four consecutive rays intersecting  $S_1$  intersect  $S_2$  as well. Denote the intersection points on  $S_2$  by  $x_1, x_2, x_3, x_4$  in this order. Computing the  $\mu_1$  measures of the sectors we get, with some positive  $h$  and suitable  $j$ ,

$$x_2 - x_1 = h\alpha_j, \quad x_3 - x_2 = h\alpha_{j+1}, \quad x_4 - x_3 = h\alpha_{j+2},$$

where  $j + 1, j + 2$  are taken mod 5. The  $\mu_2$  measures of the sectors give

$$F(x_2) - F(x_1) = \alpha_j, \quad F(x_3) - F(x_2) = \alpha_{j+1}, \quad F(x_4) - F(x_3) = \alpha_{j+2}.$$

This shows four points on the graph of  $F(t)$  contained in a line, namely the ones corresponding to  $x_1, x_2, x_3, x_4$ .

**Two measures by convex 4-fan.** For part (d), we use the previous setting but the second measure  $\mu_2$  will be different: let its support,  $S_2$ , be the segment  $[(-2, 0), (0, 2)]$  and let it be uniform on  $S_2$  for the time being. We will modify it soon. Assume there is convex  $\alpha$ -partition by a 4-fan. Then three consecutive rays intersect each support and the center cannot be between the two supports. It cannot be below  $S_2$  either because then one sector would meet  $S_2$  in an interval

too short to have the prescribed  $\mu_2$ -measure. So the center is above the line of  $S_1$ . Then there are three downward rays and the fourth goes upwards to make the partition convex. It is evident that the three downward rays, together with the center, are uniquely determined by  $\alpha$ . Now we modify  $\mu_2$  near the intersection of the middle downward ray with  $S_2$ , by pushing a little mass from the left to the right. This changes only the middle downward ray, and this ray will not pass through the intersection of the other two downward rays.

### 3 Preliminary reductions

In this section, we make some preliminary steps for the proof of the positive results in Theorem 1.1.

**A reformulation on the sphere.** First, we transfer the problem to the 2-dimensional sphere  $S^2$ . Let  $S^2$  be the unit sphere in  $\mathbf{R}^3$  centered at the origin, and let  $\mathbf{R}^2$  be embedded in  $\mathbf{R}^3$  as the horizontal plane  $\rho$  tangent to  $S^2$  from below (i.e. with equation  $z = -1$ ). Let  $\pi$  denote the central projection from the origin. This  $\pi$  gives a homeomorphism of the upper open hemisphere of  $S^2$  onto  $\rho$ , and similarly for the lower open hemisphere. A given Borel measure in  $\mathbf{R}^2$  is transferred by  $\pi^{-1}$  to a Borel measure in the upper open hemisphere.

A *k-fan* in  $S^2$  is a point  $x \in S^2$  and a collection  $(\ell_1, \ell_2, \dots, \ell_k)$  of great semicircles emanating from  $x$ , in such a way that the semicircles  $\ell_1, \dots, \ell_k$  are ordered clockwise around  $x$  when viewed from the center of the sphere. We write this *k-fan* as  $(x; \ell_1, \dots, \ell_k)$ . The sectors of a *k-fan* and  $\alpha$ -partition of a measure by a *k-fan* are defined in an obvious analogy to the planar case.

To any *k-fan*  $(x; \ell_1, \dots, \ell_k)$ , we assign a *k-fan* in the plane as follows. The center of the corresponding planar *k-fan* is  $\pi(x)$  (for  $x$  on the equator,  $\pi(x)$  is formally at the infinity). The image of the ray  $\ell_i$  is obtained as the intersection of  $\rho$  with the halfplane that intersects  $S^2$  in the great semicircle  $\ell_i$ . Also the orientation of the *k-fan* is transferred by the projection (hence *k-fans* with centers at the upper hemisphere induce counterclockwise planar *k-fans*, and *k-fans* with centers on the lower hemisphere induce clockwise planar *k-fans*).

If any  $m$  measures on  $S^2$  can be  $\alpha$ -partitioned by *k-fans*, then any  $m$  planar measures can be  $\alpha$ -partitioned by *k-fans*. In fact, the above correspondence of planar and spherical *k-fans* is bijective, but the spherical problem is more general, since the measures obtained from planar situations only live in the upper open hemisphere.

**Nice measures.** Next, we note that by standard arguments, we can restrict ourselves to special measures. We formulate a somewhat stronger result than we actually need.

**Lemma 3.1** *Let  $k$ ,  $m$ , and  $\alpha$  be given. Any  $m$  Borel probability measures in  $S^2$  can be  $\alpha$ -partitioned by a *k-fan* if (and only if) all  $m$ -tuples of measures from the*

following special classes can be so partitioned:

(i) Measures concentrated on finitely many points in  $S^2$ .

(ii) Measures that are absolutely continuous with respect to the Lebesgue measure and such that any nonempty open set has a strictly positive measure.

**Sketch of proof.** It follows easily from the results of Vapnik and Chervonenkis [VC71] that given a Borel probability measure  $\mu$  on  $S^2$ , for any  $\varepsilon > 0$  there is a measure  $\mu_\varepsilon$  concentrated on finitely many points such that for any sector  $\sigma$  (the open region delimited by two great semicircles), we have  $|\mu(\sigma) - \mu_\varepsilon(\sigma)| \leq \varepsilon$ . The argument is finished by letting  $\varepsilon \rightarrow 0$  and noting that the space of all spherical  $k$ -fans with the natural topology is compact.

The transformation to measures as in (ii) is used in almost all of the equipartition results in the literature. For a measure concentrated on finitely many points, we can replace each point by a spherical cap of radius  $\varepsilon$  with the appropriately scaled Lebesgue measure on it, and then add  $\varepsilon$ -times the Lebesgue measure on  $S^2$ . Letting  $\varepsilon \rightarrow 0$  and using compactness works again. Let us remark that for an arbitrary Borel probability measure, we can take convolution with a suitable measure  $\nu_\varepsilon$  (whose density function is a narrow peak) and obtain a measure as in (ii) directly.  $\square$

## 4 Easy positive results

First we prove part (a) of Theorem 1.1(ii),  $\alpha$ -partitions of 2 measures by 2-fans. This result can easily be proved by the methods below involving equivariant topology, but here we show a simple averaging argument suggested by Attila Pór. Fix the center  $x$  of the considered 2-fans, and let  $\gamma$  be the circle of unit length centered at  $x$ . We may assume that  $\mu_1$  and  $\mu_2$  are measures on  $\gamma$ , and by considerations analogous to those in the proof of Lemma 3.1, we may suppose that they are sufficiently nice. Namely, we suppose that after a suitable reparameterization of  $\gamma$ ,  $\mu_2$  is the one-dimensional Lebesgue measure on  $\gamma$ , and that  $\mu_1$  is given by a density function  $g$  on  $\gamma$  (with  $\int_\gamma g(t) dt = 1$ ).

We want to show that there is an arc  $a = (s, s + \alpha_1) \subset \gamma$  of length  $\alpha_1$  with  $\mu_1(a) = \alpha_1$ . Define the function  $f(s) = \mu_1((s, s + \alpha_1))$ . This is a continuous function, and if attains no value  $\alpha_1$  then it is always strictly below  $\alpha_1$  or always strictly above  $\alpha_1$ . But we have

$$\int_\gamma f(s) ds = \int_\gamma \int_s^{s+\alpha_1} g(t) dt ds = \int_\gamma \int_{t-\alpha_1}^t g(t) ds dt = \alpha_1.$$

This proves part (a) of Theorem 1.1(ii).

Another easy case is the  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ -partition of 2 measures by 3-fans in part (c) of Theorem 1.1(ii). Two measures can be simultaneously bisected by a line by the ham-sandwich theorem. We bisect the two halves in one of the halfplanes by another line, and we obtain a (convex) 3-fan providing the desired partition.

## 5 Tools from equivariant topology

We now interrupt our discussion of  $k$ -fans and discuss some general results from equivariant topology.

Let  $G$  be a group and  $X$  a topological space. We recall that an *action*  $\omega$  of  $G$  on  $X$  is a homomorphism from  $G$  into the group of homeomorphisms of  $X$ . Explicitly, for each  $g \in G$ , we have a homeomorphism  $\omega_g: X \rightarrow X$ , and  $\omega_g \circ \omega_h = \omega_{gh}$  holds for all  $g, h \in G$ . The action  $\omega$  is called *free* if for  $g \in G$  distinct from the unit element, the homeomorphism  $\omega_g$  has no fixed points.

If  $G = \mathbf{Z}_q$  is a cyclic group with generator  $g$ , it is sufficient to specify the single homeomorphism  $\omega_g$ . In this case, as is usual in the literature, we write  $\omega$  instead of  $\omega_g$  and we speak of “the  $\mathbf{Z}_q$ -action  $\omega$ .”

If  $X$  is a topological space with an action  $\omega$  of  $G$  and  $Y$  is another topological space with an action  $\nu$  of  $G$ , a  *$G$ -equivariant map* (with respect to  $\omega$  and  $\nu$ ), or simply a  *$G$ -map*, from  $X$  to  $Y$  is a continuous mapping  $f: X \rightarrow Y$  that commutes with the actions; that is,  $f \circ \omega_g = \nu_g \circ f$  for all  $g \in G$ .

There are numerous results in combinatorics and in geometry that are derived by showing the non-existence of an equivariant map between suitable topological spaces. This method was elaborated in a number of papers by Lovász, Alon, Bárány, Shlosman, Szücs, Sarkaria, Živaljević, Vrećica, Ramos, and others; a recent survey is [Živ97] (also see [Živ96] for an expanded version), and fairly advanced tools are discussed in its continuation [Živ].

The following theorem of Dold [Dol83] can be used to exclude the existence of equivariant mappings  $X \rightarrow Y$ , provided that the actions of  $G$  on both  $X$  and  $Y$  are free. It is a far-reaching generalization of the famous Borsuk–Ulam theorem (the latter claims that there is no  $\mathbf{Z}_2$ -equivariant mapping  $S^n \rightarrow S^{n-1}$ , where both spheres are considered with the antipodal actions  $x \mapsto -x$ ).

We recall that a topological space  $X$  is  *$n$ -connected* if for each  $j \leq k$ , any continuous mapping  $f: S^j \rightarrow X$  can be extended continuously to  $\bar{f}: B^{j+1} \rightarrow X$ , where  $B^{j+1}$  is the  $(j+1)$ -dimensional ball bounded by the  $S^j$ .

**Theorem 5.1** *Let  $G$  be a finite group,  $|G| > 1$ , let  $X$  be an  $n$ -connected space with a free action of  $G$ , and let  $Y$  be a (paracompact) topological space of dimension at most  $n$  with a free action of  $G$ . Then there is no  $G$ -equivariant map  $f: X \rightarrow Y$ .*

In our applications of the theorem, we will deal with a space  $X$  which is homeomorphic to the real projective space  $PR^3$ , and thus it is not even 1-connected (the fundamental group is  $\mathbf{Z}_2$ ). But there is a double-covering map  $\pi: S^3 \rightarrow PR^3$ , and  $S^3$  is 2-connected. The following lemma shows that for odd  $q$ , a  $\mathbf{Z}_q$ -action can be lifted from our  $X$  to  $S^3$ , and so  $X$  is as good as a 2-connected space in this case.

**Lemma 5.2** *Let  $\pi: X \rightarrow Y$  be a  $p$ -fold covering map, with  $X$  arcwise connected and simply connected, and let  $\omega$  be a  $\mathbf{Z}_q$ -action on  $Y$ , where  $(p, q) = 1$ . Then there is a  $\mathbf{Z}_q$ -action  $\tilde{\omega}$  on  $X$  such that  $\pi$  is a  $\mathbf{Z}_q$ -map. If  $\omega$  is free then  $\tilde{\omega}$  is free too.*

This simple result is most likely known, but we haven't found a reference.

**Proof.** Fix  $x_0 \in X$ . Let  $y_0 = \pi(x_0)$ ,  $y_1 = \omega(y_0)$ , and fix some  $x_1 \in \pi^{-1}(y_1)$ . Define a map  $\tilde{\omega}: X \rightarrow X$ , as follows. For  $x \in X$ , choose a path  $\gamma$  in  $X$  connecting  $x_0$  to  $x$ . The path  $\omega(\pi(\gamma))$  in  $Y$  has endpoints  $y_1$  and  $\omega(\pi(x))$ . Lift the endpoint  $y_1$  to  $x_1$ ; the other endpoint of this lifted path defines  $\tilde{\omega}(x)$ ; see Fig 6.




Figure 6: Lifting a  $\mathbf{Z}_q$ -action.

This is a well-defined mapping. If  $\gamma'$  is another path from  $x_0$  to  $x$ , then it is homotopic to  $\gamma$  (rel  $\{x_0, x\}$ ). Applying  $\pi$  and then  $\omega$  to the system of paths witnessing the homotopy of  $\gamma$  and  $\gamma'$  yields a system of homotopic paths from  $y_1$  to  $\omega(\pi(x))$ , and the liftings thus all have the same endpoints.

Clearly,  $\tilde{\omega}$  is continuous. By the construction, we have  $\tilde{\omega} \circ \pi = \pi \circ \omega$ . (From this, we also have  $\tilde{\omega}^k \circ \pi = \pi \circ \omega^k$ , for all  $k \geq 1$ .)

One can define a continuous inverse to  $\tilde{\omega}$ : to find the inverse image of  $z \in X$ , consider a path  $\gamma$  from  $x_1$  to  $z$ , apply  $\pi$ , then  $\omega^{-1}$ , then lift to  $X$  with  $y_0$  lifted to  $x_0$ . Thus,  $\tilde{\omega}$  is a homeomorphism.

Consider  $\nu = \tilde{\omega}^q$ . This is a homeomorphism lifting the identity. It acts on  $F = \pi^{-1}(y_0)$  as a permutation. Moreover, if some  $\nu^k$  has a fixed point then it is the identity map, as is easy to check. Hence all cycles of the permutation  $\nu|_F$  have the same length, and consequently  $\nu^k$  is the identity for any  $k$  divisible by  $p$ . Choosing  $k$  as a multiple of  $p$  with  $k \equiv 1 \pmod{q}$ , we get that  $\tilde{\omega} = \tilde{\omega}^k$  satisfies  $\tilde{\omega}^q = 1_X$  and  $\pi \circ \tilde{\omega} = \omega \circ \pi$ .  $\square$

Let us remark that, for example, a  $\mathbf{Z}_2$ -action generally cannot be lifted from  $PR^3$  to  $S^3$ .

## 6 Fan partitions by equivariant topology

Here we resume our discussion of (spherical)  $k$ -fans. From now on, we assume that the measures  $\mu_1, \dots, \mu_m$  on  $S^2$  are as in Lemma 3.1(ii). Actually, we use the following two properties: any nonempty sector has a strictly positive measure, and if the angle of a sector goes to 0, then its measure goes to 0.

**The candidate space and test maps.** Let  $q \geq 2$  be a given integer and let the probability measures  $\mu_1, \dots, \mu_m$  on  $S^2$  be fixed. To each pair  $x, y \in S^2$  of orthonormal unit vectors we assign a  $q$ -fan in  $S^2$ , as follows. The center is  $x$ , and  $\ell_1$  is the intersection of  $S^2$  with the halfplane having 0 and  $x$  on its boundary and containing  $y$ . The rays  $\ell_2, \dots, \ell_q$  are uniquely determined by the condition that the resulting  $q$ -fan equipartition the measure  $\mu_m$ . In this way, the space  $X_q$  of all  $q$ -fans equipartitioning  $\mu_m$  is identified with the space  $V_2(\mathbf{R}^3)$  of all pairs  $(x, y)$  of orthonormal vectors in  $\mathbf{R}^3$ , with the natural topology ( $V_2(\mathbf{R}^3)$  is called the *Stiefel manifold* of orthonormal 2-frames in  $\mathbf{R}^3$ ). Write  $\iota_q: V_2(\mathbf{R}^3) \rightarrow X_q$  for this identification, that is,

$$\iota_q(x, y) = (x; \ell_1, \dots, \ell_q).$$

It is easy to see that  $V_2(\mathbf{R}^3)$  is homeomorphic with the group  $SO(3)$  of rotations around the origin in  $\mathbf{R}^3$ , and as is well-known, this is homeomorphic with the projective space  $P\mathbf{R}^3$  (see e.g. Bredon [Bre93] p. 164).

For assessing the “quality” of a given  $q$ -fan from  $X_q$  with respect to the other measures, we introduce *test maps*  $f_j: X_q \rightarrow \mathbf{R}^q$ ,  $j = 1, 2, \dots, m-1$ , by letting

$$f_j(F) = \left( \mu_j(\sigma_1) - \frac{1}{q}, \mu_j(\sigma_2) - \frac{1}{q}, \dots, \mu_j(\sigma_q) - \frac{1}{q} \right),$$

where  $\sigma_1, \dots, \sigma_q$  are the sectors of the  $k$ -fan  $F$ . As is easy to check, the conditions on the measures  $\mu_j$  imply that each  $f_j$  is continuous. We also observe that the image of each  $f_j$  is actually contained in the hyperplane  $Z = \{y \in \mathbf{R}^q : y_1 + y_2 + \dots + y_q = 0\}$ .

If we want to prove the existence of a  $q$ -fan equipartitioning  $\mu_1$  through  $\mu_m$ , we must show that  $f_1, \dots, f_{m-1}$  have a common zero. More generally, for proving the existence of simultaneous  $\alpha$ -partitions by  $k$ -fans, where  $\alpha = (\frac{a_1}{q}, \frac{a_2}{q}, \dots, \frac{a_k}{q})$  with  $a_1, a_2, \dots, a_k$  being natural numbers summing up to  $q$ , it is sufficient to show that there is a  $q$ -fan  $F \in X_q$  such that  $f_j(F) \in L$  for all  $j = 1, 2, \dots, m-1$ , where  $L$  is the linear subspace

$$L = L(\alpha) = \{x \in \mathbf{R}^q : x_1 + x_2 + \dots + x_{a_1} = 0, x_{a_1+1} + \dots + x_{a_1+a_2} = 0, \dots, x_{a_1+\dots+a_{k-1}+1} + \dots + x_q = 0\}. \quad (1)$$

The results in Theorem 1.1 are obtained by considering suitable group actions on the candidate space  $X_q$  such that the test maps are equivariant, and then using the results in Section 5 for showing that no equivariant map missing  $L$  can exist.

**The group action.** On our space  $X_q$ , we have a natural free action  $\omega$  of the group  $\mathbf{Z}_q$  (integers with addition modulo  $q$ ). It is given by “turning by one sector;” formally,  $\omega(x; \ell_1, \dots, \ell_q) = (x; \ell_2, \ell_3, \dots, \ell_q, \ell_1)$ . Let  $F \in X_q$  be a  $q$ -fan, and let  $f_j(F) = (y_1, \dots, y_q) \in Z$ . We have  $f_j(\omega(F)) = (y_2, y_3, \dots, y_q, y_1)$ . Thus, if we define the  $\mathbf{Z}_q$ -action  $\nu$  on the hyperplane  $Z$  by  $\nu(y_1, y_2, \dots, y_q) = (y_2, y_3, \dots, y_q, y_1)$ , then  $f_j$  is equivariant.

We mention that there is a natural  $\mathbf{Z}_q$  action  $\rho_q$  on  $V_2(\mathbf{R}^3)$ :  $\rho_q(x, y) = (x, y')$  where  $y'$  is obtained from  $y$  by turning clockwise by angle  $2\pi/q$  around  $x$  when viewed from the center of the sphere. With this map the identification  $\iota_q$  becomes a  $\mathbf{Z}_q$  map:

$$\iota_q \circ \rho_q = \omega \circ \iota_q.$$

The advantage is that the  $\mathbf{Z}_q$  action does not depend on the measure  $\mu_m$  any more.

We should remark that the  $\mathbf{Z}_q$ -action  $\nu$  is free on  $Z \setminus \{0\}$  if and only if  $q$  is a prime, and so we can expect that Dold’s theorem 5.1 will only be applicable for prime  $q$ ’s.

Next, we discuss the few specific cases where we can prove the existence of  $k$ -fan partitions by the equivariant topology methods.

**Equipartition of 2 measures by 3-fans.** The possibility of equipartitioning 2 measures on  $\mathbf{R}^2$  by a 3-fan was proved in [BKS98], [IUY98], [Sa98]; they even get a convex 3-fan. Here we give a topological proof, of the slightly more general case of  $S^2$ , as it is the simplest case in our discussion. We consider the space  $X_3$  of 3-fans that equipartition  $\mu_2$  as introduced above. It suffices to show that there is no  $\mathbf{Z}_3$ -map  $f_1: X_3 \rightarrow Z \setminus \{0\}$ , where  $Z$  is the plane  $\{y_1 + y_2 + y_3 = 0\}$ . Supposing that  $f_1$  exists and using Lemma 5.2, we can lift the  $\mathbf{Z}_3$ -action from  $X_3$  to  $S^3$  (obtaining a free action), and define a  $\mathbf{Z}_3$ -map  $\tilde{f}_1: S^3 \rightarrow Z \setminus \{0\}$ . Finally, it is convenient (although not strictly necessary here) to reduce the dimension of the target space. Namely, if  $S(Z) = \{y \in Z : \|y\| = 1\}$  denotes the unit sphere in  $Z$  (an  $S^1$  in this case), we define  $h: Z \setminus \{0\} \rightarrow S(Z)$  by  $h(y) = \frac{y}{\|y\|}$ . Then  $h \circ \tilde{f}_1: S^3 \rightarrow S(Z)$  is a  $\mathbf{Z}_3$ -map. Since the  $\mathbf{Z}_3$ -action on  $S(Z)$  is free, Theorem 5.1 applies and excludes the existence of such a map (as  $S^3$  is 2-connected and  $S(Z)$  is 1-dimensional; this is even more than we need).

There is another, almost elementary, argument showing the non-existence of a  $\mathbf{Z}_3$ -map  $h \circ f_1: X_3 \rightarrow S^1$ , which we now describe. Assume there is such a map and combine it with the identification  $\iota_3$  to obtain a  $\mathbf{Z}_3$ -map  $g: V_2(\mathbf{R}^3) \rightarrow S^1$ . Consider the set  $S(a) = \{(a, y) \in V_2(\mathbf{R}^3)\}$  with  $a \in S^2$  fixed; this is an  $S^1$ , and so we have a  $\mathbf{Z}_3$ -map  $g_1: S^1 \rightarrow S(a)$ , say the isometric imbedding. Then  $g_1 \circ g: S^1 \rightarrow S^1$  is also a  $\mathbf{Z}_3$ -map, where the action on both  $S^1$ s is the rotation by  $2\pi/3$ . Then the degree of  $g_1 \circ g$  is 1 mod 3, as it is well known and easy to check (cf. [KZ75], or [BSS81]). Define now  $S^*$  as the union of two copies of  $S^1$  glued together at a single point. Define  $g_2: S^* \rightarrow V_2(\mathbf{R}^3)$  as winding twice around  $S^1$  while mapping to  $S(a) \subset V_2(\mathbf{R}^3)$  the same way as  $g_1$  does. The degree of  $g_2 \circ g$  is

clearly twice that of  $g_1 \circ g$ , and so it is 2 mod 3. But as the fundamental group of  $V_2(\mathbf{R}^3)$  is  $\mathbf{Z}_2$ , the cycle that goes around  $S(a)$  twice is homotopic to zero. Thus the degree of  $g_2 \circ g$  is zero. Contradiction.

Remark. Using this proof one can place extra restrictions on the 3-fan realizing the equipartition: any subset of  $V_2(\mathbf{R}^3)$  in which twice  $S(a)$  can be deformed to a point ought to contain such a 3-fan.

**Other partitions by Dold's theorem.** Another case where Dold's theorem can be applied are  $\alpha$ -partitions of 2 measures by 4-fans with  $\alpha = (\frac{2}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})$  (consequently, we get also all  $\alpha$ -partitions by 3-fans, where all the components of  $\alpha$  are fractions with denominator 5).

In this case, we choose  $q = 5$ , and we let  $L = L((\frac{2}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})) = \{y \in \mathbf{R}^5 : y_1 + y_2 = 0, y_3 = y_4 = y_5 = 0\}$  and  $\mathcal{L} = \{L, \nu(L), \nu^2(L), \dots, \nu^{q-1}(L)\}$ , where  $\nu$  is the coordinate shift action on  $\mathbf{R}^q$ . If the desired 4-fan  $\alpha$ -partition didn't exist, we would get a  $\mathbf{Z}_5$ -map  $f_1: X_5 \rightarrow Z \setminus \cup \mathcal{L}$ . By lifting the  $\mathbf{Z}_5$ -action from  $X_5$  to  $S^3$ , we obtain a  $\mathbf{Z}_5$ -map  $\tilde{f}_1: S^3 \rightarrow Z \setminus \cup \mathcal{L}$ , with free  $\mathbf{Z}_5$ -actions on both sides.

This time, even if we compose  $\tilde{f}_1$  with the mapping  $h: Z \setminus \{0\} \rightarrow S(Z)$  as in the previous case, the target space is still 3-dimensional; it is a  $S^3$  with 10 points (i.e. 5 copies of  $S^0$ ) deleted. The dimension is too large to apply Dold's theorem directly, but it can be easily reduced by 1, by equivariantly contracting  $Z \setminus \cup \mathcal{L}$  to a 2-dimensional subspace  $Y \subset S(Z)$ . In the following lemma, we give a simple geometric construction of a suitable  $Y$ .

**Lemma 6.1** *Let  $\mathcal{L}$  be a finite collection of linear subspaces of  $\mathbf{R}^n$ , each of dimension between 1 and  $n - 1$ , and let  $\mathcal{L}$  be closed under a  $\mathbf{Z}_q$ -action  $\nu$  on  $\mathbf{R}^n$ , whose homeomorphisms are isometries of  $\mathbf{R}^n$  fixing the origin. Suppose that the linear span of  $\cup \mathcal{L}$  is the whole  $\mathbf{R}^n$ . Then there is a subset  $Y \subset S^{n-1}$  of dimension at most  $n - 2$ , closed under the action  $\nu$ , and a  $\mathbf{Z}_q$ -map  $g: \mathbf{R}^n \setminus \cup \mathcal{L} \rightarrow Y$ .*

We postpone the proof to the end of this section. In the problem of  $(\frac{2}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})$ -partitioning of 2 measures by 4-fans, we thus obtain a  $\mathbf{Z}_5$ -mapping  $g \circ \tilde{f}_1: S^3 \rightarrow Y$ , where  $Y$  is a 2-dimensional space. This is ruled out by Dold's theorem 5.1.

A similar situation arises for  $(\frac{2}{3}, \frac{1}{3})$ -partitioning 3 measures by a 2-fan. Here we have two  $\mathbf{Z}_3$ -mappings  $f_1, f_2: X_3 \rightarrow Z$ , which can be regarded as a single  $\mathbf{Z}_3$ -map to the product  $Z \times Z = \{(y_1, y_2, \dots, y_6) \in \mathbf{R}^6 : y_1 + y_2 + y_3 = 0, y_4 + y_5 + y_6 = 0\}$ . This time, the excluded subspace is  $L(\frac{2}{3}, \frac{1}{3}) \times L(\frac{2}{3}, \frac{1}{3}) = \{y \in \mathbf{R}^6 : y_1 + y_2 = y_3 = y_4 + y_5 = y_6 = 0\}$ , plus its two images under the  $\mathbf{Z}_3$ -action  $\nu \times \nu$ . The sphere  $S(Z \times Z)$  is again an  $S^3$ , and 3 copies of  $S^1$  are deleted this time. By Lemma 6.1, the target space can be  $\mathbf{Z}_3$ -mapped to a 2-dimensional subspace, and an application of Lemma 5.2 and of Theorem 5.1 finishes the proof.

**Equipartitions of 3 measures by 2-fans.** Here we have the mappings  $f_1, f_2: X_2 \rightarrow Z$ , where the target space is 1-dimensional. By considering this as a single mapping into the product  $Z \times Z$  and by composing with the mapping

$h: Z \times Z \setminus \{0\} \rightarrow S(Z \times Z)$ , we obtain a mapping  $: X_2 \rightarrow S^1$ . Combining it with the identification  $\iota_2$  we get an  $f: V_2(\mathbf{R}^3) \rightarrow S^1$  map which is a  $\mathbf{Z}_2$ -map. In fact, it is straightforward to check that  $f$  is antipodal in the second variable, that is  $f(x, -y) = -f(x, y)$ .

We cannot use Lemma 5.2 since  $p$  and  $q$  are not coprime (both are equal to 2). But the second, almost elementary, proof of the 3-fan equipartition works. Namely,  $f$ , when restricted to  $S(a)$ , is an  $S^1 \rightarrow S^1$  antipodal map so it has odd degree. But when extending it to  $S^*$  by winding twice around  $S^1$  we get that this map is homotopic to zero, and its degree (which is twice that of  $f$ ) is zero. A contradiction again.

We remark that another proof can be obtained using the ideal-valued cohomological index of Fadell and Husseini (discussed in [Živ]).

**Proof of Lemma 6.1.** Define  $Y_0 \subset \mathbf{R}^n \setminus \bigcup \mathcal{L}$  as the set of all points  $y$  that have at least two nearest neighbors in  $\bigcup \mathcal{L}$  (thus,  $Y_0$  consists of the boundaries of the cells of the Voronoi diagram of  $\mathcal{L}$ ). This  $Y_0$  is a union of finitely many pieces of quadratic surfaces (the surfaces equidistant to some  $L_i, L_j \in \mathcal{L}$ ). We define a mapping  $g_0: \mathbf{R}^n \setminus \bigcup \mathcal{L} \rightarrow Y_0$ ; by composing  $g_0$  with the mapping  $h: \mathbf{R}^n \setminus \{0\} \rightarrow S^{n-1}$ , we then obtain the desired  $g$ .

For  $y \in Y_0$ , we let  $g_0(y) = y$ . For  $y \notin Y_0$ , let  $\bar{y}$  be the (unique) point of  $\bigcup \mathcal{L}$  nearest to  $y$ , and define the semiline  $s$  as the part of the line  $y\bar{y}$  starting at  $y$  and not containing  $\bar{y}$ .

We claim that  $s$  intersects  $Y_0$ . If  $L_i \in \mathcal{L}$  is the linear subspace containing  $\bar{y}$ , then  $s$  is perpendicular to  $L_i$ . If we travel distance  $t$  from  $y$  along the semiline  $s$ , the distance to  $L_i$  increases by  $t$ . On the other hand, since  $\mathcal{L}$  spans  $\mathbf{R}^n$ , not all of its spaces can be perpendicular to  $s$ , and thus there is a  $L_j \in \mathcal{L}$  such that by traveling distance  $t$  from  $y$  along  $s$ , the distance to  $L_j$  increases by no more than  $\beta t$  for a fixed  $\beta < 1$ . Therefore, by going far enough along  $s$ , we reach a point that is closer to  $L_j$  than to  $L_i$ , and we must have passed a point of  $Y_0$  on the way. This proves the claim, and we define the first intersection of  $s$  with  $Y_0$  as  $g_0(y)$ .

Since  $g_0$  is defined using metric properties and the action  $\nu$  is an isometry,  $g_0$  commutes with  $\nu$ . It remains to check the continuity of  $g_0$ . First, let  $y \in Y_0$ ; we have  $g_0(y) = y$  and so want to show that points close to  $y$  are mapped close to  $y$ . Let  $\mathcal{L}' \subseteq \mathcal{L}$  be the collection of the linear subspaces containing points nearest to  $y$ . For  $L_i \in \mathcal{L}'$ , let  $\bar{y}_i \in L_i$  be the point of  $L_i$  nearest to  $y$ . First, we note that  $y\bar{y}_i$  cannot be perpendicular to all  $L_j \in \mathcal{L}'$ . Indeed, if it were the case,  $\mathcal{L}'$  would be contained in the hyperplane  $h$  perpendicular to  $y\bar{y}_i$  and containing  $\bar{y}_i$ , and  $\bar{y}_i$  is the unique point of  $h$  nearest to  $y$ , which contradicts  $y$  having at least two nearest neighbors in  $\bigcup \mathcal{L}$ . Now if we choose a point  $y'$  in a sufficiently small  $\delta$ -neighborhood of  $y$ , the nearest point  $\bar{y}' \in \bigcup \mathcal{L}$  used in the definition of  $g_0(y')$  lies in some  $L_i \in \mathcal{L}'$ . Moreover, there is an  $L_j \in \mathcal{L}'$  such that the line  $y'\bar{y}'$  is not perpendicular to it, and so by traversing distance  $t$  along the line  $y'\bar{y}'$ , the

distance to  $L_j$  changes by at most  $\beta t$  for some  $\beta < 1$  (independent of  $\delta$ ). Since the distances of  $y'$  to  $L_i$  and to  $L_j$  differ by no more than  $2\delta$ , we see that  $g_0(y')$  lies at distance at most  $2\delta/(1-\beta)$  from  $y'$ . This proves the continuity of  $g_0$  at  $y$ .

Finally, we consider  $y \notin Y_0$ , and let  $y'$  lie very close to  $y$ . We let  $\bar{y}'$  be the point of  $\bigcup \mathcal{L}$  nearest to  $y'$ , and we let  $y''$  be the point of the line  $y'\bar{y}'$  closest to  $g_0(y)$ . By choosing  $y'$  sufficiently near to  $y$ , we can guarantee that  $y''$  and  $g_0(y)$  are arbitrarily close. Then we can apply the continuity of  $g_0$  at  $g_0(y)$ .  $\square$

## 7 Discussion and open problems

The most natural open question is probably whether, for two measures, a four-fan equipartition exists. The reasons why our topological approach cannot provide a positive answer will be briefly discussed below.

Our results show that  $\alpha$ -partitions of two measures by 3-fans and by 4-fans are possible for *some* values of  $\alpha$ . A very intriguing question is whether they exist for *all* values of  $\alpha$ . It seems hard to imagine why some  $\alpha$ 's shouldn't work when others do. On the other hand, our proof method cannot provide many more values of  $\alpha$ ; the reasons are indicated below. Maybe equivariant topology is not the right tool here.

**Equipartitions by 4-fans.** The space  $X_4$  of 4-fans is naturally equipped by the  $\mathbf{Z}_4$ -action  $\omega$ , but the corresponding coordinate shift action  $\nu$  of the target sphere  $S(Z)$  is not free. There are some theorems excluding the existence of an equivariant map even for non-free action, discovered by Özaydin, by Sarkaria (unpublished manuscripts), and by Volovikov [Vol96] (see [Živ] for discussion). These theorems involve actions of non-cyclic groups like  $\mathbf{Z}_p \times \mathbf{Z}_p$ . In our case, there are natural  $\mathbf{Z}_2 \times \mathbf{Z}_2$ -actions on the spaces involved.

On  $X_q$ , we have the free  $\mathbf{Z}_2$ -action  $\omega'$  corresponding to “changing the orientation” of a  $q$ -fan. Namely,  $\omega'(x; \ell_1, \dots, \ell_q) = (-x; \ell_1, \ell_q, \ell_{q-1}, \dots, \ell_2)$ . We have  $f_j(\omega'(F)) = \nu'(y_1, \dots, y_q) = (y_q, y_{q-1}, \dots, y_2, y_1)$ , and so the mappings  $f_j$  are also equivariant with respect to  $\omega'$  and the action  $\nu'$  given by reversing the order of coordinates.

For  $q = 4$ , we can take the  $\mathbf{Z}_2$ -action  $\omega^2$  of  $X_4$  (turning by two sectors) and combine it with the  $\omega'$  defined above. As is easy to check, we obtain a free action of the group  $G = \mathbf{Z}_2 \times \mathbf{Z}_2$  (direct product) on  $X_4$ , and  $f_1$  becomes a  $G$ -map. Unfortunately, we found that there is a  $G$ -map  $X_4 \rightarrow S(Z)$ . While this provides no counterexample to the 4-fan equipartitioning problem, it shows that this particular proof method fails.

Here comes an explicit description of such a  $G$ -map. Recall that  $X_4$  is just  $V_2(\mathbb{R}^3) = \{(x, y) \in S^2 \times S^2 \mid \langle x, y \rangle = 0\}$ . The actions of the generators of  $G$  are:

$$\omega^2(x, y) = (x, -y), \quad \omega'(x, y) = (-x, y).$$

In this case  $S(Z)$  is a  $S^2$  and the actions corresponding to  $\omega^2$ , resp.  $\omega'$  on the target space are  $g$  and  $h$  with

$$g(u, v, w) = (-u, -v, w), \quad h(u, v, w) = (-u, v, -w).$$

With  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$  we define  $f: V_2(\mathbf{R}^3) \rightarrow S^2$  as

$$f(x, y) = (|x_1|a - x_2b, |x_1|b + x_2a, x_3)$$

where  $a = (x_1y_2 - x_2y_1)/\sqrt{x_1^2 + x_2^2}$  and  $b = \text{sgn}(y_3)\sqrt{1 - a^2}$ . The expression for  $f(x, y)$  is undefined if  $x_3 = \pm 1$  but one can check that for all  $y$ ,  $f(x, y) \rightarrow (0, 0, \pm 1)$  as  $x_3 \rightarrow \pm 1$ , respectively, and so  $f$  extends continuously to these points. An easy calculation verifies that  $\|f(x, y)\| = 1$  for all unit  $x, y$ . Finally,  $f$  is continuous at  $y_3 = 0$  too since, as further simple calculation reveals,  $(x_1^2 + x_2^2)(1 - a^2) = 0$  whenever  $y_3 = 0$ . And clearly,  $f(x, y)_1$  is odd in both  $x$  and  $y$  and  $f(x, y)_2$  is even in  $x$  and odd in  $y$ .

**Equivariant maps for larger  $q$ .** In the above proofs, the existence of equivariant maps could be excluded using Dold's theorem or other tools. Unfortunately, it turns out that for sufficiently large  $q$ , equivariant maps between the considered  $\mathbf{Z}_q$ -spaces do exist, although it is not clear if such maps can actually arise in the  $k$ -fan partition problem. We describe one concrete example concerning the  $\alpha$ -partitioning of 3 measures by 2-fans, where the presentation of the mapping is particularly simple. Similar examples can be given for partitioning of 2 measures by 3-fans and 4-fans.

Let  $\alpha = (\frac{1}{q}, 1 - \frac{1}{q})$ ,  $q \geq 3$  odd. Recall that for  $\alpha$ -partitioning of 3 measures by a 2-fan, the test mappings provide a  $\mathbf{Z}_q$ -map  $S^3 \rightarrow Z \times Z$ , where  $Z$  is the hyperplane  $\{x_1 + x_2 + \dots + x_q = 0\}$  in  $\mathbf{R}^q$  and where the  $\mathbf{Z}_q$ -action  $\nu$  on  $\mathbf{R}^q$  is the coordinate shift. The nonexistence of the  $\alpha$ -partition implies that this mapping avoids the subspace  $L(\frac{1}{q}, 1 - \frac{1}{q}) \times L(\frac{1}{q}, 1 - \frac{1}{q}) = \{(x, y) : x \in Z, y \in Z, x_1 = 0, y_1 = 0\}$ . The following example shows that for  $q \geq 6$ , an equivariant map exists in this situation.

**Proposition 7.1** *For any integer  $q \geq 6$ , there exists a  $\mathbf{Z}_q$ -map  $f: S^3 \rightarrow \mathbf{R}^q \times \mathbf{R}^q$  (where  $S^3$  has a free  $\mathbf{Z}_q$ -action and both the  $\mathbf{R}^q$ s in the target space are equipped with the coordinate shift action), such that for all  $x \in S^3$ ,  $\sum_{i=1}^q f(x)_{1i} = \sum_{i=1}^q f(x)_{2i} = 0$  and  $f(x)_{11}$  and  $f(x)_{21}$  are never simultaneously 0 (the points of the image are indexed as  $2 \times q$  matrices).*

**Proof.** For simplicity of notation, let  $q = 6$  (the construction for larger  $q$  is entirely analogous; let us stress that the primality or non-primality of  $q$  plays no role here). It is sufficient to construct a continuous map  $g: S^3 \rightarrow \mathbf{R}^2 \setminus \{0\}$  satisfying

$$\sum_{i=0}^5 g(\omega^i x) = 0 \tag{2}$$

for all  $x \in S^3$ , where  $\omega$  is a free  $\mathbf{Z}_q$ -action on  $S^3$ . Indeed, with such a  $g$  at our disposal, the mapping  $f$  given by  $f(x)_{ji} = g(\omega^i x)_j$  is as required.

Let  $A$  be the perimeter of a regular hexagon  $a_0 a_1 \dots a_5$ ; that is,  $A$  is the 1-dimensional simplicial complex with the 6 vertices  $a_0, a_1, \dots, a_5$  and with 1-simplices  $a_i a_{i+1}$ ,  $i = 0, 1, \dots, 5$  (indices taken modulo 6). Similarly,  $B$  is the perimeter of a hexagon  $b_0 b_1 \dots b_5$ . The sphere  $S^3$  is homeomorphic to the join  $A * B$ . The 3-dimensional simplices of this join have the form  $a_i a_{i+1} b_j b_{j+1}$ . A  $\mathbf{Z}_q$ -action  $\omega$  is defined on  $A * B$  by letting  $a_i \mapsto a_{i+1}$ ,  $b_i \mapsto b_{i+1}$ , and extending linearly on each simplex.

First we describe a simpler mapping  $g_0: A * B \rightarrow \mathbf{R}^2$ . Let us choose three nonzero vectors  $u, v, w \in \mathbf{R}^2$  with  $4u + v + w = 0$ . Put  $g_0(a_0) = g_0(b_0) = v$ ,  $g_0(a_1) = g_0(b_1) = w$ , and  $g_0(a_i) = g_0(b_i) = u$  for  $i = 2, 3, 4, 5$ . This defines  $g_0$  on the vertices of  $A * B$ , and we extend it linearly on each simplex of  $A * B$ . Clearly, (2) is satisfied for  $g_0$ , but there are simplices in  $A * B$  whose image contains 0. We note that all such simplices contain  $a_0$  or  $b_0$ , and also  $a_1$  or  $b_1$ .

Next, we form a simplicial complex  $K$  by subdividing some simplices in  $A * B$ . We let  $c_{01}$  be the midpoint of the edge  $a_0 b_1$ , and we let  $O_1 = \{c_{01}, c_{12}, \dots, c_{50}\}$  be the orbit of  $c_{01}$  under  $\omega$ . Similarly,  $c_{10}$  is the midpoint of the edge  $a_1 b_0$  and  $O_2 = \{c_{10}, c_{21}, \dots, c_{05}\}$  is its orbit; see the schematic Fig. 7. Each edge of




Figure 7: The subdivided edges in  $A * B$ .

$A * B$  containing a point of  $O_1 \cup O_2$  is subdivided, and the higher-dimensional simplices containing such edges are appropriately subdivided too. In particular, 3-dimensional simplices may have one subdivided edge (such as the simplex  $a_0 a_1 b_2 b_3$ ) or two (such as  $a_0 a_1 b_0 b_1$  or  $a_0 a_1 b_1 b_2$ ). Those with one subdivided edge are subdivided into two simplices as in Fig. 8(a), and those with two subdivided edges into four simplices as in Fig. 8(b) (note that the subdivided edges are never adjacent). This defines the simplicial complex  $K$ . Now we define a new mapping  $g$  on the vertex set of  $K$ ; the values at the  $a_i$  and  $b_j$  are as those for  $g_0$ , and we put  $g(c_{23}) = g(c_{32}) = w$ ,  $g(c_{45}) = g(c_{54}) = v$ , and  $g(c_{ij}) = u$  for all other  $c_{ij} \in O_1 \cup O_2$ ; see Fig. 9 (the vertex labels are as in Fig. 7 and are omitted). Finally,  $g$  is extended linearly on the whole polyhedron of  $K$ . It is easy to check




Figure 8: Subdividing the 3-dimensional simplices.




Figure 9: The values of the mapping  $g$ ; unlabeled vertices receive  $u$ .

that for each simplex of  $K$ ,  $g$  attains at most two distinct values at the vertices. Thus, no point of the polyhedron of  $K$  is mapped to 0 by  $g$ .

*Remark.* Another, somewhat more complicated example of this type was constructed independently by Attila Pór.

## Acknowledgment

We would like to thank Attila Pór, Otfried Cheong, András Szűcs, Gyula Y. Katona and Pavel Valtr for inspiring discussions and valuable remarks concerning the subject of this paper. We also thank Rade Živaljević for his kind explanations concerning the ideal-valued index and the existence of equivariant mappings.

## References

[ARNU98] J. Akiyama, E. Rivera-Campo, G. Nakamura, and J. Urrutia. Perfect divisions of a cake. Manuscript, Tokai University, 1998.

- [BSS81] I. Bárány, S. Shlosman, and A. Szücs. On a topological generalization of a theorem of Tverberg. *J. London Math. Soc.* 23 (1981), 158–164.
- [BKS98] S. Bespamyatnikh, D. Kirkpatrick, and J. Snoeyink. Generalizing ham sandwich cuts to equitable subdivisions. Manuscript, Univ. of British Columbia, 1998.
- [Bre93] G. Bredon. *Topology and Geometry (Graduate Texts in Mathematics 139)*. Springer-Verlag, Berlin etc., 1993.
- [Dol83] A. Dold. Simple proofs of some Borsuk-Ulam results. In *Contemp. Math.* 19, volume 68, pages 64–69. Amer. Math. Soc., 1983.
- [IUY98] H. Ito, H. Uehara, and M. Yokoyama. 2–dimension ham-sandwich theorem for partitioning into three convex pieces. Manuscript, Toyohashi University of Technology, 1998.
- [KK98] A. Kaneko and M. Kano. Balanced partition of point sets in the plane. Manuscript, 1998.
- [KZ75] M. A. Krasnoselsky and P. P. Zabrejko. Geometric problems of non-linear analysis. Nauka, Moscow, 1975 (in Russian).
- [Mak88] V. V. Makeev. Six-lobed partitions of three-dimensional space. *Vestn. Leningr. Univ.*, Ser. I 1988, No.2, 31-34 (1988) (in Russian; English translation in *Vestn. Leningr. Univ.*, Math. 21, No.2, 40-45 (1988)).
- [Ram96] E. A. Ramos. Equipartition of mass distributions by hyperplanes. *Discrete Comput. Geom.*, 15:147–167, 1996.
- [Sa98] T. Sakai. Radial partitions of point sets in  $\mathbf{R}^2$ . Manuscript, Tokoha Gakuen University, 1998.
- [Sch93] L. J. Schulman. An equipartition of planar sets. *Discrete Comput. Geom.*, 9:257–266, 1993.
- [VC71] V. N. Vapnik and A. Ya. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory Probab. Appl.*, 16:264–280, 1971.
- [Vol96] A. Yu. Volovikov. On a topological generalization of the Tverberg theorem. *Math. Notes*, 59(3):324–326, 1996. Translation from *Mat. Zametki* 59, No.3, 454-456 (1996).
- [Živ] R. T. Živaljević. User’s guide to equivariant methods in combinatorics II. *Publ. Inst. Math. Beograd*. To appear.

- [Živ96] R. T. Živaljević. User's guide to equivariant methods in combinatorics. *Publ. Inst. Math. Beograd*, 59(73):114–130, 1996.
- [Živ97] R. T. Živaljević. Topological methods. In Jacob E. Goodman and Joseph O'Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 11, pages 209–224. CRC Press LLC, Boca Raton, FL, 1997.