

A stronger form of the theorem constructing a rigid binary relation on any set

Apoloniusz Tyszk

Summary. On every set A there is a rigid binary relation i.e. such a relation $\mathcal{R} \subseteq A \times A$ that there is no homomorphism $\langle A, \mathcal{R} \rangle \rightarrow \langle A, \mathcal{R} \rangle$ except the identity (Vopěnka et al. [1965]). We prove that for each infinite cardinal number κ if $\text{card } A \leq 2^\kappa$, then there exists a relation $\mathcal{R} \subseteq A \times A$ with the following property:

$$\forall x \in A \exists \left\{ \begin{array}{l} \{x\} \subseteq A(x) \subseteq A \\ \text{card } A(x) \leq \kappa \end{array} \right. \forall f : A(x) \rightarrow A \quad f \text{ is not a homomorphism of } \mathcal{R} \\ f \neq \text{id}_{A(x)}$$

which implies that \mathcal{R} is rigid. If a relation $\mathcal{R} \subseteq A \times A$ has the above property, then $\text{card } A \leq 2^\kappa$.

On every set A there is a rigid binary relation, i.e. such a relation $\mathcal{R} \subseteq A \times A$ that there is no homomorphism $\langle A, \mathcal{R} \rangle \rightarrow \langle A, \mathcal{R} \rangle$ except the identity ([8] and also [2],[3] [5]). Conjectures 1 and 2 below strengthen this theorem.

Mathematics Subject Classification (2000): 03E05, 08A35.

Keywords and phrases: rigid binary relation, rigid symmetric relation.

Conjecture 1 ([6],[7]). If κ is an infinite cardinal number and $\text{card } A \leq 2^{2^\kappa}$, then there exists a relation $\mathcal{R} \subseteq A \times A$ which satisfies the following condition:

$$(\kappa^*) \quad \forall x, y \in A \quad \exists \{x\} \subseteq A(x, y) \subseteq A \quad \forall f : A(x, y) \rightarrow A \\ x \neq y \quad \text{card } A(x, y) \leq \kappa \quad f(x) = y$$

f is not a homomorphism of \mathcal{R} .

Proposition 1a ([7]). If κ is an infinite cardinal number, $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^*) and $\text{card } \tilde{A} \leq \text{card } A$, then there exists a relation $\tilde{\mathcal{R}} \subseteq \tilde{A} \times \tilde{A}$ which satisfies condition (κ^*) .

Proposition 1b ([6]). If $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^*) , then \mathcal{R} is rigid. If κ is an infinite cardinal number and a relation $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^*) , then $\text{card } A \leq 2^{2^\kappa}$.

Theorem 1 ([7]). Conjecture 1 is valid for $\kappa = \omega$.

Conjecture 2 ([6],[7]). If $\kappa \neq 0$ is a limit cardinal number and $\text{card } A \leq 2^{\sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}}$, then there exists a relation $\mathcal{R} \subseteq A \times A$ which satisfies the following condition:

$$(\kappa^{**}) \quad \forall x, y \in A \quad \exists \{x\} \subseteq A(x, y) \subseteq A \quad \forall f : A(x, y) \rightarrow A \\ x \neq y \quad \text{card } A(x, y) < \kappa \quad f(x) = y$$

f is not a homomorphism of \mathcal{R} .

Proposition 2a ([7]). If $\kappa \neq 0$ is a limit cardinal number, $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^{**}) and $\text{card } \tilde{A} \leq \text{card } A$, then there exists a relation $\tilde{\mathcal{R}} \subseteq \tilde{A} \times \tilde{A}$ which satisfies condition (κ^{**}) .

Proposition 2b ([6]). If $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^{**}) , then \mathcal{R} is rigid. If $\kappa \neq 0$ is a limit cardinal number and a relation $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^{**}) , then $\text{card } A \leq 2^{\sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}}$.

Theorem 2 ([6],[7]). Conjecture 2 is valid for $\kappa = \omega$.

In this article we prove a changed form of Conjecture 1 which holds for all infinite cardinal numbers κ , see Theorem 3, Corollary and Theorem 5.

Theorem 3. If κ is an infinite cardinal number and $\text{card } A \leq 2^\kappa$, then there exists a relation $\mathcal{R} \subseteq A \times A$ which satisfies the following condition:

$$(\kappa^\diamond) \quad \forall x \in A \quad \exists \{x\} \subseteq A(x) \subseteq A \quad \forall f : A(x) \rightarrow A \\ \text{card } A(x) \leq \kappa \quad f \neq \text{id}_{A(x)}$$

f is not a homomorphism of \mathcal{R} .

Proof. It is known ([1] and also [2],[5]) that for each infinite cardinal number κ there exists a rigid symmetric relation $R \subseteq \kappa \times \kappa$. Let Φ denote the family of all relations $S \subseteq \kappa \times \kappa$ which satisfy:

- (1) $R \subseteq S$,
- (2) for each $\alpha, \beta \in \kappa$ if $\alpha \neq \beta$, then $\alpha S \beta$ or $\beta S \alpha$,
- (3) for each $\alpha, \beta \in \kappa$ if $\alpha S \beta$ and $\beta S \alpha$, then $\alpha R \beta$ and $\beta R \alpha$.

Since R is rigid

- (4) $R \subseteq \{(\alpha, \beta) : \alpha, \beta \in \kappa, \alpha \neq \beta\}$.

By (1) and (3) the following Lemma 1 holds true.

Lemma 1. If $S_1, S_2 \in \Phi$ and $f : \langle \kappa, S_1 \rangle \rightarrow \langle \kappa, S_2 \rangle$ is a homomorphism, then $f : \langle \kappa, R \rangle \rightarrow \langle \kappa, R \rangle$ is a homomorphism.

Lemma 2. For every $S_1, S_2 \in \Phi$ if $S_1 \neq S_2$, then $\text{id}_\kappa : \langle \kappa, S_1 \rangle \rightarrow \langle \kappa, S_2 \rangle$ is not a homomorphism.

Proof. Applying (3) and (4) we obtain two cases. First case: there exist $\alpha, \beta \in \kappa$, $\alpha \neq \beta$ such that $(\alpha, \beta) \in S_1$ and $(\alpha, \beta) \notin S_2$, so id_κ is not a homomorphism. Second case: there exist $\alpha, \beta \in \kappa$, $\alpha \neq \beta$ such that $(\alpha, \beta) \in S_2$ and $(\alpha, \beta) \notin S_1$. By (2) $(\beta, \alpha) \in S_1$. It suffices to prove that $(\beta, \alpha) \notin S_2$. Suppose, on the contrary, that $(\beta, \alpha) \in S_2$. By (3) $(\alpha, \beta) \in R$, so by (1) $(\alpha, \beta) \in S_1$, a contradiction.

Lemma 3. $\text{card } \Phi = 2^\kappa$.

Proof. Let $T := \{(\alpha, \beta) : \alpha, \beta \in \kappa, \alpha \neq \beta, (\alpha, \beta) \notin R\}$. It suffices to prove that $\text{card } T = \kappa$. Suppose, on the contrary, that $\text{card } T < \kappa$. Hence $\text{card } \bigcup T < \kappa$ and consequently $\text{card } (\kappa \setminus \bigcup T) = \kappa$. For each $\alpha, \beta \in \kappa \setminus \bigcup T$ if $\alpha \neq \beta$, then $(\alpha, \beta) \in R$. From this and (4) any non-identical injection from κ into $\kappa \setminus \bigcup T$ is a homomorphism of R . This contradiction completes the proof of Lemma 3.

Now we turn to the main part of the proof. For each $\emptyset \neq \Psi \subseteq \Phi$ we define the relation $R_\Psi \subseteq (\kappa \times \Psi) \times (\kappa \times \Psi)$ by the following formula:

$$\forall \alpha, \beta \in \kappa \forall S_1, S_2 \in \Psi \left(((\alpha, S_1), (\beta, S_2)) \in R_\Psi \iff (\alpha, \beta) \in S_1 = S_2 \right).$$

In other words, the graph corresponding to the relation R_Ψ is a disjoint union of graphs belonging to Ψ . By Lemma 3 it suffices to prove that R_Ψ satisfies condition (κ^\diamond) . Let $(\lambda, S_1) \in \kappa \times \Psi$. We prove that $(\kappa \times \Psi)((\lambda, S_1)) := \kappa \times \{S_1\}$ satisfies condition (κ^\diamond) .

Suppose, on the contrary, that $f : \kappa \times \{S_1\} \rightarrow \kappa \times \Psi$ is a homomorphism of R_Ψ and $f \neq \text{id}_{\kappa \times \{S_1\}}$. Then there exist $\alpha, \beta \in \kappa$ and

$S_2 \in \Psi$ such that $f((\alpha, S_1)) = (\beta, S_2)$ and $(\alpha, S_1) \neq (\beta, S_2)$. By (2) for each $\gamma \in \kappa \setminus \{\alpha\}$ $\alpha S_1 \gamma$ or $\gamma S_1 \alpha$. From this for each $\gamma \in \kappa \setminus \{\alpha\}$ $(\alpha, S_1) R_\Psi(\gamma, S_1)$ or $(\gamma, S_1) R_\Psi(\alpha, S_1)$. Therefore $f((\alpha, S_1)) R_\Psi f((\gamma, S_1))$ or $f((\gamma, S_1)) R_\Psi f((\alpha, S_1))$ and consequently $(\beta, S_2) R_\Psi f((\gamma, S_1))$ or $f((\gamma, S_1)) R_\Psi(\beta, S_2)$. In both cases there exists a $\delta \in \kappa$ such that $f((\gamma, S_1)) = (\delta, S_2)$. It implies that f maps $\kappa \times \{S_1\}$ into $\kappa \times \{S_2\}$. Let $\pi : \{S_1\} \rightarrow \{S_2\}$. There is a uniquely determined transformation $\tilde{f} : \kappa \rightarrow \kappa$ such that $f = \langle \tilde{f}, \pi \rangle$. Obviously, $\tilde{f}(\alpha) = \beta$ and $\tilde{f} : \langle \kappa, S_1 \rangle \rightarrow \langle \kappa, S_2 \rangle$ is a homomorphism. By Lemma 1 $\tilde{f} : \langle \kappa, R \rangle \rightarrow \langle \kappa, R \rangle$ is a homomorphism. Since R is rigid $\tilde{f} = \text{id}_\kappa$. Therefore $\alpha = \tilde{f}(\alpha) = \beta$ and $\text{id}_\kappa : \langle \kappa, S_1 \rangle \rightarrow \langle \kappa, S_2 \rangle$ is a homomorphism. On the other hand, $\alpha = \beta$ and $(\alpha, S_1) \neq (\beta, S_2)$ implies $S_1 \neq S_2$. It is impossible by Lemma 2. This contradiction completes the proof of Theorem 3.

Remark 1. It is easy to observe that condition (κ^\diamond) implies condition (κ^*) . Obviously, if $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^\diamond) , then \mathcal{R} is rigid.

Theorem 3 follows easily from Perminov's ([4]) Theorem 4 on rigid tournaments. By a tournament on A we will understand a relation $Z \subseteq A \times A$ satisfying:

$$\begin{aligned} \forall \substack{x, y \in A \\ x \neq y} \quad & (x, y) \in Z \vee (y, x) \in Z, \\ \forall x, y \in A \quad & (x, y) \in Z \Rightarrow (y, x) \notin Z. \end{aligned}$$

Tournaments in [4] are defined differently, but Theorem 4 holds true for our definition. Unfortunately, the proof in [4] is only sketched.

Theorem 4 ([4]). For each infinite cardinal number κ there exists a family Γ of 2^κ rigid tournaments on κ such that there is no homomorphism between any two distinct tournaments belonging to Γ .

Corollary. For every $\Delta \subseteq \Gamma$ a disjoint union of tournaments belonging to Δ satisfies condition (κ^\diamond) . It gives a new proof of Theorem 3.

We will show an alternative, algebraic method for proving Theorem 3. This method described in the proof of Theorem 5 gives more, namely a symmetric relation satisfying condition (κ^\diamond) . Unfortunately, in contradiction to the relation constructed in the proof of Theorem 3, a direct description of such a relation is very complicated.

Theorem 5. If κ is an infinite cardinal number and $\text{card } A \leq 2^\kappa$, then

there exists a symmetric relation $\mathcal{R} \subseteq A \times A$ which satisfies condition (κ^\diamond) .

Proof. Following [5] let **Graph** denote the category of graphs and their homomorphisms. The objects of **Graph** are couples (X, R) with $R \subseteq X \times X$, the morphisms from (X, R) to (X', R') are triples $((X', R'), f, (X, R))$ with $f : X \rightarrow X'$ such that $(f(x), f(y)) \in R'$ whenever $(x, y) \in R$, and it is viewed as a concrete category endowed with the natural forgetful functor. Let \mathcal{C}_n ($n \geq 3$) denote the category of connected n -chromatic undirected graphs and their homomorphisms. It is known (see [5], Theorem 4.12 on page 113), that for every $n \geq 3$ there is a strong embedding $F : \mathbf{Graph} \rightarrow \mathcal{C}_n$ that transforms objects of the cardinality κ into objects of the cardinality κ .

Let $R \subseteq \kappa \times \kappa$ be a rigid symmetric relation (undirected graph). Considering all possible orientations of R we obtain 2^κ rigid graphs with the property that there is no homomorphism between any two distinct graphs. Using the strong embedding $F : \mathbf{Graph} \rightarrow \mathcal{C}_n$ ($n \geq 3$) constructed in [5] we obtain a family Γ of 2^κ rigid undirected graphs on κ with the property that there is no homomorphism between any two distinct graphs. Since Γ consists of connected graphs, for every $\Delta \subseteq \Gamma$ a disjoint union of graphs belonging to Δ satisfies condition (κ^\diamond) , the proof is similar to the proof that R_Ψ satisfies condition (κ^\diamond) . This completes the proof of Theorem 5.

Theorem 6. If κ is an infinite cardinal number and a relation $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^\diamond) , then $\text{card } A \leq 2^\kappa$.

Proof. Suppose, on the contrary, that $\mathcal{R} \subseteq A \times A$ satisfies condition (κ^\diamond) and $\text{card } A > 2^\kappa$. For each $x \in A$ we choose the set $A(x)$ from condition (κ^\diamond) . Let $\mathcal{B} := \{A(x) : x \in A\}$. Since $\bigcup \mathcal{B} = A$ we conclude that $\text{card } \mathcal{B} = \text{card } A$. For each $B \in \mathcal{B}$ we choose a bijective $f_B : \text{card } B \rightarrow B$ and define the relation $R_B \subseteq \text{card } B \times \text{card } B$ by the following formula:

$$\forall \alpha, \beta \in \text{card } B \quad ((\alpha, \beta) \in R_B \iff (f_B(\alpha), f_B(\beta)) \in \mathcal{R}).$$

Let

$$\mathcal{B} \ni B \xrightarrow{h} (\text{card } B, R_B) \in \bigcup_{\substack{\alpha \in \text{Card} \\ \alpha \leq \kappa}} \{\alpha\} \times \mathcal{P}(\alpha \times \alpha).$$

Since

$$\text{card } \mathcal{B} = \text{card } A > 2^\kappa = \text{card} \bigcup_{\substack{\alpha \in \text{Card} \\ \alpha \leq \kappa}} \{\alpha\} \times \mathcal{P}(\alpha \times \alpha)$$

we conclude that there exist $B_1, B_2 \in \mathcal{B}$ such that $B_1 \neq B_2$ and $h(B_1) = h(B_2)$. Hence $\langle B_1, \mathcal{R} \rangle \xrightarrow{f_{B_2} \circ (f_{B_1})^{-1}} \langle B_2, \mathcal{R} \rangle$ is a non-identical isomorphism. This contradiction completes the proof.

Conjecture 3. If $\kappa \neq 0$ is a limit cardinal number and $\text{card } A \leq \sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}$, then there exists a relation $\mathcal{R} \subseteq A \times A$ which satisfies the following condition:

$$(\kappa^{\diamond\diamond}) \quad \forall x \in A \quad \exists \begin{array}{l} \{x\} \subseteq A(x) \subseteq A \\ \text{card } A(x) < \kappa \end{array} \quad \forall f : A(x) \rightarrow A \\ f \neq \text{id}_{A(x)} \\ f \text{ is not a homomorphism of } \mathcal{R}.$$

Remark 2. Obviously, if $\mathcal{R} \subseteq A \times A$ satisfies condition $(\kappa^{\diamond\diamond})$, then \mathcal{R} is rigid. By Theorem 3 Conjecture 3 holds true if $\text{card } A < \sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}$.

Theorem 7. If $\kappa \neq 0$ is a limit cardinal number and a relation $\mathcal{R} \subseteq A \times A$ satisfies condition $(\kappa^{\diamond\diamond})$, then $\text{card } A \leq \sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}$.

Proof. Suppose, on the contrary, that $\mathcal{R} \subseteq A \times A$ satisfies condition $(\kappa^{\diamond\diamond})$ and $\text{card } A > \sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\}$. For each $x \in A$ we choose the set $A(x)$ from condition $(\kappa^{\diamond\diamond})$. Let $\mathcal{B} := \{A(x) : x \in A\}$. Since $\bigcup \mathcal{B} = A$ we conclude that $\text{card } \mathcal{B} = \text{card } A$. For each $B \in \mathcal{B}$ we choose a bijective $f_B : \text{card } B \rightarrow B$ and define the relation $R_B \subseteq \text{card } B \times \text{card } B$ by the following formula:

$$\forall \alpha, \beta \in \text{card } B \quad ((\alpha, \beta) \in R_B \iff (f_B(\alpha), f_B(\beta)) \in \mathcal{R}).$$

Let

$$\mathcal{B} \ni B \xrightarrow{h} (\text{card } B, R_B) \in \bigcup_{\substack{\alpha \in \text{Card} \\ \alpha < \kappa}} \{\alpha\} \times \mathcal{P}(\alpha \times \alpha).$$

Since

$$\text{card } \mathcal{B} = \text{card } A > \sup\{2^\alpha : \alpha \in \text{Card}, \alpha < \kappa\} = \text{card} \bigcup_{\substack{\alpha \in \text{Card} \\ \alpha < \kappa}} \{\alpha\} \times \mathcal{P}(\alpha \times \alpha)$$

we conclude that there exist $B_1, B_2 \in \mathcal{B}$ such that $B_1 \neq B_2$ and $h(B_1) = h(B_2)$. Hence $\langle B_1, \mathcal{R} \rangle \xrightarrow{f_{B_2} \circ (f_{B_1})^{-1}} \langle B_2, \mathcal{R} \rangle$ is a non-identical isomorphism. This contradiction completes the proof.

Proposition 3. By Theorems 3 and 7, if $\kappa \neq 0$ is a limit cardinal number, $\mathcal{R} \subseteq A \times A$ satisfies condition $(\kappa^{\diamond\diamond})$ and $\text{card } \tilde{A} \leq \text{card } A$, then there exists a relation $\tilde{\mathcal{R}} \subseteq \tilde{A} \times \tilde{A}$ which satisfies condition $(\kappa^{\diamond\diamond})$.

Theorem 8. Conjecture 3 is valid for $\kappa = \omega$ i.e. there exists a relation $\mathcal{R} \subseteq \omega \times \omega$ satisfying condition $(\omega^{\diamond\diamond})$.

Proof. The relation $\mathcal{R} := \{(i, i+1) : i \in \omega\} \cup \{(0, 2)\} \subseteq \omega \times \omega$ satisfies condition $(\omega^{\diamond\diamond})$. Indeed, for each $i \in \omega$ the set $A(i) := \{j \in \omega : j \leq i+2\}$ is adequate for property $(\omega^{\diamond\diamond})$.

References

- [1] Z. Hedrlín and A. Pultr, *Symmetric relations (undirected graphs) with given semigroups*, Monatsh. Math. 69 (1965), 318–322.
- [2] J. Nešetřil, *Aspects of structural combinatorics (Graph homomorphisms and their use)*, Taiwanese J. Math. 3 (1999), 381–423.
- [3] J. Nešetřil, *A rigid relation for every set*, J. Graph Theory, to appear.
- [4] E. A. Perminov, *On the number of pairwise nonembeddable rigid graphs*, Sov. Math. 29 (1985), No. 8, 107–110; translation from Izv. Vyssh. Uchebn. Zaved. Mat. 1985, No. 8 (279), 77–78.
- [5] A. Pultr and V. Trnková, *Combinatorial, algebraic and topological representations of groups, semigroups and categories*, North-Holland, Amsterdam, 1980.
- [6] A. Tyszka, *Some remarks on endomorphism rigid relations*, Univ. Iagel. Acta Math. 31 (1994), 7–20.
- [7] A. Tyszka, *On binary relations without non-identical endomorphisms*, Aequationes Math., to appear.
- [8] P. Vopěnka, A. Pultr and Z. Hedrlín, *A rigid relation exists on any set*, Comment. Math. Univ. Carolinae 6 (1965), 149–155.

Apoloniusz Tyszka
 Technical Faculty
 Hugo Kołłątaj University
 Balicka 104, 30-149 Kraków, Poland
 E-mail: rtyyszka@cyf-kr.edu.pl