Triangles in random graphs

Martin Loebl, Jiří Matoušek, and Ondřej Pangrác *

Abstract

We show the number of triangles of $G_{n,\frac{1}{2}}$ is almost uniformly distributed among residue classes modulo q, where q is a prime number bounded by $\Theta(\log n)$. This implies a consequence of a conjecture of Bollobás, Pebody and Riordan (that almost every random graph $G_{n,\frac{1}{2}}$ is uniquely determined by its Tutte polynomial): almost every pair of independently chosen random graphs $G_{n,\frac{1}{2}}$ has different Tutte polynomials.

1 Introduction

The Tutte polynomial of a graph $G = (V, E)$ is the bivariate polynomial

$$T(G; x, y) = \sum_{A \subseteq E} (x-1)^{r(E)-r(A)} (y-1)^{|A|-r(A)},$$

where $r(A) = |V| - k(A)$ and $k(A)$ denotes the number of components of the graph (V, A).

It is clear that two isomorphic graphs have the same Tutte polynomial. On the other hand, there are examples of non-isomorphic graphs with the same Tutte polynomial (see [3, 1]). In this context, the question of how much information about the graph its Tutte polynomial contains is natural.

A random graph $G_{n,\frac{1}{2}}$, for n natural number, is a graph on an n element vertex set such that any pair of vertices forms an edge with probability $\frac{1}{2}$ independently of other pairs. Let $\mathcal{G}_{n,\frac{1}{2}}$ denote the corresponding probability

*Department of Applied Mathematics and Institute for Theoretical Computer Science Charles University, Malostranské nám. 25, 118 00 Prague, Czech Republic. Institute for Theoretical Computer Science is supported by Ministry of Education of Czech Republic as project LN00A056.

space. Note that such a random graph dense; the expectation of the number of edges is $\frac{1}{2}\binom{n}{2}$ and the expectation of the number of triangles is $\frac{1}{8}\binom{n}{3}$. The following conjecture was presented in [1].

Conjecture 1.1 *Almost every graph $G \in \mathcal{G}_{n,\frac{1}{2}}$ is such that $T(G;x,y) = T(G';x,y)$ implies $G \cong G'$.*

The main result of this paper stated below implies a weaker result.

Theorem 1.2 *There are numbers q_0 and n_0 and a function $f \in \Theta(\log n)$ such that for any $n > n_0$, any prime number q, $q_0 < q < f(n)$, and any choice of $k \in \{0,1,\ldots,q-1\}$ the probability that a random graph $G_{n,\frac{1}{2}}$ contains k triangles modulo q is $q^{-1}(1+o(1))$.*

Since the number of triangles of any simple graph can be easily derived from its Tutte polynomial, we can state this easy consequence of Theorem 1.2.

Corollary 1.3 *The probability that two independently chosen random graphs $G_{n,\frac{1}{2}}$ have the same Tutte polynomial is of order $O(1/\log n)$.*

The another open problem is to find or to prove an existence of a family containing almost every graph such that the Tutte polynomial distinguish any pair of graphs inside this family. Solving this problem affirmatively may give us better understanding and a hope for positive answer to the Conjecture 1.1.

2 Counting 2-paths

Let G be a random graph from $\mathcal{G}_{n,\frac{1}{2}}$ with the vertex set V. Let V_1 be a set of $\lfloor \frac{n}{4} \rfloor$ pairs of vertices $V_1 = \{u_1, v_1, u_2, v_2, \ldots, u_{\lfloor \frac{n}{4} \rfloor}, v_{\lfloor \frac{n}{4} \rfloor}\}$ (suppose n large enough) and let V_2 be a subset of $V - V_1$. Denote the size of V_2 by l. Fix a pair u_i, v_i of vertices of V_1. For $w \in V_2$, let X_w be an indicator of the existence of both edges $\{u_i, w\}$ and $\{v_i, w\}$. Then $X = \sum_{w \in V_2} X_w$ counts the number of 2-paths connecting vertices $\{u, v\}$ and having the middle vertex in V_2. Note that the probability of such a path through a fixed vertex w is $\left(\frac{1}{2}\right)^2$. Denote by $p_j(l)$ the probability that $X = j$. Since edges from distinct vertices of V_1 going to V_2 are independent, the number

of such 2-paths is independent for distinct pairs of vertices of V_1, and $p_j(l)$ is also independent of the exact choice of V_2.

Assume q to be an arbitrary fixed integer greater than 1 and smaller than n. Denote by $s_i(l) = \sum_{j \equiv i (\mathrm{mod}\ q)} p_j(l)$ the probability that the number of connecting 2-path is equal to i modulo q.

Lemma 2.1 *Let $1 \leq q < n$. There exists a number $c = c(q) > 0$ such that $s_i(l) \geq c$ for $i = 0, 1, \ldots, q-1$ and $l \geq q-1$.*

Proof: Using the recursion $p_i(l) = \frac{1}{4} p_{i-1}(l-1) + \frac{3}{4} p_i(l-1)$ one can derive $s_i(l) = \frac{1}{4} s_{i-1}(l-1) + \frac{3}{4} s_i(l-1)$ (where $s_{-1}(l)$ means $s_{q-1}(l)$).

It is easy to observe that if $s_i(l_0) \geq c$ for all possible i's and some l_0 and c, then also $s_i(l) \geq c$ for all $l \geq l_0$. Note that $s_i(q-1) = p_i(q-1)$, and clearly, each $p_i(q-1) \geq \left(\frac{1}{4}\right)^{q-1}$. Hence for any $l \geq q-1$ and any $i = 1, 2, \ldots, q-1$ we have $s_i(l) \geq \left(\frac{1}{4}\right)^{q-1}$. By setting $c(q) = \left(\frac{1}{4}\right)^{q-1}$ we complete the proof.

□

Lemma 2.2 *Let n be sufficiently large natural number. Then with probability at least $1 - \frac{1}{15} \cdot \frac{\log_2 n}{n^{1/3}}$ a graph $G \in \mathcal{G}_{n,\frac{1}{2}}$ has, for any $q \leq q_0 = \lfloor \log_2 n^{1/3} \rfloor$, disjoint pairs u_k, v_k of vertices of G ($k = 1, 2, \ldots, q-1$) where the number of 2-paths connecting u_k and v_k is k modulo q.*

Proof: Let V_1 be the set of $\lfloor \frac{n}{4} \rfloor$ pair of vertices as described above and let $V_2 = V - V_1$. We want to chose pairs u_k, v_k from the pairs of vertices forming V_1. By Lemma 2.1 the probability that the number of 2-paths connecting u_i and v_i is equal to k modulo q is at least $c(q)$ for any $k = 0, 1, \ldots, q-1$ (and hence at most $1-(q-1)c(q)$). Moreover, these probabilities are independent for distinct pairs u_i, v_i and u_j, v_j.

Denote the event that there is a pair u_i, v_i (among the pairs from V_1) connected by exactly k 2-paths (counted modulo q) by Y_k ($Y_k = 1$ if such a pair exists, $Y_k = 0$ otherwise). Then the probability $P[Y_1, Y_2, \ldots, Y_{q-1}]$ is equal to $1 - P\left[\exists k : \overline{Y_k}\right]$. Assume the edges inside V_1 are fixed. The probability that vertices u_i, v_i are connected by k 2-paths is $s_{i'}(|V_2|)$, where i' is such that the number of 2-paths with the middle vertex inside V_1 plus i' is equal to k modulo q. Hence $P\left[\overline{Y_k}\right] = \prod_{i=1}^{\lfloor \frac{n}{4} \rfloor} (1 - s_{i'}(|V_2|)) \leq (1-c(q))^{\lfloor \frac{n}{4} \rfloor}$.

$$P\left[\exists k : \overline{Y_k}\right] \leq \sum_{k=1}^{q-1} P\left[\overline{Y_k}\right] \leq \sum_{k=1}^{q-1} (1-c(q))^{\lfloor \frac{n}{4} \rfloor} = (q-1)(1-c(q))^{\lfloor \frac{n}{4} \rfloor}.$$

Let $q_0 = \lfloor \log_2 n^{1/3} \rfloor$. Since $q \leq q_0$ and $c(q) \geq c(q_0)$ we can bound

$$\begin{aligned} P\left[\exists k : \overline{Y_k}\right] &\leq q_0 \left(1 - c(q_0)\right)^{\lfloor \frac{n}{4} \rfloor} = q_0 \frac{\left(1 - c(q_0)^2\right)^{\lfloor \frac{n}{4} \rfloor}}{\left(1 + c(q_0)\right)^{\lfloor \frac{n}{4} \rfloor}} \leq \\ &\leq \frac{q_0}{\left(1 + c(q_0)\right)^{\lfloor \frac{n}{4} \rfloor}} \leq \frac{q_0}{1 + \lfloor \frac{n}{4} \rfloor \, c(q_0)} \, . \end{aligned}$$

From Lemma 2.1 we have $c(q_0) = \left(\frac{1}{4}\right)^{q_0 - 1} \geq \left(\frac{1}{4}\right)^{\log n^{1/3}} = 2^{-2 \log n^{1/3}} = n^{-2/3}$. Hence

$$P\left[\exists k : \overline{Y_k}\right] \leq \frac{\log n^{1/3}}{1 + \lfloor \frac{n}{4} \rfloor \, n^{-2/3}} \leq \frac{\log n^{1/3}}{\frac{n}{5} n^{-2/3}} = \frac{1}{15} \frac{\log n}{n^{1/3}} \, .$$

□

3 Proof of the Theorem

Using the results of [2] we can state the following useful proposition.

Proposition 3.1 *Let q be a sufficiently large prime number; then the number of subsets $S \subseteq \{1, 2, \ldots, q-1\}$ such that the sum $\sum_{i \in S} i \equiv k \bmod q$ is $2^{q-1} q^{-1} (1 + o(1))$ for any $k = 0, 1, \ldots, q-1$.*

Combining the result of Lemma 2.2 with the previous proposition we can prove Theorem 1.2.

Proof of Theorem 1.2: Let us consider a graph $G \in \mathcal{G}_{n,\frac{1}{2}}$. It is possible to find disjoint pairs of vertices u_i, v_i such that the number of connecting 2-paths between u_i and v_i is equal to i modulo q for all $i = 1, 2, \ldots, q-1$ with probability at least $1 - \frac{1}{15} \frac{\log n}{n^{1/3}}$. Let us define graphs G_S for any $S \subseteq \{1, 2, \ldots, q-1\}$ obtained from G by adding all edges $\{u_i, v_i\}$ with $i \in S$ (if they are not in G) and by deleting all edges $\{u_i, v_i\}$ with $i \notin S$ (if they are in G). The probability of all G_S in the probability space $\mathcal{G}_{n,\frac{1}{2}}$ is the same.

From now on, all computation will be done modulo q. Let t_0 be the number of triangles of $G_\emptyset$. Since the pairs u_i, v_i are disjoint, edges $\{u_i, v_i\}$ do not affect the number of 2-paths between these pairs. Then the number of triangles of G_S is $t_0 + \sum_{i \in S} i$. By Proposition 3.1, the probability that

the number of triangles of G_S is equal to k modulo q is $q^{-1}(1+o(1))$, where S is chosen randomly so that each element is chosen independently with probability $\frac{1}{2}$.

Now, consider an arbitrary ordering of pairs of vertices and to each graph $G \in \mathcal{G}_{n,\frac{1}{2}}$ assign (if possible) the pairs u_i, v_i, $i = 1, 2, \ldots, q-1$, described above which are minimum in the fixed ordering. Note that graphs G_1 and G_2 are assigned the same pairs u_i, v_i and are identical on all other pairs if and only if they are G_{S_1} and G_{S_2} for some G and $S_1, S_2 \subseteq \{1, 2, \ldots, q-1\}$. Hence they have the same probability in $\mathcal{G}_{n,\frac{1}{2}}$.

Up to a $\left(\frac{1}{15}\frac{\log n}{n^{1/3}}\right)$-fraction, we have partitioned the graphs of $\mathcal{G}_{n,\frac{1}{2}}$ into classes of equal size such that in each class the number of triangles computed modulo q is almost uniformly distributed. Since $\frac{\log n}{n^{1/3}}$ tends to 0 for n going to infinity, we can conclude that the probability that $G_{n,\frac{1}{2}}$ contains exactly k triangles modulo q is $q^{-1}(1+o(1))$ for any $k = 0, 1, \ldots, q-1$.

□

4 Acknowledgements

The authors would like to thank to D. Welsh for helpful discussions and to M. Klazar for pointing out the paper of Erdős and Heilbronn.

References

[1] B. Bollobás, L. Pebody, O. Riordan. Contraction-deletion invariants for graphs. *J. Combin. Theory Ser. B* **80** (2000), 320–345

[2] P. Erdős and H. Heilbronn. On the addition of residue clesses mod p. *Acta Arithmetica* **9** (1964), 149–159

[3] H. Whitney. 2-isomorphic graphs. *Amer. J. Math.* **55** (1933), 245–254