

On Brown's Conjecture on Accessible Sets

Veselin Jungić*

Abstract

In this note we use a sequence constructed by H. Furtsenberg in 1981 to disprove the following conjecture posted by T. Brown: If a set of positive numbers L is such that for any finite coloring of \mathbb{N} there are arbitrarily long monochromatic sequences with all gaps in L , then for any finite coloring of \mathbb{N} there are arbitrarily long monochromatic arithmetic progressions whose common differences belong to L .

1 Introduction

Let \mathbb{N} be the set of positive integers. For $r \in \mathbb{N}$, an r -coloring of \mathbb{N} is a function $f : \mathbb{N} \rightarrow A$, with $|A| = r$. A finite coloring is an r -coloring for some r . If f is a finite coloring and if $B \subseteq \mathbb{N}$ satisfies $|f(B)| = 1$, we say that B is f -monochromatic. An arithmetic progression of length k and common difference d , $k, d \in \mathbb{N}$, is a set of the form $\{a + (i - 1)d : i \in [1, k]\}$, for some $a \in \mathbb{N}$.

Van der Waerden's theorem [5] on arithmetic progressions says that for any finite coloring f and any $k \in \mathbb{N}$ there is an f -monochromatic arithmetic progression of length k . Brown, Graham, and Landman in [2] study subsets L of \mathbb{N} such that van der Waerden's theorem can be strengthened to guarantee the existence of arbitrarily long monochromatic arithmetic progressions having common differences in L . Sets with this property are called *large*. Somehow surprisingly there are many large sets. For example, by the Polynomial van der Waerden's Theorem [1], if p is a polynomial with rational coefficients taking integer values on the integers and satisfying $p(0) = 0$ then $|p(\mathbb{N})|$ is large.

*Department of Mathematics, Simon Fraser University, Burnaby, BC, V5A 1S6, Canada E-mail: vjungic@sfu.ca

For a given set D of positive integers Landman and Robertson ([4], Definition 10.12) define a k -term D -diffsequence as a sequence $x_1 < x_2 < \dots < x_k$ such that $x_i - x_{i-1} \in D$ for all $i = 2, 3, \dots, k$. D is said to be *accessible* if for any finite coloring f of positive integers there are arbitrarily long f -monochromatic D -diffsequences.

It is known ([4], Theorem 10.27) that for any infinite set T of positive integers, the difference set $T - T = \{|t - s| : s, t \in T\}$ is accessible.

T. Brown conjectured ([4], Research Problem 10.9) that every accessible set is large.

We use a sequence of positive numbers constructed by H. Furstenberg [3] to disprove Brown's conjecture. In [3] this sequence is used to show that there is a set that intersects each IP -set of \mathbb{Z} , but does not intersect each difference set of \mathbb{Z} . A set $Q \subseteq \mathbb{Z}$ is an IP -set of \mathbb{Z} if there is sequence $\{a_i\}_{i \in \mathbb{Z}}$ of not necessarily distinct integers so that $Q = \{\sum_{i \in F} a_i : F \subseteq \mathbb{N} \text{ and } |F| < \infty\}$ for some $S \subseteq \mathbb{Z}$. (IP stands for *infinite-dimensional parallelepiped*.)

2 Not All Accessible Sets Are Large

In this section we show that there is an accessible set that is not large.

It is not difficult to check that

$$\|x\| = \min\{|x + n| : n \in \mathbb{Z}\}$$

is a norm on \mathbb{R} . It is known ([3], page 22) that for any $\alpha, a \in (0, 1)$, with α irrational, and any $\varepsilon > 0$ there is $n \in \mathbb{N}$ such that

$$\max\{\|n\alpha\|, \|n^2\alpha - a\|\} < \varepsilon.$$

Let $\alpha \in (0, 1)$ be irrational and let $\varepsilon \in (0, \frac{1}{8})$. We define the set $S = \{s_i\}_{i \in \mathbb{N}}$ inductively in the following way. Let $s_1 \in \mathbb{N}$ be such that

$$\max\left\{\|s_1\alpha\|, \left\|s_1^2\alpha - \frac{1}{4}\right\|\right\} < \varepsilon.$$

If s_1, \dots, s_k are defined, then $s_{k+1} \in \mathbb{N}$ is such that

$$\max\left\{\|s_{k+1}\alpha\|, \left\|s_{k+1}^2\alpha - \frac{1}{4}\right\|\right\} < \frac{\varepsilon}{\prod_{i=1}^k s_i}.$$

We note that for all $n \in \mathbb{N}$

$$\left\|s_n^2\alpha - \frac{1}{4}\right\| < \varepsilon$$

and, for all $m, n \in \mathbb{N}$ such that $m < n$,

$$\|s_m s_n \alpha\| \leq s_m \|s_n \alpha\| < \frac{\varepsilon}{\prod_{i \neq m} s_i} < \varepsilon.$$

Thus, for $m \neq n$ we have that

$$\left\| (s_m - s_n)^2 \alpha - \frac{1}{2} \right\| \leq \left\| s_m^2 \alpha - \frac{1}{4} \right\| + 2 \|s_m s_n \alpha\| + \left\| s_n^2 \alpha - \frac{1}{4} \right\| < 4\varepsilon.$$

In particular, since $\varepsilon < \frac{1}{8}$, we have that $s_n \neq s_m$ if $n \neq m$. Hence, S is an infinite set and by ([4], Theorem 10.27), $L = S - S = \{|s_m - s_n| : m \neq n\}$ is accessible.

We claim that there is a finite coloring of \mathbb{N} with no monochromatic 3-term arithmetic progression having its common difference in L .

For $m \in \mathbb{N}$ we define an m -coloring $f_m : \mathbb{N} \rightarrow \{1, \dots, m\}$ in the following way. By definition $f_m(n) = i$ if and only if $\left\| \frac{n(n-1)}{2} \alpha \right\| \in \left(\frac{i-1}{2m}, \frac{i}{2m} \right)$.

Suppose that $n, p \in \mathbb{N}$ are such that $\{n, n+p, n+2p\}$ is f_m -monochromatic and suppose that i is such that $f_m(\{n, n+p, n+2p\}) = \{i\}$. Then, for all $k \in \{n, n+p, n+2p\}$, $\left\| \frac{k(k-1)}{2} \alpha - \frac{i-1}{2m} \right\| \in \left(0, \frac{1}{2m} \right)$.

Particularly, for $k = n+p$

$$\begin{aligned} \frac{1}{2m} &> \left\| \frac{n(n-1)}{2} \alpha - \frac{i-1}{2m} + pn\alpha + \frac{p(p-1)}{2} \alpha \right\| \geq \\ &\geq \left\| pn\alpha + \frac{p(p-1)}{2} \alpha \right\| - \left\| \frac{n(n-1)}{2} \alpha - \frac{i-1}{2m} \right\|. \end{aligned}$$

It follows that $\left\| pn\alpha + \frac{p(p-1)}{2} \alpha \right\| < \frac{1}{m}$, or, equivalently, $\|2pn\alpha + p(p-1)\alpha\| < \frac{2}{m}$.

From

$$\begin{aligned} \frac{1}{2m} &> \left\| \frac{(n+2p)(n+2p-1)}{2} \alpha - \frac{i-1}{2m} \right\| \geq \\ &\geq \|p^2\alpha\| - \left\| \frac{n(n-1)}{2} \alpha - \frac{i-1}{2m} \right\| - \|2pn\alpha + p(p-1)\alpha\| > \\ &> \|p^2\alpha\| - \frac{1}{2m} - \frac{2}{m} \end{aligned}$$

we have that if p is the common difference of a f_m -monochromatic 3-term arithmetic progression, then $\|p^2\alpha\| < \frac{3}{m}$.

Let $k \in \mathbb{N}$ be such that $\frac{1}{k} < \frac{1}{3} (\frac{1}{2} - 4\varepsilon)$ and let $l, n \in \mathbb{N}, l \neq n$. From

$$\frac{1}{2} - \left\| (s_l - s_n)^2 \alpha \right\| \leq \left\| (s_l - s_n)^2 \alpha - \frac{1}{2} \right\| < 4\varepsilon$$

we have that

$$\left\| (s_l - s_n)^2 \alpha \right\| > \frac{1}{2} - 4\varepsilon > \frac{3}{k}.$$

Thus, f_k is a finite coloring of \mathbb{N} such that there is no f_k -monochromatic 3-term arithmetic progression having common difference in L .

Therefore, L is not large.

3 Brown-Graham-Landman Conjecture

For $k \in \mathbb{N}$, a set of positive integers L is said to be chromatically k -intersective if for any coloring f of positive integers there is an f -monochromatic k -term arithmetic progression whose common difference belongs to L . Clearly, L is large if it is chromatically k -intersective for all k . The difference set of Furstenberg's sequence from the previous section is an example of a set that is chromatically 2-intersective, but not chromatically 3-intersective.

Another way to define large sets is to start by fixing the number of colors and then to vary the length of monochromatic arithmetic progressions. For $r \in \mathbb{N}$, a set of positive integers L is said to be r -large if for any r -coloring f of positive integers there are arbitrarily long f -monochromatic arithmetic progressions whose common differences belong to L . L is large if it is r -large for all r . It is not known if there is an r -large set that is not large.

Brown, Graham, and Landman posted the following conjecture [2].

Conjecture 1 *Every 2-large set is large.*

References

- [1] V. Bergelson and A Leibman. Polynomial extension of van der Waerden's and Szemerédi's theorems. *J. Amer. Math. Soc.*, 9:725–753, 1996.
- [2] T. C. Brown, R. L. Graham, and B. Landman. On the set of common differences in van der Waerden's theorem on arithmetic progressions. *Canadian Math. Bull.*, 42:25–36, 1991.

- [3] H. Furstenberg. *Recurrence in Ergodic Theory and Combinatorial Number Theory*. Princeton University Press, 1981.
- [4] B. M. Landman and A. Robertson. *Ramsey Theory on the Integers*. American Mathematical Society, 2003.
- [5] B. L. van der Waerden. Beweis einer Baudetschen Vermutung. *Nieuw Archief voor Wiskunde*, 15:212–216, 1927.