

# THE CHROMATIC POLYNOMIAL OF FATGRAPHS AND ITS CATEGORIFICATION

MARTIN LOEBL AND IAIN MOFFATT

ABSTRACT. We introduce a homology theory for embedded graphs whose graded Euler characteristic is the chromatic polynomial, and whose Poincaré polynomial is invariant on different planar embeddings of the same graph. We conjecture that the homology can detect the genus of the embedding. We show that for planar embedded graphs our homology is obtained by ‘adding coefficients’ to the Khovanov’s categorification of the Jones polynomial. At present there are extensive efforts by mathematicians and physicists to extend the Khovanov homology theory. Both the chromatic polynomial and the Jones polynomial are determined by the Tutte polynomial on the (signed) graphs, or equivalently by the  $+ - J$  Potts partition function. We extend this to the homology: we categorify the Bollobás-Riordan topological Tutte polynomial of signed embedded graphs so that the homology of the chromatic polynomial can be recovered by a projection and the Khovanov homology theory of the Jones polynomial can be recovered using a ‘universal coefficients theorem’.

## CONTENTS

1. Introduction	2
1.1. Fatgraphs	3
1.2. Graph Polynomials	4
1.3. Statement of Results	5
1.4. Acknowledgement	6
2. Construction of the Homology	6
3. Properties of the Homology	11

---

*Date:* This edition: June 4, 2006      First edition: November 22, 2005.

2000 *Mathematics Classification.* Primary 05C10. Secondary 57M15.

*Key words and phrases:* chromatic polynomial, fatgraphs, Khovanov homology, categorification, Bollobás-Riordan polynomial .

4. The Bollobás-Riordan Polynomial	14
4.1. Chromatic Homology as an Extension of Bollobás-Riordan Homology	14
4.2. Categorification of the Multivariate Bollobás-Riordan Polynomial	16
5. Independence of Planar Embeddings	21
5.1. The Relation to Khovanov Homology and Knots	21
5.2. Proof of Theorem 5.1	23
6. Helme-Guizon and Rong's Chromatic Homology	30
Appendix A. Table of Polynomials	31
References	32

## 1. INTRODUCTION

Over the last few years there has been a lot of activity on the categorification of knot invariants. This exciting area of mathematics was instigated by M. Khovanov's construction of a bigraded homology theory whose graded Euler characteristic is equal to the Jones polynomial ([Kh]). Khovanov's homology groups are themselves knot invariants and are in fact strictly stronger than the Jones polynomial. At present there are extensive efforts by mathematicians and physicists to extend the Khovanov homology theory, for example [DGR, GW, KR1, KR2]. In 2005, categorification passed over into graph theory with the introduction of a homology theory for abstract graphs whose Euler characteristic is the chromatic polynomial. This was introduced in [HGR1] and graph homology has been further developed in [CCR, CU, HGR2, HGRP, JHR, Pr, Ro, St1, St2].

Here we present a homology theory for embedded graphs whose Euler characteristic is the chromatic polynomial. As with Khovanov homology, the homology groups themselves are invariants of the embedded graphs and are strictly stronger than the chromatic polynomial. This homology theory is closely related to Khovanov homology when the graphs are planar and the Bollobás-Riordan polynomial for higher genus graphs. The Bollobás-Riordan polynomial is equivalent to the  $+ - J$  Potts partition function with an additional topological information. This relation enables us to propose a categorification of the full Bollobás-Riordan polynomial of signed embedded graphs which contains Khovanov homology and also our homology of the chromatic polynomial of embedded graphs.

There are interesting and well known connections between link polynomials and graph polynomials. Perhaps the best known of these is the expression of the Jones polynomial as an evaluation of the Tutte polynomial of its medial graph (see e.g. [Bo]). Our results can be interpreted as an extension of such connections between link and graph polynomials on the homological level. The connection of the homology of the chromatic polynomial of embedded graphs with Khovanov homology is central to the proof of Theorem 5.1, which shows that the Poincaré polynomial is independent of the choice of the planar embedding of a graph. Another key ingredient in this result is the classical combinatorial understanding of the structure of the planar embeddings of a given planar graph. Such structural combinatorial understanding is missing for the higher genus graphs, and our homology theory may be a new useful tool. In particular we conjecture that the homology theory can detect the genus of the embedded graph.

Most of the formalism concerning the embedded graphs is taken from [CP], although there is a basic notational difference. Embedded graphs are called *ribbon graphs* in [CP], however we chose the term *fatgraphs* instead which is standard in theoretical physics (see e.g. [DiF]).

**1.1. Fatgraphs.** Let  $G = (V, E)$  be an undirected graph, possibly with loops and multiple edges. Each subgraph  $(V, W)$ ,  $W \subset E$  of  $G$  is called a *spanning subgraph*. Let us denote by  $\mathcal{S}(G)$  the set of all spanning subgraphs of  $G$ . A graph  $F$  is called a *fatgraph* if for each vertex  $v \in V$ , there is a fixed cyclic order on edges adjacent to  $v$  (loops are counted twice). A fatgraph  $F$  may be regarded as a 2-dimensional surface with boundary, which will also be denoted by  $F$ . The surface is obtained from the fatgraph by fattening the vertices into discs (we will call these *islands*) and connecting them by untwisted fattened edges (which we call *bridges*) as prescribed by the cyclic orders. The *genus*,  $g(F)$  of a fatgraph  $F$  is defined to be the genus of this surface. It will always be clear from the context whether by  $F$  we mean the fatgraph or the surface. Note that we restrict to orientable surfaces here. For a fatgraph  $F$  we will usually denote its underlying graph by  $G = G(F)$ . Let  $V(F)$  be its set of vertices,  $E(F)$  its set of edges, and let  $v(F) = |V(F)|$ ,  $e(F) = |E(F)|$ ,  $r(F) = |V| - k(F)$  and  $n(F) = |E(F)| - r(F)$ . We denote the number of connected components of  $F$  by  $k(F)$ , and the number of connected components of the boundary of surface  $F$  by  $p(F)$ . The functions  $v, e, r, n, k$  will be used for graphs as well. We denote by  $\mathcal{S}(F)$  the set of the spanning fatsubgraphs of  $F$ , as for graphs.

**1.2. Graph Polynomials.** Let us recall the definitions of the Tutte and the chromatic polynomials of a graph  $G = (V, E)$ :

$$T(G, x, y) = \sum_{H \in \mathcal{S}(G)} (x-1)^{r(G)-r(H)} (y-1)^{n(H)},$$

$$M(G, u) = \sum_{H \in \mathcal{S}(G)} (-1)^{e(H)} u^{k(H)}.$$

The chromatic polynomial  $M(G, u)$  is a straightforward evaluation of  $T(G, x, y)$ .

In [BR], Bollobás and Riordan defined a fatgraph generalization of the Tutte polynomial. This three variable polynomial is defined by the state sum

$$(1) \quad R(F, x, y, z) = \sum_{H \in \mathcal{S}(F)} x^{r(F)-r(H)} y^{n(H)} z^{k(H)-p(H)+n(H)}.$$

The following lemma tells us that the coefficient of  $z$  is equal to twice the genus of the fatgraph  $H$  and we may therefore write

$$R(F, x, y, z) = \sum_{H \in \mathcal{S}(F)} x^{r(F)-r(H)} y^{n(H)} z^{2g(H)}.$$

**Lemma 1.1.** *The genus,  $g(F)$ , of a fatgraph  $F$  is determined by the formula  $g(F) = \frac{1}{2}(k(F) - p(F) + n(F))$ .*

*Proof.* Suppose  $F = \cup_{i=1}^k F_i$ , where each  $F_i$  is connected. By regarding  $F$  as an orientable surface we see  $2g(F) = \sum_i 2g(F_i) = \sum_i 2 - \chi(F_i)$ . If  $T_i$  is a triangulation of the surface  $F_i$ , this is equal to  $\sum_i 2 - (v(T_i) - e(T_i) + f(T_i) + \partial(F_i))$ , where  $\partial(F_i)$  denotes the number of boundary components of  $F_i$ . Now since the underlying map  $G_i$  of  $F_i$  is a deformation retract and the homology determines the Euler characteristic (see eg. [Ha]) we see that this is equal to  $\sum_i 2 - p(F_i) + e(F_i) - v(F_i) = 2k(F) - p(F) + e(F) - v(F)$  as required.  $\square$

If  $F$  is a fatgraph and  $G$  its underlying graph, then one can express the chromatic polynomial of  $G$  in terms of geometric information from  $F$ . The next lemma expresses the chromatic polynomial in the evaluation we use.

**Lemma 1.2.** *Let  $F$  be a fatgraph and  $G$  be its underlying graph. Then*

$$(2) \quad (q + q^{-1})^{e(F)} M(G, (q + q^{-1})^2) =$$

$$(-1)^{e(F)} (q + q^{-1})^{v(F)} \sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{(p(H) + 2g(H))} (-q)^{e(F) - e(H)} (1 + q^{-2})^{e(F) - e(H)}.$$

*Proof.* Using the formula  $2g(H) = k(H) - p(H) + n(H)$  and the definitions of  $n(H), r(H)$  above, we have  $2k(H) = p(H) + 2g(H) - e(H) + v(H)$ . Hence

$$M(G, u) = u^{1/2v(F)} \sum_{H \in \mathcal{S}(F)} u^{1/2p(H)+g(H)} [(-1)u^{1/2}]^{-e(H)}.$$

Substituting  $u^{1/2} = (q + q^{-1})$  we get

$$M(G, (q+q^{-1})^2) = (q+q^{-1})^{v(F)} \sum_{H \in \mathcal{S}(F)} (q+q^{-1})^{p(H)+2g(H)} (-1)^{-e(H)} (q+q^{-1})^{-e(H)}.$$

This easily implies 2. □

The main object of our study is exactly this evaluation and scaling of the chromatic polynomial. We set

$$\begin{aligned} Z(F, q) &= (q + q^{-1})^{e(F)} M(G, (q + q^{-1})^2) = (-1)^{e(F)} (q + q^{-1})^{v(F)} \\ &\quad \sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{p(H)+2g(H)} (-q)^{e(F)-e(H)} (1 + q^{-2})^{e(F)-e(H)}. \end{aligned}$$

**1.3. Statement of Results.** Let  $M = \bigoplus_{i \in \mathbb{Z}} M_i$  be a graded  $\mathbb{Z}$ -module. The *graded dimension* of  $M$  is defined by

$$\text{qdim}(M) := \sum_{i \in \mathbb{Z}} q^i \text{rk}(M_i) = \sum_{i \in \mathbb{Z}} q^i \dim_{\mathbb{Q}}(M_i \otimes_{\mathbb{Z}} \mathbb{Q}),$$

and if  $H = (H^i)_{i \in \mathbb{Z}}$  is the homology of some chain complex of graded  $\mathbb{Z}$ -modules, the *Poincaré polynomial* is defined by

$$P(H) = \sum_{i \in \mathbb{Z}} t^i \text{qdim}(H^i) \in \mathbb{Z}[q, q^{-1}, t, t^{-1}].$$

The Poincaré polynomial encodes all of the torsion-free information of the homology groups. The *Euler characteristic* is defined by  $\chi(H) = P(H)(t = -1)$ . It generalizes the usual Euler characteristic of graphs and surfaces.

In Section 2 we construct a graded chain complex for every fatgraph. The following theorem describes the main results of the paper.

**Theorem 1.3.** *Let  $F$  be a fatgraph and  $G$  be its associated graph. Let  $\mathcal{C}(F)$  be its chain complex as constructed in Section 2. Then the following hold:*

- (1) *the Euler characteristic of the homology  $(H^i(\mathcal{C}(F)) := (\ker d^i)/(\text{im } d^{i-1}))$  is equal to the chromatic polynomial  $Z(F, q)$ ;*
- (2) *the homology groups  $H$  are strictly stronger graph invariants than the chromatic polynomial;*

- (3) *the Poincaré polynomial is invariant on different planar embeddings of a planar graph  $G$ . However, the homology is dependent upon the genus of the embedding of a graph;*
- (4) *this homology theory may be extended to a categorification of the Bollobás-Riordan polynomial of the signed fat graphs, from which the Khovanov homology may be recovered.*


This theorem will follow from Theorems 2.2, which contains statement 1; Proposition 3.1, which gives some properties of the homology; Theorem 5.1 which contains statement 3 and section 4 which contains the construction for statement 4 above.

In the final section we show how the Bollobás-Riordan polynomial provides a framework in which to relate Helme-Guizon and Rong’s chromatic homology with the Khovanov homology of a certain associated link. This addresses the question ”What is the relationship [of Helme-Guizon and Rong’s chromatic homology] with the Khovanov homology for knots?”, which was posed in [HGR1].


**1.4. Acknowledgement.** We would like to thank Jo Ellis-Monaghan, Bojan Mohar and Irasema Sarmiento for helpful discussions. M. L. gratefully acknowledges the support of CONICYT via grant Anillo en Redes.

## 2. CONSTRUCTION OF THE HOMOLOGY

Let  $F$  be a fatgraph and  $G = (V, E)$  be its underlying graph. We call a spanning fatsubgraphs of a fatgraph a *state*. We see that The chromatic polynomial  $Z(F, q)$  is expressed in (2) as a sum over all states. Clearly each state of a fatgraph  $F$  is obtained by the removal of a set of bridges of  $F$ . For example:



*A fat-graph.*



*A state.*

We call the total number of edges of  $F$  minus the number of bridges in a state  $H$  the *height*, denoted by  $h$ , so  $h(H) = e(F) - e(H)$ .

Let us consider the following example of the state sum  $Z(F, q)$ :

$$(q + q^{-1})^5 - 3(q + q^{-1})^4(1 - q^{-2})(-q) + 3(q + q^{-1})^3(1 - q^{-2})^2(-q)^2 - (q + q^{-1})^2(1 - q^{-2})^3(-q)^3 = 0,$$

which, of course, is equal to zero. We will use this example to illustrate how the state sum for  $Z(F, q)$  gives rise to a chain complex. The approach here is similar to Bar-Natan's exposition of Khovanov homology in [BN1].

Notice that each state of the fatgraph in Equation 2 gives rise to (up to sign) a Laurent polynomial of the form

$$(3) \quad (q + q^{-1})^{(v+p+2g)}(1 + q^{-2})^h q^h,$$

where  $v = v(H)$ ,  $p = p(H)$ , etc.. We want to replace each such polynomial term of the state sum with a graded module whose graded dimension is equal to the polynomial.

To do this we define  $V$  to be the free, graded  $\mathbb{Z}$ -module with two basis elements  $v_-$  and  $v_+$  in graded degrees  $-1$  and  $+1$  respectively, and  $R$  to be the free, graded  $\mathbb{Z}$ -module with basis elements  $x_{-2}$  and  $x_0$  in graded degrees  $-2$  and  $0$ . Notice that  $\text{qdim}(V) = q + q^{-1}$  and  $\text{qdim}(R) = 1 + q^{-2}$ .

The *degree shift* operation  $\{\cdot\}$  on graded modules is defined by setting  $M\{l\}_m := M_{m-l}$ . Clearly  $\text{qdim}(M\{l\}) = q^l \cdot \text{qdim}(M)$ . We see that  $R = V\{-1\}$  and therefore every occurrence of  $R$  in this paper could be replaced with  $V\{-1\}$ . Although we retain the use of  $R$  for clarity, the reader should bear in mind that the two modules only differ by a grading shift. We will see that, in some sense,  $R$  plays the role of coefficients in the homology theory.

Now, upon observing that for two graded modules  $M$  and  $N$ ,  $\text{qdim}(M \otimes N) = \text{qdim}(M) \cdot \text{qdim}(N)$  and  $\text{qdim}(M \oplus N) = \text{qdim}(M) + \text{qdim}(N)$ , it is


easily seen that the modules

$$(4) \quad V^{\otimes(v+p+2g)} \otimes R^{\otimes h}\{h\}$$

have graded dimensions equal to the Laurent polynomial 3.

In order to simplify the text, we abuse notation and identify the state of a fatgraph with its assigned polynomial term (3) and its assigned graded module (4).

Next, as in the calculation for  $Z(F, q)$  above, we arrange the states into  $n = e(F)$  columns according to the height of the corresponding state of the fatgraph, so the  $i$ -th column contains all modules which come from states of height  $i$ . We then define the  $i$ -th chain module  $\tilde{\mathcal{C}}^i(F)$  to be the direct sum of all of the modules corresponding to states of height  $i$ . For example (ignoring the maps in the figure for the time being), the above calculation becomes:



Note that although the tensor powers of  $V$  are monotone in this example, this is not the case in general.


Our next task is to define chain maps. We begin with the observation that removing a bridge from a state of height  $h$  determines a state of height  $h + 1$ . The state obtained depends upon which bridge is removed. Notice that the state determined by the removal of a bridge will have one more or one less boundary component than the original state, and the genus of the

state will either be unchanged or will decrease by one. Whenever we can move from a state of height  $h$  to a state of height  $h + 1$  by the removal of an edge we will define a *per-edge map*

$$\delta : V^{\otimes(v+p+2g)} \otimes R^{\otimes h}\{h\} \longrightarrow V^{\otimes(v+p'+2g')} \otimes R^{\otimes(h+1)}\{h+1\},$$

where  $p' = p \pm 1$  and  $g' = g$  or  $g - 1$ . In order to define  $\delta$ , we first describe three maps which correspond to the tensor factors  $V^{\otimes p}$ ,  $V^{\otimes 2g}$  and  $R^{\otimes h}$ .

$V^{\otimes p}$ : In terms of the boundary components of the states, one of two things can happen when we move from one state to another by the deletion of a bridge: either two boundary components will be merged into one component or one component will be split into two components. For example,



Notice that by (4), a copy of the module  $V$  is assigned to each boundary component of a state. We fix a correspondence between components and the modules  $V$  once and for all (the homology will be independent of this choice). When two components merge we need a map  $\tilde{m} : V^{\otimes p} \rightarrow V^{\otimes(p-1)}$ . We define the map to be the identity on all copies of  $V$  which are unchanged by the addition of an edge, and to act on the two merging components by the multiplication defined on basis elements by

$$\tilde{m}' : \begin{cases} v_- \otimes v_- \mapsto 0 & v_+ \otimes v_- \mapsto v_- \\ v_+ \otimes v_+ \mapsto v_+ & v_- \otimes v_+ \mapsto v_- \end{cases} .$$

In the case where one component is split into two, we define a map  $\tilde{\Delta} : V^{\otimes p} \rightarrow V^{\otimes(p+1)}$  which is the identity on all factors of  $V^{\otimes p}$  except on the component being split where it acts on basis elements by

$$\tilde{\Delta}' : \begin{cases} v_+ \mapsto v_+ \otimes v_- + v_- \otimes v_+ \\ v_- \mapsto v_- \otimes v_- \end{cases} .$$

Notice that the two maps  $\tilde{m}'$  and  $\tilde{\Delta}'$  are the maps used in the definition of Khovanov homology ([Kh, BN1]). We will use this observation later.

$V^{\otimes 2g}$ : We identify  $V^{\otimes 0}$  with  $\mathbb{Z}$ . There are two cases. If  $g = g'$  then we set  $\hat{m}_g : V^{\otimes 2g} \rightarrow V^{\otimes 2g}$  equal to  $1^{\otimes 2g}$ . If  $g' = g - 1$  then we define  $\hat{m}_g : V^{\otimes 2g} \rightarrow V^{\otimes 2(g-1)}$  on its basis elements by

$$\hat{m}_g : v_{i_1} \otimes \cdots \otimes v_{i_{2g}} \mapsto \sum v_{j_1} \otimes \cdots \otimes v_{j_{2g-2}},$$

where the sum is over all basis elements of  $V^{\otimes 2(g-1)}$  whose graded dimension is equal to the graded dimension of the basis element  $v_{i_1} \otimes \cdots \otimes v_{i_{2g}}$  of  $V^{2g}$ , ie.  $\widehat{m}_g$  is a map of degree 0.

$R^{\otimes h}$ : Setting  $R^0 = \mathbb{Z}$ , we define  $\overline{\Delta}_h : R^{\otimes h} \rightarrow R^{\otimes(h+1)}$  by

$$\overline{\Delta}_h : \begin{cases} 1 \mapsto x_0, & \text{when } h = 0 \\ y \otimes x_i \mapsto y \otimes \sum_{k+l=i} x_k \otimes x_l, & \text{otherwise} \end{cases} .$$

Again this is a degree 0 map.

We take the tensor of these maps and define maps from  $V^{\otimes(v+p+2g)} \otimes R^{\otimes h}\{h\}$  to  $V^{\otimes(v+p'+2g')} \otimes R^{\otimes(h+1)}\{h+1\}$  by

$$m = 1^{\otimes v} \otimes \widehat{m} \otimes \widehat{m}_g \otimes \overline{\Delta}_h \quad \text{and} \quad \Delta = 1^{\otimes v} \otimes \widetilde{\Delta} \otimes \widehat{m}_g \otimes \overline{\Delta}_h .$$

Next we need to assign a sign to each of the maps  $m$  and  $\Delta$  to obtain the per-edge maps  $\delta$ . This is done as in [Kh, BN1] by realizing the states as vertices of an  $n$ -dimensional cube and the per-edge maps as its edges. To do this we label the bridges of  $F$  with  $1, \dots, n$ . The homology is independent of the choice of labeling, the proof of this follows [HGR1] Theorem 2.12. Then each state of  $F$  can be represented by the vertex of a  $n$ -dimensional cube  $(\alpha_1, \dots, \alpha_n)$ , by setting  $\alpha_i = 0$  if the bridge labelled  $i$  is in the state, and  $\alpha_i = 1$  if it is not. Now a per-edge map is a map from a state labelled  $(\alpha_1, \dots, \alpha_{j-1}, 0, \alpha_{j+1}, \dots, \alpha_n)$  to one labeled  $(\alpha_1, \dots, \alpha_{j-1}, 1, \alpha_{j+1}, \dots, \alpha_n)$ . The sign  $(-1)^{\sum_{i < j} \alpha_i}$  is then assigned to each of the maps  $m$  and  $\Delta$  in the complex. This gives the per-edge maps  $\delta$ .

Finally the differential  $\widetilde{\partial}^h : \widetilde{\mathcal{C}}^h \rightarrow \widetilde{\mathcal{C}}^{h+1}$  is obtained as the sum of all of the per-edge maps between the tensor factors of  $\widetilde{\mathcal{C}}^h$  and  $\widetilde{\mathcal{C}}^{h+1}$ .

Let  $\widetilde{\mathcal{C}}(F)$  denote the complex  $(\widetilde{\mathcal{C}}^h, \widetilde{\partial}^h)$  constructed as above.

**Lemma 2.1.**  $\widetilde{\mathcal{C}}$  is a chain complex (ie.  $\widetilde{\partial} \circ \widetilde{\partial} = 0$ ) and the differentials are of graded degree 0.

*Proof.* The per-edge maps are (co)associative and (co)commutative, the first statement then follows as the per-edge maps around each state anti-commute. The second statement follows since  $\widehat{m}$  and  $\widetilde{\Delta}$  are of degree  $-1$  and therefore are of degree zero once the target is shifted by  $\{1\}$ , and  $\widehat{m}_g$  and  $\overline{\Delta}_r$  are of degree 0.  $\square$

Finally, in order to deal with the factor  $(-1)^{e(F)}$  of  $Z(F, q)$ , we define the *height shift* operation  $[\cdot]$  on chain complexes by  $(\mathcal{C}^i, \partial^i)[s] := (\mathcal{C}^{i-s}, \partial^{i-s})$ .

We can then normalize the chain complex by  $[-e(F)]$  and define

$$\mathcal{C} = (\mathcal{C}^h, \partial^h) = \tilde{\mathcal{C}}(F)[-e(F)] = (\tilde{\mathcal{C}}^h, \tilde{\partial}^h)[-e(F)].$$

Recall that the *homology* of a chain complex  $\mathcal{C} = (\mathcal{C}^h, \partial^h)$  is the sequence  $H(\mathcal{C}) = (H^i(\mathcal{C}))_{i \in \mathbb{Z}}$ , where  $H^i(\mathcal{C}) = \ker(\partial^i)/\text{im}(\partial^{i-1})$ .

**Theorem 2.2.** *Let  $F$  be a fatgraph,  $G$  its associated graph and  $\mathcal{C}(F)$  its chain complex. Then the Euler characteristic of the homology  $H(\mathcal{C}(F))$  is equal to the scaled chromatic polynomial  $Z(F, q)$ . Moreover the homology is an invariant of fatgraphs and is a strictly stronger invariant than the chromatic polynomial  $\chi_G$ .*

*Proof.* The first statement follows essentially by construction: it is well known that when the differentials are of graded degree 0,  $\text{qdim}(H^i(\mathcal{C}))$  is equal to  $\text{qdim}(\mathcal{C}^i)$ , and since in turn this is equal to the sum of the graded dimensions of the states of height  $i$ , the result follows (this is the graded extension of a classical result in topology, see eg [Ha] page 146. The shift  $[-e(F)]$  ensures that the alternating sum of  $\text{qdim}(H^i(\mathcal{C}))$  has the correct sign. The second statement follows by a calculation (see Proposition 3.1).  $\square$

### 3. PROPERTIES OF THE HOMOLOGY

The following proposition summarizes the basic properties of our homology theory.

**Proposition 3.1.** *The following hold for the homology  $H(\mathcal{C}(F))$ :*



- (1) *the homology groups are strictly stronger than the chromatic polynomial;*
- (2) *the homology will differentiate between graphs which differ only by multiple edges or loops;*
- (3) *the homology is not a Tutte-Grothendieck invariant;*
- (4) *if  $F$  is the disjoint union of fatgraphs  $F_1$  and  $F_2$ , then*

$$H^i(\mathcal{C}(F)) = \left( \bigoplus_{p+q=i} H^p(\mathcal{C}(F_1)) \otimes H^q(\mathcal{C}(F_2)) \right) \oplus \left( \bigoplus_{p+q=i-1} \text{Tor}_1(H^p(\mathcal{C}(F_1)), H^q(\mathcal{C}(F_2))) \right);$$



- (5) let  $e$  be a bridge of a fatgraph  $F$  and let  $F - e$  denote  $F$  with the bridge  $e$  deleted and  $F/e$  denote  $F$  contracted along the edge  $e$ , then there exists a deletion-contraction exact sequence

$$\begin{array}{ccc} H^*(\mathcal{C}(F - e)) \otimes R & \longrightarrow & H^*(\mathcal{C}(F)) \\ & \searrow & \swarrow \\ & H^*(\mathcal{C}(F/e)) \otimes V & \end{array} ;$$

- (6) If  $F_1$  is a subfatgraph of  $F_2$  the inclusion map induces a homomorphism  $H(i) : H(F_1) \rightarrow H(F_2)$  in homology.

*Proof. 1.* This follows by a calculation. For example the fatgraphs  and  have the same chromatic polynomials but different homology.

2. This follows since the homology in the highest degree is non-zero (since  $\bar{\Delta}_{n-1}$  is not surjective).

3. Again this follows by a calculation. For example the fatgraphs  and  have different homology. Note that this example also shows that the Poincaré polynomial is not a Tutte-Grothendieck invariant.

4. As in [HGR1], the chain complex of  $F$  can be written  $\mathcal{C}(F) = \mathcal{C}(F_1) \otimes \mathcal{C}(F_2)$ . Then, since the chain complexes are free, the result follows by an application of the Künneth formula (see eg. [Br]).

5. We may assume that  $e$  is the bridge which is labelled last so that  $(\alpha_1, \dots, \alpha_{n-1}, 0)$  is the vertex label of a state containing the bridge  $e$  while  $(\alpha_1, \dots, \alpha_{n-1}, 1)$  is the label of the corresponding state with the bridge  $e$  deleted. Now, let  $\alpha = (\alpha_1, \dots, \alpha_{n-1})$  be a state of  $F - e$  of height  $h$ . Then  $\alpha' = (\alpha_1, \dots, \alpha_{n-1}, 1)$  gives a state of  $F$ . Notice that the fatgraphs of the states  $\alpha$  and  $\alpha'$  are identical but the state  $\alpha'$  is of height  $h + 1$ . Therefore  $V^{\otimes(v+p+2g)} \otimes R^{\otimes h}\{h\}$  is the module assigned to the state  $\alpha$ . The map

$$\eta_\alpha : V^{\otimes(v+p+2g)} \otimes R^{\otimes h}\{h\} \otimes R \longrightarrow V^{\otimes(v+p+2g)} \otimes R^{\otimes(h+1)}\{h+1\},$$

which takes basis elements to themselves, then induces a map  $\eta : \mathcal{C}(F - e) \otimes R \rightarrow \mathcal{C}(F)$ .

We also need to define a map  $\nu : \mathcal{C}(F) \rightarrow \mathcal{C}(F/e) \otimes V$ . To do this, suppose that  $\alpha' = (\alpha_1, \dots, \alpha_n)$  is assigned to a state of  $F$ . Then  $\alpha'' = (\alpha_1, \dots, \alpha_{n-1})$  is a state of  $F/e$ . If  $\alpha_n = 0$ , then the fatgraph in the state  $\alpha''$  is equal to the fatgraph of  $\alpha'$  contracted along the bridge  $e$ . Notice that the height of the two states  $\alpha'$  and  $\alpha''$  are equal, however the fatgraphs  $F$  has one more island than  $F/e$ . We then define a map  $\nu_{\alpha'} : V^{\otimes(v+p+2g)} \otimes R^h\{h\} \longrightarrow$

$V^{\otimes((v-1)+p+2g)} \otimes R^h\{h\} \otimes V$  by

$$\begin{aligned} & \nu_{\alpha'} : v_1 \otimes \cdots \otimes v_{v+p+2g} \otimes x_1 \otimes \cdots \otimes x_h \longmapsto \\ & \longmapsto \begin{cases} v_2 \otimes \cdots \otimes x_h \otimes v_1 & \text{if } \alpha' = (\alpha_1, \dots, \alpha_{n-1}, 0) \\ 0 & \text{otherwise} \end{cases}, \end{aligned}$$

where  $v_1 \otimes \cdots \otimes v_{v+p+2g} \otimes x_1 \otimes x_h$  denotes a basis element of  $V^{\otimes(v+p+2g)} \otimes R^h\{h\}$ . These induce  $\nu : \mathcal{C}(F) \rightarrow \mathcal{C}(F/e) \otimes V$ .

The next step is to show that  $\eta$  and  $\nu$  are chain maps, that is  $\partial \circ \eta = \eta \circ \partial$  and  $\partial \circ \nu = \nu \circ \partial$ . It is enough to show this for the per-edge maps, *ie.* that  $\eta$  and  $\nu$  commute with  $\delta$ . Now since each  $\eta_\alpha$  maps a state to a state with the same fatgraph, the corresponding per-edge maps in the two complexes will both be of the form of  $\pm m$  or  $\pm \Delta$ , and since we chose the edge  $e$  to be the last element  $\alpha_n$  in the vertex labeling, the maps will occur with the same sign in both complexes. Therefore it is enough to show  $m \circ \eta = \eta \circ (m \otimes 1)$  and  $\Delta \circ \eta = \eta \circ (\Delta \otimes 1)$ , which is easily verified.

Similarly, since  $\nu$  maps a fatgraph to 0 or the fatgraph contracted along the edge  $e$ , it is enough to show  $m \circ \nu = \nu \circ (m \otimes 1)$  and  $\Delta \circ \nu = \nu \circ (\Delta \otimes 1)$ . Again this is easily verified.

It is not hard to see that the chain maps  $\eta$  and  $\nu$  form a short exact sequence of chain complexes:

$$0 \rightarrow \mathcal{C}^{i-1}(F - e) \otimes R \rightarrow \mathcal{C}^i(F) \rightarrow \mathcal{C}^i(F/e) \otimes V \rightarrow 0.$$

This induces the long exact sequence in homology, completing the proof.

6. Since there is an inclusion  $F_1 \hookrightarrow F_2$ , for each state  $H$  of  $F_1$  there is a state  $H_2$  of  $F_2$  whose fatgraph consists of  $H$  and  $k$  independent islands, where  $k = v(F_2) - v(F_1)$ . This inclusion between states of fatgraphs gives an inclusion  $V^{\otimes v(H)+p(H)+2g(H)} \otimes R^{\otimes h(H)}\{h(H)\} \hookrightarrow V^{\otimes v(H_2)+p(H_2)+2g(H_2)} \otimes R^{\otimes h(H_2)}\{h(H_2)\}$  between modules and therefore a map  $\mathcal{C}(F_1) \hookrightarrow \mathcal{C}(F_2)$  of chain complexes. This is clearly a chain map and therefore induces a homomorphism in homology. (Note that this property also arises in [JHR] Section 6.3.)  $\square$

*Remark 3.2.* The homology theory is genuinely different from the categorification of the chromatic polynomial for abstract graphs constructed in [HGR1]. This can be seen immediately from properties 2. and 3. of Proposition 3.1. In fact Helme-Guizon and Rong's homology groups are trivial on graphs which contain loops and can not distinguish graphs which differ only by multiple edges (see [HGR1]).

*Remark 3.3.* As with [HGR2], our construction works over a variety of algebras. However, Khovanov’s homology of links requires a unique choice of algebra for isotopy invariance, and since our proof of the invariance of the Poincaré polynomial for planar fatgraph homology (Theorem 5.1) is dependent upon the invariance of Khovanov homology, we do not know if the corresponding result holds over different algebras.

As mentioned several times, our homology theory is motivated by Khovanov’s categorification of the Jones polynomial. For *planar fatgraphs*, the relation is made explicit by Theorem 5.2 which states that for fatgraphs of genus zero, our homology is obtained by “adding coefficients” to the Khovanov’s homology of the corresponding alternating link diagram.

We also show that for general genus, the Jones polynomial may be replaced by a suitable evaluation and scaling of the Bollobás-Riordan polynomial  $R$  defined in the introduction. We remark that Chmutov and Pak [CP] have proven recently that the Jones polynomial (represented by the Kauffman bracket as in the work of Khovanov) of (a class of) alternating links in the 3-manifold  $\Sigma \times I$ , where  $\Sigma$  is an oriented surface and  $I$  is a unit interval, is an evaluation of  $R$ . In the next section we state and prove a “universal coefficient theorem” linking our homology with the categorification of a suitable evaluation and scaling of  $R$ .

#### 4. THE BOLLOBÁS-RIORDAN POLYNOMIAL

**4.1. Chromatic Homology as an Extension of Bollobás-Riordan Homology.** We first write a suitable evaluation of  $R(F, x, y, z)$  as a state sum.

**Lemma 4.1.** *Let  $F$  be a fatgraph and  $G$  be its underlying graph. Then*

$$\begin{aligned}
 (5) \quad & \left( x^{k(F)} y^{v(F)} [-y^{-1}(xy)^{1/2}]^{e(F)} R(F, x, y, z) \right) \Big|_{x=-q(q+q^{-1}), y=-q^{-1}(q+q^{-1}), z=1} \\
 & = (-1)^{e(F)} \sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{v(H)+p(H)+2g(H)} (-q)^{e(F)-e(H)}.
 \end{aligned}$$

*Proof.* Using  $r(F) = v(F) - k(F)$ ,  $n(F) = e(F) - r(F)$  and  $g(F) = 1/2(k(F) - p(F) + n(F))$ , we have

$$\begin{aligned}
R(F, x, y, z) &= x^{-k(F)} y^{-v(F)} \sum_{H \in \mathcal{S}(F)} (xy)^{k(H)} y^{e(H)} z^{2g(H)} \\
&= x^{-k(F)} y^{-v(F)} \sum_{H \in \mathcal{S}(F)} (xy)^{1/2(v(H)+p(H)+2g(H))} [y(xy)^{-1/2}]^{e(H)} z^{2g(H)} \\
&= x^{-k(F)} y^{-v(F)} [y^{-1}(xy)^{1/2}]^{-e(F)} \\
&\quad \sum_{H \in \mathcal{S}(F)} (xy)^{1/2(v(H)+p(H)+2g(H))} [y^{-1}(xy)^{1/2}]^{e(F)-e(H)} z^{2g(H)}.
\end{aligned}$$

The lemma follows upon substituting  $x = -q(q + q^{-1})$ ,  $y = -q^{-1}(q + q^{-1})$  and  $z = 1$ .  $\square$

We define the *restricted Bollobás-Riordan polynomial* by

$$\widehat{R}(F, q) = \sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{v(H)+p(H)+2g(H)} (-q)^{e(F)-e(H)}.$$

We may categorify this in a similar way as in Section 2, by assigning the modules  $V^{\otimes(v(H)+p(H)+2g(H))} \{h\}$  to the state  $(q+q^{-1})^{v(H)+p(H)+2g(H)} (-q)^h$ . The differentials are defined through a restriction of the per-edge maps used in the above complex for the chromatic polynomial  $Z(F, q)$ , so that  $m(\widehat{R}) = 1^{\otimes v(H)} \otimes \widetilde{m} \otimes \widehat{m}_g$  and  $\Delta(\widehat{R}) = 1^{\otimes v(H)} \otimes \widetilde{\Delta} \otimes \widehat{m}_g$ . One then obtains a chain complex  $\widehat{\mathcal{C}}(F)$  which we call the *restricted Bollobás-Riordan chain complex*. We denote the homology of this complex by  $\widehat{H}(\widehat{\mathcal{C}}(F))$ . Just as before we have:

**Theorem 4.2.** *Let  $F$  be a fatgraph and  $\widehat{\mathcal{C}}(F)$  its restricted Bollobás-Riordan chain complex. Then the Euler characteristic of the homology  $\widehat{H}(\widehat{\mathcal{C}}(F))$  is equal to the restricted Bollobás-Riordan polynomial  $\widehat{R}(F)$ .*

The following universal coefficient type theorem relates the categorification of the chromatic polynomial above to this categorification of the restricted Bollobás-Riordan polynomial  $\widehat{R}(F)$ . It follows from proposition 4.5.

**Proposition 4.3.** *Let  $F$  be a fatgraph. Then*

$$\begin{aligned}
&\widetilde{H}_j^i(\widetilde{\mathcal{C}}(F)) \\
&= \bigoplus_{p+q=j} \left( \left( \widehat{H}_p^i(\widehat{\mathcal{C}}(F)) \otimes \bar{\Delta}(R_q^{\otimes(i-1)}) \right) \oplus \left( Z_p^i(\widehat{\mathcal{C}}(F)) \otimes R_q^{\otimes i} / \bar{\Delta}(R_q^{\otimes(i-1)}) \right) \right),
\end{aligned}$$

where  $M_p$  denotes the degree  $p$  part of a graded module  $M$ ,  $\widehat{H}(\widehat{\mathcal{C}}(F))$  denotes the homology of the restricted Bollobás-Riordan complex and  $Z_p^i(\widehat{\mathcal{C}}(F)) = \ker(d^i(\widehat{\mathcal{C}}(F))_p)$  are its cycles. Moreover  $H_j^{i+e(F)}(\widehat{\mathcal{C}}(F)) = H_j^i(\mathcal{C}(F))$ .

We also note that the following deletion-contraction exact sequence holds for the homology of the restricted Bollobás-Riordan complex.

**Theorem 4.4.** *Let  $e$  be a bridge of a fatgraph  $F$  and let  $F - e$  denote  $F$  with the bridge  $e$  deleted and  $F/e$  denote  $F$  contracted along the edge  $e$ , then there exists a deletion-contraction exact sequence*

$$\begin{array}{ccc} \widehat{H}^*(\widehat{\mathcal{C}}(F - e)) & \longrightarrow & \widehat{H}^*(\widehat{\mathcal{C}}(F)) \\ & \searrow & \swarrow \\ & \widehat{H}^*(\widehat{\mathcal{C}}(F/e)) \otimes V & \end{array}$$

The proof of this is similar to the proof of property (5) of Proposition 3.1 and is therefore excluded.

**4.2. Categorification of the Multivariate Bollobás-Riordan Polynomial.** Notice that if in the proof of Lemma 4.1 we set  $z = (r + r^{-1})/(q + q^{-1})^{\frac{1}{2}}$  instead of 1 we obtain the two variable polynomial  $\sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{v(H)+p(H)} (-q)^{e(F)-e(H)} (r + r^{-1})^{2g(H)}$ .

We modify it to the three variable polynomial

$$R'(F, q, r, s) =$$

$$\sum_{H \in \mathcal{S}(F)} (q + q^{-1})^{v(H)+p(H)} (-q)^{e(F)-e(H)} (r + r^{-1})^{2g(H)} (1 + s^{-2})^{e(F)-e(H)}.$$

It is straightforward to check that  $R'$  is equivalent to the full Bollobás-Riordan polynomial (1).

This polynomial can be categorified using a straight forward modification of the construction in Section 2. Essentially all this involves is replacing the graded modules in the construction of the chain complex with trigraded modules. To do this define  $V$ ,  $U$  and  $R$  to be the trigraded free modules of rank two with generators in degrees  $(\pm 1, 0, 0)$ ,  $(0, \pm 1, 0)$  and  $(0, 0, -2)$  and  $(0, 0, 0)$  respectively.

Then assign the module  $V^{\otimes v(H)+p(H)} \otimes U^{\otimes 2g(H)} \otimes R^{\otimes h(H)} \{h(H)\}$  to each state. The per-edge maps are defined as before but acting in the relevant

grading. Finally defining the graded dimension as

$$\text{qdim}(M) = \sum_{i,j,k} q^i r^j s^k \text{rk}(M_{(i,j,k)}),$$

we obtain the desired categorification of  $R'(F, q, r, s)$ .

Notice that our chromatic homology is recovered by projecting the arising trigraded homology groups onto a single grading.

The idea of using multi-graded modules was also used in [JHR], to categorify the Tutte polynomial and [Ro] to categorify the Bollobás Riordan polynomial.

In fact, keeping an analogy with Khovanov homology in mind (this will be discussed in detail in the following section), we can extend the function  $R'$  to fat graphs with *signed edges*, that is fatgraphs such that each bridge of  $F$  is decorated with plus or minus sign. Let us reserve the symbol  $F_s$  for signed fatgraphs. For a state  $H$  of a signed fatgraph, let  $e_-(H)$  (respectively  $e_+(H)$ ) denote the number of bridges in  $H$  with a negative (resp. positive) sign. We then define the *height* of a state  $H$  as  $h_s(H) = e_-(F) - e_-(H) + e_+(H)$ . The construction of the homology described above with respect to this new height function, gives a categorification of the polynomial

$$R'(F_s, q, r, s) = (-q - qs^{-2})^{e_-(F_s)} \sum_{H \in \mathcal{S}(F_s)} (q + q^{-1})^{v(H) + p(H)} (r + r^{-1})^{2g(H)} (-q(1 + s^{-2}))^{e_+(H) - e_-(H)}.$$

We note that up to normalization this can be seen to be equivalent to the more simple polynomial

$$\sum_{H \in \mathcal{S}(F)} x^{k(H)} y^{g(H)} \prod_{e \in E(H)} z_e,$$

where  $z_e$  equals  $x^{-1}z$  for an edge of positive weight and  $x^{-1}z^{-1}$  for an edge of negative weight.

In order to write down a universal coefficient theorem for the categorification of the three variable polynomial  $R'(F_s, q, r, s)$  we introduce some notation. Let  $HB(\mathcal{D}(F_s))$  be the homology of the complex  $\mathcal{D}(F_s)$  associated with the three variable Bollobás-Riordan polynomial of signed graphs  $R'$  described above. Clearly  $\mathcal{D}(F_s) = (V^{\otimes v} \otimes \widehat{\mathcal{D}}^i(F_s) \otimes R^{\otimes i}, 1^{\otimes v} \otimes \widehat{d}^i \otimes \bar{\Delta}_i)$ .

Let  $\widehat{HB}^*$  denote the homology of the subcomplex  $(\widehat{\mathcal{D}}(F_s)^i, \widehat{d}^i)$ . Its Euler characteristic is equal to the polynomial

$$\widehat{R}'(F_s, q, r) = \sum_{H \in \mathcal{S}(F_s)} (q + q^{-1})^{p(H)} (r + r^{-1})^{2g(H)} (-q)^{h_s(H)}.$$

**Proposition 4.5.** *Let  $F_s$  be a fatgraph. Then*

$$\begin{aligned} HB_{(j,k,l)}^i(\mathcal{D}(F_s)) &= \bigoplus_{p+q=j} \left( \left( \widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes \overline{\Delta}(R_{(0,0,l)}^{\otimes(i-1)}) \right) \right. \\ &\quad \left. \oplus \left( \widehat{Z}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes R_{(0,0,l)}^{\otimes i} / \overline{\Delta}(R_{(0,0,l)}^{\otimes(i-1)}) \right) \otimes V_{(q,0,0)}^{\otimes v} \right), \end{aligned}$$

where  $M_{(j,k,l)}$  denotes the degree  $(j, k, l)$  part of a trigraded module  $M$  and  $\widehat{Z}_{(j,k,l)}^i(\widehat{\mathcal{D}}(F_s)) = \ker(\widehat{d}_{(j,k,l)}^i)$  are the cycles of  $\widehat{\mathcal{D}}(F_s)$ .

*Proof.* Let us consider the subcomplex  $\mathcal{D}'(F_s) = (\widehat{\mathcal{D}}^i(F_s) \otimes R^{\otimes i}, \widehat{d}^i \otimes \overline{\Delta}_i)$ .

Let  $(HB')^*$  denote its homology.

By the universal coefficient theorem (see eg. [Br])

$$HB_{(j,k,l)}^i(\mathcal{D}(F_s)) = \bigoplus_{p+q=j} \left( (HB')_{(p,k,l)}^i(\mathcal{D}'(F_s)) \otimes V_{(q,0,0)}^{\otimes v} \right),$$

and we see that it remains to understand  $(HB')_{(p,k,l)}^i(\mathcal{D}'(F_s))$ . Now let  $\widehat{Z}^i = \ker(\widehat{d}^i)$  and  $\widehat{B}^i = \text{im}(\widehat{d}^{i-1})$ . For convenience we will denote  $\overline{\Delta}_i$  by  $\overline{\Delta}$ . Then since  $\widehat{d}^i$  and  $\overline{\Delta}_i$  are of degree zero we have

$$(HB')_{(p,k,l)}^i(\widehat{\mathcal{D}}(F_s)) = \left( \frac{\widehat{Z}^i \otimes R^{\otimes i}}{\widehat{B}^i \otimes \overline{\Delta}(R^{\otimes(i-1)})} \right)_{(p,k,l)} = \frac{\widehat{Z}_{(p,k,0)}^i \otimes (R^{\otimes i})_{(0,0,l)}}{\widehat{B}_{(p,k,0)}^i \otimes \overline{\Delta}(R^{\otimes(i-1)})_{(0,0,l)}}.$$

Now  $\widehat{Z}_{(p,k,0)}^i$  and  $\widehat{B}_{(p,k,0)}^i$  are free abelian groups with  $\widehat{Z}_{(p,k,0)}^i \subset \widehat{B}_{(p,k,0)}^i$ , and so we have  $\widehat{Z}_{(p,k,0)}^i = \bigoplus_{\alpha \in I} n_\alpha \mathbb{Z}$  and  $\widehat{B}_{(p,k,0)}^i = \bigoplus_{\alpha \in I} m_\alpha \mathbb{Z}$ , where the sum is over the same finite index  $I$  and  $m_\alpha | n_\alpha$  for each  $\alpha \in I$ . Also it is easy to show that  $(R^{\otimes(i-1)})_{(0,0,l)} = \mathbb{Z}^{\oplus N}$  and  $\overline{\Delta}(R^{\otimes(i-1)})_{(0,0,l)} = \mathbb{Z}^{\oplus M}$  for some  $M$  and  $N$  (i.e. the generators of these groups are 1). We can then write

$$\frac{\widehat{Z}_{(p,k,0)}^i \otimes (R^{\otimes i})_{(0,0,l)}}{\widehat{B}_{(p,k,0)}^i \otimes \overline{\Delta}(R^{\otimes(i-1)})_{(0,0,l)}} = \frac{(\bigoplus_{\alpha \in I} n_\alpha \mathbb{Z}) \otimes \mathbb{Z}^{\oplus N}}{(\bigoplus_{\alpha \in I} m_\alpha \mathbb{Z}) \otimes \mathbb{Z}^{\oplus M}},$$

where  $M|N$ . Using standard properties of the tensor product, we see that this can be written as

$$\bigoplus_{\alpha \in I} \frac{(n_\alpha \mathbb{Z})^{\oplus N}}{(m_\alpha \mathbb{Z})^{\oplus M}} = \bigoplus_{\alpha \in I} \frac{(n_\alpha \mathbb{Z})^{\oplus (M+P)}}{(m_\alpha \mathbb{Z})^{\oplus M}} = \left( \bigoplus_{\alpha \in I} \frac{n_\alpha \mathbb{Z}}{m_\alpha \mathbb{Z}} \right)^{\oplus M} \oplus \left( \bigoplus_{\alpha \in I} n_\alpha \mathbb{Z} \right)^{\oplus P},$$

which by definition is equal to

$$\begin{aligned} & \left( \widehat{Z}_{(p,k,0)}^i / \widehat{B}_{(p,k,0)}^i \right)^{\oplus M} \oplus \left( \widehat{Z}_{(p,k,0)}^i \right)^{\oplus P} = \\ & \left( \widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \right)^{\oplus M} \oplus \left( Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \right)^{\oplus P} \\ & = \left( \widehat{HB}_{(p,k,0)}^i(\widetilde{\mathcal{D}}(F_s)) \otimes \mathbb{Z}^{\oplus M} \right) \oplus \left( Z_{(p,k,0)}^i(\widetilde{\mathcal{D}}(F_s)) \otimes \mathbb{Z}^{\oplus P} \right). \end{aligned}$$

Now since  $R_{(0,0,l)}^{\otimes i} = \mathbb{Z}^{\oplus N}$  and  $\bar{\Delta}(R_{(0,0,l)}^{\otimes (i-1)}) = \mathbb{Z}^{\oplus M}$  we have that  $R_{(0,0,l)}^{\otimes i} / \bar{\Delta}(R_{(0,0,l)}^{\otimes (i-1)}) = \mathbb{Z}^{\oplus (N-M)} = \mathbb{Z}^{\oplus P}$ , we see that the above is equal to

$$\left( \widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F)) \otimes \bar{\Delta}(R_{(0,0,l)}^{\otimes (i-1)}) \right) \oplus \left( Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F)) \otimes R_{(0,0,l)}^{\otimes i} / \bar{\Delta}(R_{(0,0,l)}^{\otimes (i-1)}) \right),$$

as required.  $\square$

Note that in the above proof we do need to be careful about the order of the summands in the direct sums.

Proposition 4.3 is a corollary of this result by projecting the trigraded modules onto a single graded dimension as follows.

*Proof of Proposition 4.3.* Regard the fatgraph  $F$  as a signed fatgraph  $F_s$  by assigning a negative sign to each edge. We have

$$\widetilde{H}_j^i(\widetilde{\mathcal{C}}(F)) = \bigoplus_{p+q+r=j} HB_{(p,q,r)}^i(\mathcal{D}(F_s)).$$

An application of the above proposition gives

$$\begin{aligned} & \bigoplus_{p+q+r=j} \bigoplus_{s+t=p} \left( \left( \widehat{HB}_{(s,q,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes \bar{\Delta}(R_{(0,0,r)}^{\otimes (i-1)}) \right) \oplus \right. \\ & \left. \left( Z_{(s,q,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes R_{(0,0,r)}^{\otimes i} / \bar{\Delta}(R_{(0,0,r)}^{\otimes (i-1)}) \right) \otimes V_{(t,0,0)}^{\otimes v} \right). \end{aligned}$$

Projection onto a single graded variable then gives

$$\bigoplus_{p+q+r=j} \bigoplus_{s+t=p} \left( \left( \widehat{HB}_{s+q}^i(\widehat{\mathcal{D}}(F_s)) \otimes \bar{\Delta}(R_r^{\otimes (i-1)}) \right) \oplus \right.$$

$$\left( Z_{s+q}^i(\widehat{\mathcal{D}}(F_s)) \otimes R_r^{\otimes i} / \bar{\Delta}(R_r^{\otimes(i-1)}) \right) \otimes V_t^{\otimes v}.$$

Which after reindexing gives the required formula.  $\square$

The convenience of our homology theory for the Bollobás-Riordan polynomial of weighted graphs is described in the following theorem.

**Theorem 4.6.** *The Khovanov categorification of the Jones polynomial as well as the categorification of the chromatic polynomial described in the previous section may be recovered from the categorification of  $R'$  just described.*

*Proof.* We begin by using proposition 4.5 to determine the homology  $\widehat{HB}(\widehat{\mathcal{D}}(F_s))$  from  $HB(\mathcal{D}(F_s))$ . To do this suppose that  $r_{-2}$  and  $r_0$  are the generators of  $R$  in graded dimension  $(0, 0, -2)$  and  $(0, 0, 0)$  respectively.

First, consider the map  $f$  defined by  $r_{-2} \mapsto 0$ . Applying this to the homology gives

$$\begin{aligned} & f \left( HB_{(j,k,l)}^i(\mathcal{D}(F_s)) \right) \\ = & \bigoplus_{p+q=j} \left( \left( \widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes \mathbb{Z} \right) \oplus \left( Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes (\mathbb{Z}/\mathbb{Z}) \right) \right) \otimes V_{(q,0,0)}^{\otimes v}, \\ = & \bigoplus_{p+q=j} \left( \left( \widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \oplus Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \right) \otimes V_{(q,0,0)}^{\otimes v} \right), \end{aligned}$$

since only basis elements in graded degree  $(j, k, 0)$  are not killed by  $f$ . Secondly, notice that  $R_{(0,0,-2i)}^{\otimes i} = \mathbb{Z}$  and  $\bar{\Delta}(V^{\otimes(i-1)})_{(0,0,-2i)} = 0$ , therefore proposition 4.5 also gives

$$HB_{(j,k,-2i)}^i(\mathcal{D}(F_s)) = \bigoplus_{p+q=j} Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s)) \otimes \mathbb{Z} \otimes V_{(q,0,0)}^{\otimes v}$$

Finally, since we know each free module  $V_{(q,0,0)}^{\otimes v}$  we know  $Z_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s))$  and hence  $\widehat{HB}_{(p,k,0)}^i(\widehat{\mathcal{D}}(F_s))$ .

The Khovanov homology of an associated link can be recovered from this as in the discussion in Section 5.1: associate the crossing  $\times$  to a edge of negative weight and  $\times$  to an edge of positive weight. The height then equals the number of 1-smoothings of the associated link. The Khovanov homology is then a normalization of  $\widehat{HB}$  after projection onto a single grading.

The second statement follows by regarding an unsigned fatgraph as a signed fatgraph whose edges all have negative weight and projecting onto a single graded dimension.  $\square$



## 5. INDEPENDENCE OF PLANAR EMBEDDINGS


We begin this section by describing the relationship of our homology theory with Khovanov's categorification of the Jones polynomial ([Kh]). We will use this relation to prove the following theorem in Subsection 5.2.

**Theorem 5.1.** *Let  $F$  and  $F'$  be two genus 0 fatgraphs with the same associated graph  $G$ , and let  $P$  be the Poincaré polynomial of the homology. Then*









$$P(F) = P(F'),$$

ie. *the Poincaré polynomial is independent of the planar embedding of the graph  $G$ .*

**5.1. The Relation to Khovanov Homology and Knots.** Let  $F$  be a genus  $g$  fatgraph. As mentioned earlier,  $F$  is equivalent to a genus  $g$  surface which we will denote  $\Sigma_g$ . Notice that  $F$  gives rise to an alternating link  $L \subset \Sigma_g \times I$ , and a canonical projection onto  $F$  by associating a crossing  to each bridge . We call this the *associated link*. The following figure shows a fatgraph with one island of degree 4 and two islands of degree 2 and its associated link.



Just as with link projections on  $S^2$ , we can define the notion of a *smoothing* of a crossing. A *0-smoothing* is defined locally on a link by changing a crossing which looks like  $\times$  to look like  $\smile$ , and for a *1-smoothing* replacing it with  $\frown$ . A *state* of a link diagram is what is obtained by smoothing all of the crossings of the link projection. There is a clear correspondence between the states of a fatgraph and the states of the associated link. This is summarized in the table below.

smoothing	ass. link	fatgraph
0	 $\rightarrow$ 	 $\rightarrow$ 
1	 $\rightarrow$ 	 $\rightarrow$ 

From now on we will now restrict ourselves to genus 0 fatgraphs. The associated links can then be regarded as links in  $S^3$ .

The Khovanov homology of a link is constructed in essentially the same way as the complex in Section 2: given a link diagram  $L$  we define the *height*,  $h(S)$ , of a state  $S$  to be the number of 1-smoothings used in its construction and we let  $p(S)$  denote the number of cycles in the state  $S$ . The chain modules are constructed by assigning the graded module  $V^{\otimes p(S)}\{h(S)\}$ , which has graded dimension  $q^{h(S)}(q+q^{-1})^{p(S)}$ . Again the  $i$ -th chain group is defined to be the direct sum of all modules assigned to states of height  $i$ . Just as in Section 2, one can move from a state of height  $i$  to  $i+1$  by merging or splitting cycles and we use the per-edge maps  $\tilde{m}$  and  $\tilde{\Delta}$  from Section 2 to move between the corresponding modules. The differentials are then obtained by summing over all of the per-edge maps of the appropriate height as before. This gives a chain complex  $\tilde{\mathcal{C}}(L) = (\tilde{\mathcal{C}}^i(L), \tilde{d}^i)$ . Let  $\widetilde{HK}(L)$  denote its homology.

If  $n_{\pm}$  denotes the number of  $\pm$ -crossings of  $L$  (see Subsection 5.2), then the Khovanov complex is defined as the following normalization of  $\tilde{\mathcal{C}}$ ,

$$\mathcal{C}(L) = (\mathcal{C}^i(L), d^i) = \tilde{\mathcal{C}}(L)[-n_-]\{n_+ - 2n_-\} = (\tilde{\mathcal{C}}^i(L), \tilde{d}^i)[-n_-]\{n_+ - 2n_-\}.$$

The homology of this complex is called *Khovanov homology*,  $HK(L)$ . It can be shown ([Kh, BN1]) that the Euler characteristic  $\chi(HK(L))$  is equal to the Jones polynomial of  $L$  and that the homology itself is a knot invariant which is strictly stronger than the Jones polynomial.

Notice that by the correspondence between the states of a fatgraph and of an associated link described above, we have

$$\tilde{\mathcal{C}}(F) = \left( V^{\otimes v} \otimes \tilde{\mathcal{C}}^i(L) \otimes R^{\otimes h}, 1^{\otimes v} \otimes \tilde{d}^i \otimes \bar{\Delta}_h \right),$$

where  $L$  is the associated link of a planar fatgraph.

This observation leads to the following theorem, which is a corollary of Theorem 4.3.

**Theorem 5.2.** *Let  $F$  be a genus 0 fatgraph and  $L \subset S^3$  be the associated link with an arbitrary orientation. Then*

$$H_j^i(\tilde{\mathcal{C}}(F)) =$$

$$\bigoplus_{p+q+r=j} \left( \left( \left( \widetilde{HK}_p^i(L) \otimes \bar{\Delta} R_q^{\otimes(i-1)} \right) \oplus \left( Z_p^i(L) \otimes R_q^{\otimes i} / \bar{\Delta} R_q^{\otimes(i-1)} \right) \right) \otimes V_r^{\otimes v} \right),$$

Where  $M_p$  denotes the degree  $p$  part of a graded module  $M$ ,

$$Z_p^i(L) = \ker(\tilde{d}_p^i) \quad \left( = \ker \left( d_{p+2n_- - n_+}^{i+n_-} \right) \right)$$

are cycles determined by the Khovanov complex and for the planar fatgraph  $F_s$  corresponding to  $L$ ,


$$\widetilde{HK}_p^i(L) = HK_{p+2n_- - n_+}^{i+n_-}(L),$$

where  $HK$  denotes Khovanov homology. Moreover  $H_j^i(\mathcal{C}) = H_j^{i+e(F)}(\tilde{\mathcal{C}})$ .


Note that one can prove an analogous result for the full (ie. without the restriction to alternating links) Jones polynomial using the homology of the signed Bollobás-Riordan polynomial.

**5.2. Proof of Theorem 5.1.** Our technique is to reduce the problem to one of knot theory.

First we need to understand how the two fatgraphs  $F$  and  $F'$  and their corresponding links are related. For this we find it convenient to switch to the language of embedded graphs. Recall that a genus 0 fatgraph is equivalent to an embedding of the associated graph  $G \hookrightarrow S^2$ . We will need the following two local moves on embedded graphs. Let  $G \subset S^2$  be a connected embedded graph. A *1-flip* is a move which replaces a 1-connected component of the map  $G$  with its rotation by  $\pi$  around the axis in the  $xy$ -plane which intersects the 1-connecting vertex. For example,



A *2-flip* is a move which replaces a 2-connected component of  $G$  with its rotation by  $\pi$  around the axis determined by its 2-connecting vertices. For example,




The following theorem relates two planar embeddings of the same graph (see [MT], [MM]).

**Theorem 5.3.** *Let  $G$  be a connected graph and  $f, f' : G \rightarrow S^2$  be two planar embeddings. Then  $f(G)$  and  $f'(G)$  are related by a sequence of 1-flips and 2-flips.*


We need to understand how a flip changes the associated link. First consider the 1-flip. By regarding a 1-connected graph as two components with a vertex identified, it is easy to see that the associated link is non-prime with a connect sum determined by the connecting vertex, and that the embeddings of the two components determine where the connect sum occurs. (Recall that the sum of two links is the link formed cutting open an arc of each link and identifying the free ends in a way consistent with orientation. Although this process is well defined with respect to isotopy for knots, in general it will depend upon which components of the link have been identified under the sum.)

It is then clear that in terms of the associated link, a 1-flip simply changes the way we connect the two links in the connect sum. We will discuss this in more detail in the proof of Lemma 5.7.

Now, by considering the example



it is easy to see that a 2-flip corresponds to a move which replaces a 2-tangle with its rotation by  $\pi$  around the axis between the tangle ends:



This is a form of Conway mutation of the link ([Co]). We refer to this specific move on the link simply as *mutation*.

This discussion gives the following lemma.


**Lemma 5.4.** *Two genus 0 fatgraphs have the same associated connected graph if and only if their associated links are obtained as the connect sum of the same set of links and by a sequence of mutations.*

So far we have only discussed un-oriented links, but our application of knot theory requires a choice of orientation of the links. We need to be careful in this choice. Recall that the *sign* of a crossing of an oriented link is the assignment of  $\pm 1$  according to the following scheme:



If the components of a link  $L$  are labelled  $\{1, \dots, n\}$ , then the *linking number*  $lk_L(i, j)$  is defined to be the sum of the signs over all crossings between the components labelled  $i$  and  $j$ .

Given two links which are related as in lemma 5.4, we require that the corresponding crossings in the two links have the same sign. To do this we orient the summands of links arbitrarily. First we deal with the case of a connected sum. If a connect sum requires the reversal of the orientation of a component, then we reverse all of the components of that summand. The case for mutation is a little more complicated. Let  $R$  denote the tangle on which the mutation acts. Then, if the two free arcs at the top  $R$  are both oriented into or out of the tangle, we retain all orientations. If one of the arcs at the top  $R$  is oriented into the tangle and one out of the tangle, then we reverse the orientations of *all* the components of the tangle  $R$ . See the figure below. We call such an orientation of the mutant the *induced orientation*. It is immediate that the sign of each crossing before and after the mutation is the same.



Further to this, the labeling of  $R$  induces a labeling of the components after the mutation which we call the *induced labeling*. With this information we can now prove the theorem.

*Proof of Theorem 5.1.* Let  $f(G)$  and  $f'(G)$  be the embeddings corresponding to the associated graphs  $F$  and  $F'$ . The following lemma reduces the problem.

**Lemma 5.5.** *Let  $L$  and  $L'$  be the two links associated with the fatgraphs  $F$  and  $F'$ . Then to prove the theorem, it is enough to show that  $P(HK(L)) = P(HK(L'))$ .*

*Proof.* Assume  $P(HK(L)) = P(HK(L'))$ . Recall that if  $H = (H^i)_{i \in \mathbb{Z}}$  is the homology of some chain complex of graded  $\mathbb{Z}$ -modules, the Poincaré polynomial is defined by

$$P(H) = \sum_{i \in \mathbb{Z}} t^i \text{qdim}(H^i).$$

The Poincaré polynomial encodes all of the torsion-free information of the homology groups. By Theorem 5.2, to prove the lemma it suffices to show that  $\text{rk}(\widetilde{HK}(L)_j^i) = \text{rk}(\widetilde{HK}(L')_j^i)$ , and that  $\text{rk}(\ker(\tilde{d}_j^i)) = \text{rk}(\ker(\tilde{d}_j^i))$ .

Clearly, in any graded degree we have

$$\text{rk}(HK_j^i) = \text{rk}(\ker(d_j^i)) - \text{rk}(\text{im}(d_j^{i-1})),$$

and by classic linear algebra we know that

$$\text{rk}(\mathcal{C}_j^i) = \text{rk}(\ker(d_j^i)) + \text{rk}(\text{im}(d_j^i)).$$

Suppose we know the ranks of  $\mathcal{C}_j^k$  and  $HK_j^k$  for some  $k$ . If in addition to this we know the value  $\text{rk}(\text{im}(d_j^{k-1}))$ , then by the above we can determine the values of  $\text{rk}(\ker(d_j^k))$  and  $\text{rk}(\text{im}(d_j^k))$ . Therefore if we know the ranks of each  $\mathcal{C}_j^i$  and  $HK_j^i$  and one value of  $\text{rk}(\text{im}(d_j^{k-1}))$  then we can determine every value  $\text{rk}(\ker(d_j^i))$  and  $\text{rk}(\text{im}(d_j^i))$  and we then know the torsion-free information of the entire complex.

Now it is easily seen (recall from the introduction that  $v+p+2g-e=2k$ ) that each term  $\text{rk}(\mathcal{C}_j^i)$  is equal for the two chain complexes from the two associated links  $L$  and  $L'$ . Also by assumption we have that each value of  $\text{rk}(HK_j^i)$  is the same in the two complexes. The above argument then tells us that if there exists a value  $\text{rk}(\text{im}(d_l^{k_l}))$  which is equal for the two complexes associated with  $L$  and  $L'$  for each graded dimension  $l$ , then the torsion-free parts of the two complexes are equal. Finally since the chain complexes in any graded degree only have finitely many non-zero terms clearly such values  $\text{rk}(\text{im}(d_l^{k_l})) = 0$  exist, completing the proof of the lemma.  $\square$

We need to use the following theorem of Lee, which proved a result conjectured by Garoufalidis in [Ga].

**Theorem 5.6** ([Le2]). *For a reduced alternating link,  $P(HK)$  is determined by the Jones polynomial, the signature of the link and the linking numbers.*

Clearly the associated links are alternating. Since we are only interested in determining  $P(HK)$  and Khovanov homology is a link invariant, we can reduce the associated link (recall that a link projection is said to be *reduced* if four distinct regions meet at every crossing). It is well known and easily seen that Conway mutation does not change the Jones polynomial or the signature of the link (for a definition of the signature of a link see, for example, [Li]) and since the Jones polynomial is multiplicative and the signature additive under the connect sum, we see that these two invariants are equal for both our associated links. However, the linking numbers do change under the operations. By Theorems 1.2 and 4.5 of [Le1] and Corollary A.2 of [Ga], we see that it is enough to show that if the components of each of the associated links are labelled  $1, \dots, n$  then the following formula is equal for both links,

$$(6) \quad \sum_{E \subset \{2, \dots, n\}} \gamma_{\sum_{j \in E, k \notin E} lk_{jk}}.$$

The following two lemmas will complete the proof of Theorem 5.1.

**Lemma 5.7.** *Equation 6 does not depend upon which two components the connect sum operation acts.*

*Proof.* Suppose we have two links  $L$ , with components labeled  $a_1, \dots, a_n$ , and  $L'$  with components labeled  $b_1, \dots, b_m$ . Let  $A$  be the link obtained by connect summing with respect to the components  $a_1$  and  $b_1$  labeling this new component  $\alpha$  and  $B$  be the link obtained by connect summing with respect to the components  $a_2$  and  $b_1$  labelling this new component  $\beta$ . If  $n \leq 2$  and  $m = 1$  the result is obvious, so assume that this is not the case. Now if  $E \subset \{\alpha, a_2, \dots, a_n, b_2, \dots, b_m\}$  then construct a subset  $F$  from  $E$  by replacing the element  $\alpha$  with  $a_1$  and  $a_2$  by  $\beta$ , if  $\alpha$  or  $a_2$  are in  $E$ . Similarly if  $E \subset \{\beta, a_1, a_3, \dots, a_n, b_2, \dots, b_m\}$  then construct a subset  $F$  from  $E$  by replacing the element  $\beta$  with  $a_2$  and  $a_1$  by  $\alpha$ , if  $\beta$  or  $a_1$  are in  $E$ . Then since

$$\begin{aligned} lk_A(\alpha, a_i) &= lk_L(a_1, a_i), & lk_A(\alpha, b_i) &= lk_{L'}(b_1, b_i), \\ lk_A(a_i, a_j) &= lk_L(a_i, a_j), & lk_A(b_i, b_j) &= lk_{L'}(b_i, b_j) \end{aligned}$$

and

$$lk_B(\beta, a_i) = lk_L(a_2, a_i), \quad lk_B(\beta, b_i) = lk_{L'}(b_1, b_i),$$

$$lk_B(a_i, a_j) = lk_L(a_i, a_j), \quad lk_B(b_i, b_j) = lk_{L'}(b_i, b_j)$$

and all other linking numbers are zero, we have that


$$\gamma_{\sum_{j \in E, k \notin E} lk_{jk}} = \gamma_{\sum_{j \in F, k \notin F} lk_{jk}},$$

and therefore for each summand in the equation 6 for the link  $A$ , there is a corresponding summand of equal value in the equation for the link  $B$  and vice-versa. This completes the proof of Lemma 5.7  $\square$

Note that by our choice of orientation of the summands, to prove the theorem we are allowed to assume that the connect sum is consistent with the orientation.

**Lemma 5.8.** *If  $L$  is a link,  $L'$  is obtained from  $L$  by a mutation and  $L'$  has the canonical orientation and labeling, then the sum of formula 6 is equal for the two links  $L$  and  $L'$ ,*

*Proof.* We can regard the link  $L$  as the identification of two 2-tangles,  $R$  and  $T$ . We may assume that  $L'$  is obtained from  $L$  by a mutation which ‘flips over’ the tangle  $R$ . Then the two links  $L$  and  $L'$  and their linking numbers differ according to how the two tangles  $R$  and  $T$  are joined. This can be represented by the following figure.



The eight free ends of the tangles belong to one or two components of the final link. The remainder of the proof is split into several cases according to which components the free ends will belong to in the corresponding link. Note that the number of components of  $L$  and  $L'$  will always be equal.

Now suppose that the free ends of the tangle are labelled as in the above figure. First note that only the linking numbers which involve a component coming from one of the free ends can change under mutation.

**Case 1.** Suppose that after identification all of the free ends belong to one component labelled  $A$ . Then

$$lk_L(A, i) = lk_{L'}(A, i),$$

for an arbitrary component  $i$ . Therefore Lee's formula is unchanged by mutation.

**Case 2.** Suppose that after identification  $x_1, x_2, \alpha_1$  and  $\alpha_2$  belong to the same component labelled  $A$  and  $x_3, x_4, \alpha_3$  and  $\alpha_4$  belong to the same component labelled  $B$ . Then

$$\begin{aligned} lk_L(A, i) &= lk_{L'}(A, i) \\ lk_L(B, i) &= lk_{L'}(B, i) \end{aligned} \quad ,$$

for an arbitrary component  $i$ , and again Lee's formula is unchanged by mutation.

**Case 3.** Suppose that after identification  $x_1, \alpha_2, x_3$  and  $\alpha_4$  belong to the same component labelled  $A$  and  $x_2, \alpha_1, x_4$  and  $\alpha_3$  belong to the same component labelled  $B$ . This case is more complicated. We start with some notation. Let  $A_R$  be the segment of  $A$  in  $R$  with open ends  $x_1, x_3$  and let  $A_T, B_R, B_T$  be defined analogously. Hence the components  $A_R A_T, B_R B_T$  of  $L$  are transformed into components  $A_R B_T, B_R A_T$  of  $L'$ . If  $n = 2$  then the components of  $L$  are exactly  $A, B$  and Lemma is simply true. Hence assume further that  $n > 2$  and  $l_1 = C$  is a component of  $L$  different from both  $A, B$ . Without loss of generality assume  $C$  is a component of  $R$ . Let us denote by  $\mathcal{L}_R$  ( $\mathcal{L}_T$ ) the set of the components of  $L$  in  $R$  ( $T$  respectively).

Let  $\mathcal{E}(L)$  be the set of all  $E \subset \{2, \dots, n\}$  such that  $A \in E$  and  $B \notin E$ , or  $A \notin E$  and  $B \in E$ . Analogously let  $\mathcal{E}(L')$  be the set of all  $E \subset \{2, \dots, n\}$  such that  $A_R B_T \in E$  and  $B_R A_T \notin E$ , or  $A_R B_T \notin E$  and  $B_R A_T \in E$ .

For  $E \in \mathcal{E}(L)$  let

$$l(E) = \sum_{i \in \{A, B\} \cap E, j \notin E \cup \{A, B\}} lk_L(i, j) + \sum_{i \in E - \{A, B\}, j \in \{A, B\} - E} lk_L(i, j).$$

Analogously for  $E \in \mathcal{E}(L')$  let

$$\begin{aligned} l(E) &= \sum_{i \in \{A_R B_T, B_R A_T\} \cap E, j \notin E \cup \{A_R B_T, B_R A_T\}} lk_{L'}(i, j) + \\ &\quad \sum_{i \in E - \{A_R B_T, B_R A_T\}, j \in \{A_R B_T, B_R A_T\} - E} lk_{L'}(i, j). \end{aligned}$$

Clearly, it suffices to show:

**Claim 1.**

$$\{l(E); E \in \mathcal{E}(L)\} = \{l(E); E \in \mathcal{E}(L')\}.$$

For each  $X \subset \mathcal{L}_R$  containing  $C$  and for each  $Y \subset \mathcal{L}_T$  let  $\mathcal{E}(L, A_R, X, Y) = \{E \in \mathcal{E}(L); E = \{A_R A_T\} \cup X \cup Y\}$  and  $\mathcal{E}(L', A_R, X, Y) = \{E \in \mathcal{E}(L'); E = \{A_R B_T\} \cup X \cup Y\}$ . We define  $\mathcal{E}(L, B_R, X, Y), \mathcal{E}(L', B_R, X, Y)$  analogously.

Clearly the sets  $\mathcal{E}(L, A_R, X, Y), \mathcal{E}(L, B_R, X, Y), X \subset \mathcal{L}_R$  containing  $C$  and  $Y \subset \mathcal{L}_T$  form a partition of  $\mathcal{E}(L)$ . An analogous statement holds also for  $L'$ . Hence Claim 1 follows from the following.

**Claim 2.** For each  $X \subset \mathcal{L}_R$  containing  $C$  and  $Y \subset \mathcal{L}_T$ ,

$$\{l(E); E \in \mathcal{E}(L, A_R, X, Y)\} = \{l(E); E \in \mathcal{E}(L, A_R, X, \mathcal{L}_T - Y)\}$$

and the same is true when  $A_R$  is replaced by  $B_R$ .

Claim 2 may be verified by checking. In fact, both sides are equal to

$$\{lk_L(A_R, \mathcal{L}_R - X) + lk_L(B_R, X), lk_L(A_T, \mathcal{L}_T - Y) + lk_L(B_T, Y)\}.$$

This finishes the proof of Claim 2, Claim 1 and Case 3.

**Case 4.** The final case is when, after identification,  $x_1, \alpha_2, x_4$  and  $\alpha_3$  belong to the same component labelled  $A$  and  $x_2, \alpha_1, x_3$  and  $\alpha_4$  belong to the same component labelled  $B$ . This case is similar to Case 3 and the proof is omitted.

This finishes the proof of the lemma. □

This completes the proof of Theorem 5.1. □

*Remark 5.9.* With some easy changes, the proof of Theorem 5.1 above shows that the torsion-free part of the homology  $HB(\mathcal{D}(F_s))$  is also invariant of the choice of genus 0 embedding of a signed fatgraph all of whose edges are of the same sign.

## 6. HELME-GUIZON AND RONG'S CHROMATIC HOMOLOGY

In this final section we show how a categorification of the Bollobás-Riordan polynomial can provide a framework to unite Helme-Guizon and Rong's categorification for the chromatic polynomial ([HGR1]) and Khovanov homology. As mentioned earlier, this addresses a question in [HGR1]. We begin by recalling this categorification of the chromatic polynomial.

Helme-Guizon and Rong categorify the chromatic polynomial  $M(G, 1 + r) = \sum_{H \in \mathcal{S}(G)} (-1)^{e(H)} (1 + r)^{k(H)}$ . This is categorified by considering modules  $M$  which are free, graded, rank 2  $\mathbb{Z}$ -modules with generators  $m_0$  in graded degree zero and  $m_1$  in degree 1, so that  $\text{qdim} = 1 + r$ . The height function is  $|H|$  and the the module  $M^{\otimes k(H)}$  is attached to each state. The per-edge maps are either the identity or the map induced by the degree zero multiplication  $m'(m_0, m_0) = m_0$ , acting on merging connected components.

Consider again the Bollobás-Riordan polynomial

$$R(F, x, y, z) = \sum_H x^{r(F) - r(H)} y^{n(H)} z^{k(H) - p(H) + n(H)}.$$

Using the definitions of the rank and nullity we can write this as

$$x^{-k}(yz)^{-v} \sum_H (xyz^2)^{k(H)} (yz)^{e(H)} z^{-p(H)}.$$

Setting  $x =$ ,  $y =$ ,  $z =$  and forget about normalizations, we obtain the polynomial

$$B(F; q, r) = \sum_H (1+r)^{k(H)} (-q)^{e(H)} (q+q^{-1})^{p(H)}.$$

We concern ourselves with this evaluation of the Bollobás-Riordan polynomial, which we note is not equivalent to the 3-variable Bollobás-Riordan polynomial

Again this polynomial can be categorified. The chain complex is constructed using rank 2, free, bigraded modules  $V$  and  $M$  with basis generators  $v_{\pm}$  in graded degree  $(\pm 1, 0)$  and  $m_0$  and  $m_1$  in graded degrees  $(0, 0)$  and  $(0, 1)$  respectively. The module  $M^{\otimes k(H)} \otimes V^{\otimes p(H)}$  is assigned to each state and the per-edge maps are of the form  $m' \otimes \tilde{m}$  and  $m' \otimes \tilde{\Delta}$ , where  $\tilde{m}$ ,  $\tilde{\Delta}$  and  $m'$  are the obvious bigraded versions of the maps defined in Section 2 and above.

Notice that for a planar fatgraph  $F$ , this chain complex is of the form  $E(F) = (C^i \otimes D^i, d^i \otimes \partial^i)$  where  $C(F) = (C^i, d^i)$  is the chain complex of Helme-Guizon and Rong's chromatic homology and, by Subsection 5.1,  $D(F) = (D^i, \partial^i)$  is the unnormalized Khovanov complex of the reflection (since the height function here corresponds to the addition of edges rather than the removal) of the associated link. Also note that the maps  $f : M \rightarrow 1$  and  $g : V \rightarrow 1$  clearly induce chain maps  $f : E(G) \rightarrow D(G)$  and  $g : E(G) \rightarrow C(G)$ . Putting all of this together we obtain:

**Proposition 6.1.** *For a planar fatgraph  $F$  there is a homology theory for the Bollobás-Riordan polynomial which comes equipped with two natural homomorphisms, one to the chromatic homology of Helme-Guizon and Rong and the other to the Khovanov homology of the reflection of the associated link.*

#### APPENDIX A. TABLE OF POLYNOMIALS

To simplify the entries of the table we write  $v, p, \dots$  for  $v(H), p(H), \dots$  and sums are taken over the appropriate set of states.

polynomial	state sum	complex	homology
$Z(F, q)$	$(-1)^{e(F)} \sum (q + q^{-1})^{v+p+2g} ((-q)(1 + q^{-2}))^h$	$\mathcal{C}(F)$	$H(\mathcal{C}(F))$
-	$\sum (q + q^{-1})^{v+p+2g} ((-q)(1 + q^{-2}))^h$	$\hat{\mathcal{C}}(F)$	$\hat{H}(\hat{\mathcal{C}}(F))$
$\tilde{R}(F, q)$	$\sum (q + q^{-1})^{v+p+2g} (-q)^h$	$\hat{\mathcal{C}}(F)$	$\hat{H}\hat{B}(\hat{\mathcal{C}}(F))$
$R'(F_s, q, r, s)$	$\sum (q + q^{-1})^{v+p} (r + r^{-1})^{2g} (-q(1 + s^{-2}))^{h_s}$	$\mathcal{D}(F_s)$	$HB(\mathcal{D}(F_s))$
$\tilde{R}'(F_s, q, r)$	$\sum (q + q^{-1})^p (r + r^{-1})^{2g} (-q)^{h_s}$	$\hat{\mathcal{D}}(F_s)$	$\hat{H}\hat{B}(\hat{\mathcal{D}}(F_s))$
Jones poly.	$(-1)^{n-} q^{n+} - 2n- \sum (q + q^{-1})^p (-q)^h$	$\mathcal{C}(L)$	$HK(L)$
-	$\sum (q + q^{-1})^p (-q)^h$	$\hat{\mathcal{C}}(L)$	$\hat{H}\hat{K}(L)$

## REFERENCES

- [BN1] D. Bar-Natan, *On Khovanov's categorification of the Jones polynomial*, *Algebr. Geom. Topol* **2** (2002), 337-370, [arXiv:math.QA/0201043](#).
- [Bo] B. Bollobás, *Modern Graph Theory*, Graduate Texts in Mathematics, **184**, Springer-Verlag, New York, 1998.
- [BR] B. Bollobás and O. Riordan, *A polynomial of graphs on surfaces* *Math. Ann.* **323** (2002), no. 1, 81-96.
- [Br] K. S. Brown, *Cohomology of Groups*, Graduate Texts in Mathematics **87** Springer-Verlag, New York-Berlin, 1982.
- [CCR] M. Chmutov, S. Chmutov and Y. Rong, *Knight move for chromatic graph cohomology*, preprint, [arXiv:math.CO/0511598](#).
- [CP] S. Chmutov and I. Pak, *The Kauffman bracket and the Bollobás-Riordan polynomial of ribbon graphs*, preprint, [arXiv:math.GT/0404475](#).
- [CU] M. Chmutov and E. Udovina, *Reduced chromatic graph cohomology*, preprint, [arXiv:math.CO/0510536](#).
- [Co] J. H. Conway, *An enumeration of knots and links, and some of their algebraic properties*, *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, 329-358, Pergamon, Oxford, 1970.
- [DiF] P. Di Francesco, *2D Quantum Gravity, Matrix Models and Graph Combinatorics*, preprint, [arXiv:math-ph/0406013](#).
- [DGR] N. M. Dunfield, S. Gukov and J. Rasmussen, *The superpolynomial for knot homologies*, preprint, [arXiv:math.GT/0505662](#).
- [Ga] S. Garoufalidis, *A conjecture on Khovanov's invariants*, *Fundamenta Mathematicae*, **184** (2004) 99-101.
- [GW] S. Gukov and J. Walcher, *Matrix Factorizations and Kauffman Homology*, preprint, [arXiv:math.hep-th/0512298](#).
- [Ha] A. Hatcher, *Algebraic Topology*, Cambridge University Press, Cambridge, 2002.
- [HGR1] L. Helme-Guizon and Y. Rong, *A categorification for the chromatic polynomial*, *Algebr. Geom. Topol.* **5** (2005) 1365-1388, [arXiv:math.CO/0412264](#).
- [HGR2] L. Helme-Guizon and Y. Rong, *Graph Cohomologies from Arbitrary Algebras*, Preprint, [arXiv:math.QA/0506023](#).
- [HGRP] L. Helme-Guizon, Y. Rong and J. H. Przytycki, *Torsion in Graph Homology*, preprint, [arXiv:math.GT/0507245](#).
- [JHR] E. F. Jasso-Hernandez and Y. Rong, *A Categorification for the Tutte Polynomial*, preprint, [arXiv:math.CO/0512613](#).

- [Kh] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), 359-426, [arXiv:math.QA/9908171](#).
- [KR1] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*, preprint, [arXiv:math.QA/0401268](#).
- [KR2] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology II*, preprint, [arXiv:math.QA/0505056](#).
- [Le1] E. S. Lee *An endomorphism of the Khovanov invariant*, Adv.Math, **197** (2002), 554-586, [arXiv:math.GT/0210213](#).
- [Le2] E. S. Lee *The support of the Khovanov's invariants for alternating knots*, preprint, [arXiv:math.GT/0201105](#).
- [Li] W. B. R. Lickorish, *An introduction to knot theory*, Graduate Texts in Mathematics, **175** Springer-Verlag, New York, 1997.
- [MM] B. Mohar, *Combinatorial local planarity and the width of graph embeddings*, Canad. J. Math. 44-6 (1992), 1272- 1288.
- [MT] B. Mohar and C. Thomassen, *Graphs on surfaces*, Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, 2001.
- [Pr] J. H. Przytycki, *When the theories meet: Khovanov homology as Hochschild homology of links*, preprint, [arXiv:math.GT/0509334](#).
- [Ro] Y. Rong, *A quadruply-graded graph homology for the Bollobas-Riordan polynomial*, Talk at Knots In Washington XXI: Skein modules, Khovanov homology and Hochschild homology, December 9-11, 2005, George Washington University.
- [St1] M. Stošić, *Categorification of the Dichromatic Polynomial for Graphs*, preprint, [arXiv:math.QA/0504239](#).
- [St2] M. Stošić, *New Categorifications of the Chromatic and the Dichromatic Polynomials for Graphs*, preprint, [arXiv:math.QA/0507290](#).

DEPT. OF APPLIED MATHEMATICS AND INSTITUTE OF THEORETICAL COMPUTER SCIENCE, CHARLES UNIVERSITY, MALOSTRANSKÉ N. 25, 118 00 PRAHA 1, CZECH REPUBLIC.  
 and DEPTO. ING. MATEMATICA, UNIVERSITY OF CHILE, CHILE.  
*E-mail address:* [loebl@kam.mff.cuni.cz](mailto:loebl@kam.mff.cuni.cz)

DEPT. OF APPLIED MATHEMATICS AND INSTITUTE OF THEORETICAL COMPUTER SCIENCE, CHARLES UNIVERSITY, MALOSTRANSKÉ N. 25, 118 00 PRAHA 1, CZECH REPUBLIC.  
*E-mail address:* [iain@kam.mff.cuni.cz](mailto:iain@kam.mff.cuni.cz)