

Pointfree aspects of the T_D axiom of classical topology

Bernhard Banaschewski ^{*} Aleš Pultr ^{†‡}

Abstract

Abstraction from the condition defining T_D -spaces leads to the following notion in an arbitrary frame L : a filter F in L is called *slicing* if it is prime and there exist $a, b \in L$ such that $a \notin F$, $b \in F$, and a is covered by b . This paper deals with various aspects of these slicing filters. As a first step, we present several results about the original T_D condition. Next, concerning slicing filters, we show they are completely prime and characterize them in various ways. In addition, we prove for the frames $\mathfrak{O}X$ of open subsets of a space X that every slicing filter is an open neighbourhood filter $\mathcal{U}(x)$ and X is T_D iff every $\mathcal{U}(x)$ is slicing. Further, for \mathbf{Top}_D and \mathbf{Frm}_D the categories of T_D spaces and their continuous maps, and all frames and those homomorphisms whose associated spectral maps preserve the completely prime elements, respectively, we show that the usual contravariant functors between \mathbf{Top} and \mathbf{Frm} induce analogous functors here, providing a dual equivalence between \mathbf{Top}_D and the subcategory of \mathbf{Frm}_D given by the T_D -spatial frames (not coinciding with the spatial ones). In addition, we show that \mathbf{Top}_D is mono-coreflective in a suitable subcategory of \mathbf{Top} . Finally, we provide a comparison between T_D -separation and sobriety showing they may be viewed, in some sense, as mirror images of each other.

^{*}Department of Mathematics and Statistics, McMaster University, 1280 Main St. W, Hamilton, Ontario L8S 4K1, Canada

[†]Department of Applied Mathematics and ITI, MFF, Charles University, CZ 11800 Praha 1, Malostranské nám. 25 pultr@kam.ms.mff.cuni.cz

[‡]Thanks go to the Natural Sciences and Engineering Research Council of Canada and to the project 1M0545 of the Ministry of Education of the Czech Republic.

Introduction

This paper is motivated by the familiar topological separation axiom

T_D : Every point x has an open neighbourhood U such that $U \setminus \{x\}$ is open,

originally introduced by Aull and Thron [1] as a condition notable because it is strictly between T_0 and T_1 . On the other hand, and perhaps more importantly, there is the result that, for T_D -spaces X and Y , $X \cong Y$ whenever their lattices of open sets are isomorphic ([5, 11]). Here, this will be seen as an immediate consequence of the contravariant adjointness we establish between the category of T_D -spaces and their continuous maps, and a suitable subcategory of the category of all frames.

The tool for this is the following general concept, meaningful in pointfree topology, that is, for arbitrary frames L , abstracted from the condition which T_D imposes on the open neighbourhood filters of the points: a filter F in L is called *slicing* if it is prime and there exist $a, b \in L$ such that $a \notin F$, $b \in F$, and a is covered by b . The aim of this paper is to investigate the properties of these slicing filters and their general rôle.

As a first step, we characterize the T_D -points of a space X , that is, the $x \in X$ satisfying the condition which T_D requires of all points, in a variety of ways (2.3.2). Next, turning to the slicing filters in an arbitrary frame, we show they are completely prime (2.6.1) and then characterize them among all completely prime filters in several ways (2.6.2). Further, for the frame $\mathfrak{O}X$ of open subsets of a space X , we show that every slicing filter is the open neighbourhood filter $\mathcal{U}(x)$ of some T_D -point $x \in X$ (2.7.1) and X is T_D iff every $\mathcal{U}(x)$ is slicing (2.7.2).

Next, we consider the category \mathbf{Top}_D of all T_D -spaces and their continuous maps and the category \mathbf{Frm}_D of all frames and those homomorphisms $h : L \rightarrow M$ for which the spectral map $p \mapsto h_*(p)$ takes any *completely prime* $p \in M$ to a *completely prime* element in L (see 1.5, 3.1.1.2; the right adjoint h_* of a standard homomorphism h just takes *primes* to *primes*). We show that there are contravariant functors $\mathbf{Top}_D \rightarrow \mathbf{Frm}_D$ and $\mathbf{Frm}_D \rightarrow \mathbf{Top}_D$, adjoint on the right, restricting the usual functors between \mathbf{Top} and \mathbf{Frm} (3.5.1) which induces a dual equivalence between \mathbf{Top}_D and the subcategory of \mathbf{Frm}_D given by the T_D -spatial frames. In addition, we obtain that \mathbf{Top}_D is monoreflective in the subcategory of \mathbf{Top} given by the continuous maps $f : X \rightarrow Y$ for which $\mathfrak{O}f : \mathfrak{O}Y \rightarrow \mathfrak{O}X$ belongs to \mathbf{Frm}_D (3.7.2).

Finally, we compare T_D -separation with sobriety, based on the relation $R(X, Y)$ between spaces which says that “ X is a proper subspace of Y such that $(U \mapsto U \cap Y) : \mathfrak{O}Y \rightarrow \mathfrak{O}X$ is an isomorphism”. We show that the two properties may be regarded, in some sense, as mirror images of each other ($(X$ is sober iff $\forall Y, \neg R(X, Y)$, Y is T_D iff $\forall X, \neg R(X, Y)$, see 4.3).

1 Preliminaries

1.1. Frames and their homomorphisms. Recall that a *frame* is a complete lattice L satisfying the distributivity law

$$a \wedge \bigvee B = \bigvee \{a \wedge b \mid b \in B\}$$

for all $a \in L$ and $B \subseteq L$. A *frame homomorphism* $h : L \rightarrow M$ preserves arbitrary joins (including the bottom 0) and all *finitary* meets (including the top 1). As usual, the resulting category will be denoted by

Frm.

If X is a topological space we have the frame $\mathfrak{O}X$ of its open sets, and if $f : L \rightarrow M$ is a continuous map then $\mathfrak{O}f = (U \mapsto f^{-1}[U]) : \mathfrak{O}Y \rightarrow \mathfrak{O}X$ is a frame homomorphism. Thus we have a contravariant functor

$$\mathfrak{O} : \mathbf{Top} \rightarrow \mathbf{Frm}.$$

1.2. Primes and complete primes. An element $p \neq 1$ of a frame L is said to be *prime* (alternatively, also called *meet-irreducible*) if

$$\begin{aligned} p = a \wedge b &\Rightarrow p = a \text{ or } p = b \\ \text{(equivalently, } p \geq a \wedge b &\Rightarrow p \geq a \text{ or } p \geq b), \end{aligned}$$

and *completely prime* if

$$p = \bigwedge S \Rightarrow p \in S$$

for any subset $S \subseteq L$.

1.3. Right adjoints to frame homomorphism. Since a frame homomorphism $h : L \rightarrow M$ preserves all suprema it has a right adjoint

$$h_* : M \rightarrow L$$

(satisfying $h(a) \leq b$ iff $a \leq h_*(b)$). The following is a well-known

Fact. *If p is prime in M then $h_*(p)$ is prime in L .*

(Indeed, if $a \wedge b \leq h_*(p)$ then $h(a) \wedge h(b) \leq p$ and hence, say, $h(a) \leq p$ and $a \leq h_*(p)$.)

1.4. Spectrum. Among the various ways of defining the spectrum ΣL of a frame we have the following

$$\Sigma L = (\{p \in L \mid P \text{ prime}\}, \{\Sigma_a \mid a \in L\})$$

where $\Sigma_a = \{p \mid a \not\leq p\}$: further, for a frame homomorphism $h : L \rightarrow M$ we have and the map $\Sigma h = (p \mapsto h_*(p)) : \Sigma M \rightarrow \Sigma L$. The functor

$$\Sigma : \mathbf{Frm} \rightarrow \mathbf{Top}$$

thus obtained and the \mathfrak{D} from 1.1 are adjoint on the right.

1.5. Prime and completely prime filters. A filter F in a frame L is *prime* if it is proper and

$$a \vee b \in F \Rightarrow a \in F \text{ or } b \in F,$$

and *completely prime* if for any subset $S \subseteq L$,

$$\bigvee S \in F \Rightarrow S \cap F \neq \emptyset.$$

1.5.1. For any completely prime filter F we have a prime element

$$a = \bigvee \{x \mid x \notin F\}. \quad (*)$$

It will be called the *associated prime* of F . Note that, on the other hand, if a is a prime element then

$$F = \{x \mid x \not\leq a\} \quad (**)$$

is a completely prime filter and

the formulas () and (**) constitute a one-one correspondence between the prime elements and the completely prime filters in a frame L .*

1.5.2. The fixed filters. We will use the notation

$$\mathcal{U}(x) = \{U \in \mathfrak{O}X \mid x \in U\}$$

for the *fixed filters* in $\mathfrak{O}X$. Note that each $\mathcal{U}(x)$ is completely prime, the associated prime element of $\mathcal{U}(x)$ being $X \setminus \overline{\{x\}}$.

1.6. Special topological spaces. Since the restriction to T_0 -spaces has no impact on the point-free aspects of spaces, that is, their frames of open sets, we assume throughout that all spaces are T_0 .

Recall that a space X is *sober* if the open sets $X \setminus \overline{\{x\}}$ are precisely the prime elements of $\mathfrak{O}X$. Equivalently,

X is sober iff each completely prime filter in $\mathfrak{O}X$ is fixed

(see [6, 5, 7]). A space X is T_D if

$$\forall x \in X \exists \text{ open } U \ni x \text{ such that } U \setminus \{x\} \text{ is open.}$$

This notion was introduced independently in [1] and [5] (see also [11]).

1.7. T_D -spatial frames. Recall that a frame is *spatial* if it is isomorphic with some $\mathfrak{O}X$. If there is a T_D -space X with this property we call L *T_D -spatial*.

For more about frames see [7] or [9].

For the basic categorical facts used here see the standard texts, e.g., [8].

2 T_D -points and slicing filters

2.1. An element a in a poset is said to be *covered* by b , notation

$$a \triangleleft b,$$

if $a < b$ and $a \leq x \leq b$ implies $a = x$ or $x = b$. We say that a is covered if it is covered by some b .

2.1.1. Lemma. 1. *In a semilattice, if a prime element a is covered by b then for each $x > a$ one has $x \geq b$.*

2. In a complete lattice, if a is completely prime then it is covered; further, if a is prime and covered then it is completely prime.

Proof. 1. We have $a \leq b \wedge x \leq b$; by the primeness, $a \neq b \wedge x$, hence $b \wedge x = b$, that is, $b \leq x$.

2. $a < b = \bigwedge \{c \mid a < c\}$ ($a = b$ is excluded by complete primeness). Thus, if $a < c \leq b$ then $b \leq c \leq b$, showing $a < b$.

If $a = \bigwedge S$ and if it is prime and covered by b then $b \leq s$ for any $s \in S$ that is not a , and hence some of the $s \in S$ is equal to a . \square

2.1.2. Lemma. 1. In the frame $\mathfrak{D}X$ we have $U < V$ iff $U = V \setminus \{x\}$ for some $x \in X$.

2. If $X \setminus \overline{\{x\}} < V$ then $V = (X \setminus \overline{\{x\}}) \cup \{x\}$.

Proof. 1. Suppose $x, y \in V \setminus U$, $x \neq y$. Choose an open set W such that, say $x \notin W \ni y$. Then

$$U \subsetneq U \cup (W \cap V) \subsetneq V$$

(since $x \notin U \cup (W \cap V) \ni y$), a contradiction.

2. By 1, $V = (X \setminus \overline{\{x\}}) \cup \{y\}$ with $y \notin X \setminus \overline{\{x\}}$, that is, $y \in \overline{\{x\}}$. Thus, $x \in V$ and hence $(X \setminus \overline{\{x\}}) \cup \{x\} \subseteq V$ so that

$$((X \setminus \overline{\{x\}}) \cup \{x\}) \setminus \{y\} = X \setminus \overline{\{x\}},$$

showing $x = y$, and the statement follows. \square

2.2. A point $x \in X$ is said to be a T_D -point if there is an open $U \ni x$ such that $U \setminus \{x\}$ is open. In particular, then, X is a T_D -space iff all its points are T_D .

2.2.1. Proposition. $x \in X$ is a T_D -point iff the set $(X \setminus \overline{\{x\}}) \cup \{x\}$ is open.

Proof. If $(X \setminus \overline{\{x\}}) \cup \{x\} = U$ is open take $V = X \setminus \overline{\{x\}} = U \setminus \{x\}$. On the other hand, let x be a T_D -point, with $U \ni x$, U open and $U \setminus \{x\}$ open. Then $U \setminus \{x\} = U \setminus \overline{\{x\}}$. Now $(X \setminus \overline{\{x\}}) \cup \{x\}$ is a neighbourhood of all the $y \in X \setminus \overline{\{x\}}$. But it is a neighbourhood of x as well: we have $x \in U = (U \setminus \overline{\{x\}}) \cup \{x\} \subseteq (X \setminus \overline{\{x\}}) \cup \{x\}$. \square

2.2.2. Corollary. 1. If $x \in X$ is a T_D -point then it is a T_D -point in every subspace $Y \subseteq X$ such that $x \in Y$.

2. Consequently the subspace

$$\{x \in X \mid x \text{ is a } T_D\text{-point in } X\} \subseteq X$$

is a T_D -space.

2.3.1. Observation. In any space X , $X \setminus \overline{\{x\}}$ is a prime in $\mathfrak{D}X$.

2.3.2. Proposition. Let x be a point in a space X . Then the following statements are equivalent.

- (1) x is a T_D -point.
- (2) $X \setminus \overline{\{x\}}$ is completely prime in $\mathfrak{D}X$.
- (3) $X \setminus \overline{\{x\}}$ is covered in $\mathfrak{D}X$.

Proof. (1) \Rightarrow (2). Let $X \setminus \overline{\{x\}} = \bigwedge \mathcal{S}$ and let x be a T_D -point. Since $X \setminus \overline{\{x\}}$ is prime then by 2.1.1 and 2.1.2, each $U \in \mathcal{S}$ is either $X \setminus \overline{\{x\}}$ or $(X \setminus \overline{\{x\}}) \cup \{x\}$. But it cannot always be the latter.

(2) \Rightarrow (3) by 2.2.1, 2.

(3) \Rightarrow (1) by 2.2.2. □

2.4. Lemma. Let $a \leq b$ in a distributive lattice and let $a \vee c < b \vee c$. Then $a \vee c \leq b \vee c$.

Proof. Let $a \vee c \leq x \leq b \vee c$. Then $a = a \wedge x \leq b \wedge x \leq b$ and hence either $a = b \wedge x$ and then

$$a \vee c = (b \wedge x) \vee c = (b \vee c) \wedge (x \vee c) \geq (b \vee c) \wedge x = x,$$

or $b = b \wedge x$, that is, $b \leq x$, and since $c \leq x$ we have

$$b \vee c \leq x \leq b \vee c. \quad \square$$

2.5. For an ideal A and a filter B in a distributive lattice L define the filter

$$A \uparrow B = \{x \mid \exists a \in A \exists b \in B, a \vee x \leq b\}$$

and extend the definition for arbitrary subsets A, B by considering $A \uparrow B = \text{Idl}A \uparrow \text{Flt}B$. In particular

$$a \uparrow b = \{a\} \uparrow \{b\} = \{x \mid a \vee x \geq b\}.$$

This operation is due to Ball, [2] (see also [3]).

2.5.1. Observations. 1. $a \uparrow b$ is always non-empty, and it is proper iff $a \not\leq b$.

$$2. a \uparrow b = a \uparrow (a \vee b).$$

Consequently, since we will be interested in proper filters we will always assume that $a \not\leq b$ and in view of 2 we will typically work with $a < b$.

2.5.2. Lemma. *Let F be a completely prime filter in a frame and p its associated prime. Assume $F = a \uparrow b$ so that*

$$x \not\leq p \quad \text{iff} \quad a \vee x \geq b.$$

Then

$$a \uparrow b = p \uparrow (b \vee p)$$

and

$$p \leq b \vee p.$$

Proof. As is familiar, $p \neq 1$ and $a \vee a = a \vee 0 \not\leq b$, and hence $a \notin F$, and

$$a \leq p. \tag{*}$$

Since $p \leq p$ we have

$$p = a \vee p \not\leq b. \tag{**}$$

By Observation 2.5.1, for the first statement it suffices to prove that

$$a \uparrow b = p \uparrow b.$$

If $a \vee x \geq b$ then $p \vee x \geq b$ by (*); if $a \vee x \not\geq b$ then $x \leq p$ and $p \vee x = p \not\leq b$ by (**).

Further, let $p \leq x \leq b \vee p$. If $p < x$ we have $x \not\leq p$ and hence $p \vee x \geq b$, and finally $x = p \vee x \geq b \vee p$. \square

2.6. A *slicing filter* on a frame L is a prime filter F for which there exist $a < b$ such that

$$a \notin F \ni b.$$

2.6.1. Lemma. *Any slicing filter is completely prime.*

Proof. Let $a \notin F \ni b$ for some $a < b$. $\bigvee S \in F$ implies that

$$a \vee (b \wedge \bigvee S) = \bigvee \{a \vee (b \wedge s) \mid s \in S\} \in F.$$

For any $s \in S$ we have

$$a \leq a \vee (b \wedge s) \leq b$$

and hence

- (i) either $a = a \vee (b \wedge s)$, that is, $b \wedge s \leq a$,
- (ii) or $a \vee (b \wedge s) = b$.

If all the s followed (i) we had $b \wedge \bigvee S \leq a$ contradicting $a \notin F$. Thus, there exists an $s \in S$ such that $b = a \vee (b \wedge s) \in F$, and since F is prime and $a \notin F$, $s \geq b \wedge s \in F$. \square

2.6.2. Proposition. *Let F be a prime filter in a frame. Then the following statements are equivalent.*

- (1) F is a slicing filter.
- (2) F is a completely prime filter and the associated prime is covered.
- (3) F is a completely prime filter and the associated prime is completely prime.
- (4) F is a completely prime filter of the form $a \uparrow b$.

Proof. (1) \Rightarrow (2): By 2.6.1, F is completely prime. Let $a < b$, $a \notin F \ni b$, and let p be the element associated with F . Then $p < p \vee b$ since $b \leq p$ contradicts $p \notin F$. On the other hand, $a \leq p$ since $a \notin F$, and we have $p = a \vee p < b \vee p$, and $p < b \vee p$ by lemma 2.4.

(2) \Leftrightarrow (3) by lemma 2.1.1.

(2) \Rightarrow (4): Take an $F = \{x \mid x \not\leq p\}$ with $p < b$. If $x \in T$ (that is, $x \not\leq p$) then $p \vee x$ and b are in F and hence $(p \vee x) \wedge b = b$ (the meet is not $\leq p$) so that $p \vee x \geq b$, and $x \in p \uparrow b$.

On the other hand, if $x \in a \uparrow b (= p \uparrow b, \text{ recall 1.2.3})$ then $x \vee a \geq b \in F$, and as F is prime, $x \in F$.

(4) \Rightarrow (1) immediately follows from 2.2.2. \square

2.6.3. Notes. 1. The statement (4) is a reformulation of a property used in characterizing T_D spaces in [10].

2. The statement (3) restricts the correspondence from 1.5.1:

the formulas () and (**) constitute a one-one correspondence between the completely prime elements and the slicing filters in a frame L .*

2.7.1. Proposition. *In an arbitrary (T_0) space,*

- 1. every slicing filter in $\mathfrak{D}X$ is fixed, and

2. the fixed filter $\mathcal{U}(x)$ is slicing iff there is a $U \in \mathcal{U}(x)$ such that $U \setminus \{x\}$ is open.

Proof. 1. Let W be the prime open set associated with a slicing filter \mathcal{F} . Let $W \triangleleft V$ and let $x \in X$ be the point for which $W = V \setminus \overline{\{x\}}$ (recall 2.1.2, 2.6.2(2)). We have $W = V \setminus \overline{\{x\}}$ and hence $W = V \cap (X \setminus \overline{\{x\}})$ and as it is prime, $W = X \setminus \overline{\{x\}}$. Now

$$\mathcal{F} = \{U \in \mathfrak{O}X \mid U \not\subseteq X \setminus \overline{\{x\}}\} = \{U \mid U \cap \overline{\{x\}} \neq \emptyset\} = \mathcal{U}(x).$$

2. $\mathcal{U}(x)$ is a slicing filter iff there are $V \notin \mathcal{U}(x) \ni U$ such that $V \triangleleft U$. By 2.1.2 this amounts to $V = U \setminus \{x\}$. \square

2.7.2. Corollary. *A space X is T_D iff each fixed filter $\mathcal{U}(x)$ in $\mathfrak{O}X$ is slicing.*

3 T_D -spectrum and T_D -coreflection

3.1. The category of all T_D -spaces and continuous maps will be denoted by

Top_D.

Further, a frame homomorphism $h : L \rightarrow M$ will be called a *D-homomorphism* if for each completely prime p in M , the right adjoint image $h_*(p)$ is completely prime in L . The category of all frames and *D-homomorphisms* will be denoted by

Frm_D.

3.1.1. Remarks. 1. Note that **Top_D** is obtained from **Top**, the category of all topological spaces, by reducing the class of objects and keeping the specification of morphisms intact while in **Frm_D** we keep all the objects and reduce the class of morphisms.

2. As is familiar, for each frame homomorphism h the h_* preserves prime elements. This is because h preserves meets, see 1.3; for the same reason, if h is a complete homomorphism then h_* preserves completely prime elements. This may create the impression that being *D-homomorphism* is a strong requirement akin to completeness, but this is not the case: consider the following examples.

(a) The condition is, of course, satisfied trivially if there are no completely prime elements in M . This is trivial.

(b) A more interesting case is that of regular frames (corresponding to regular spaces: they are characterized by the property that each element x is the join $\bigvee\{y \mid y \prec x\}$ where $y \prec x$ means that $y^* \vee x$ for the pseudocomplement y^* of y) for which we have the following familiar

Fact. *In a regular frame each prime element is maximal, thus covered by 1, and hence completely prime.*

(Indeed, let a be prime and let $a < b = \bigvee\{x \mid x \prec b\}$ by regularity. Then there is an $x \not\leq a$ such that $x^* \vee b = 1$. Now $x \wedge x^* = 0 \leq a$, hence by primeness, $x^* \leq a$, and $b = a \vee b \geq x^* \vee b = 1$.)

Thus,

each frame homomorphism $h : L \rightarrow M$ with regular L is a D -homomorphism.

(c) Another case will be presented right away in 3.2.

3.2. Consider the standard functor $\mathfrak{D} : \mathbf{Top} \rightarrow \mathbf{Frm}$. For a continuous map $f : X \rightarrow Y$ we have $(\mathfrak{D}f)_*(U) = Y \setminus \overline{f[X \setminus U]}$ and hence

$$(\mathfrak{D}f)_*(X \setminus \overline{\{x\}}) = Y \setminus \overline{\{f(x)\}}.$$

By 2.3.2, the completely prime elements in $\mathfrak{D}X$ for a T_D space X are precisely the $X \setminus \overline{\{x\}}$. Consequently, we can view the restriction of \mathfrak{D} to T_D -spaces as a functor

$$\mathfrak{D} : \mathbf{Top}_D \rightarrow \mathbf{Frm}_D.$$

In the sequel the symbol \mathfrak{D} will be used in this sense.

3.3. The space ΦL . For any frame L define ΦL as the subspace of ΣL (recall 1.4) of all *completely prime* elements, with $\Phi_a = \Sigma_a \cap \Phi L = \{p \text{ completely prime} \mid a \not\leq p\}$ as the open sets. In particular then

$$\Phi_0 = \emptyset, \quad \Phi_1 = \Phi L, \quad \Phi_{a \wedge b} = \Phi_a \cap \Phi_b \quad \text{and} \quad \Phi_{\bigvee S} = \bigcup\{\Phi_a \mid a \in S\}. \quad (*)$$

3.3.1. Lemma. *For each $p \in \Phi L$, $\Phi L \setminus \overline{\{p\}} = \Phi_p$.*

Proof. $\Phi L \setminus \overline{\{p\}} = \bigcup\{\Phi_a \mid p \notin \Phi_a\} = \bigcup\{\Phi_a \mid a \leq p\} = \Phi_p$ by (*) in 3.3. \square

3.3.2. Proposition. *ΦL is a T_D -space.*

Proof. Recall 2.2.1. We have to prove that each $\{p\} \cup (\Phi L \setminus \overline{\{p\}})$ ($= \{p\} \cup \Phi_p$ by 3.3.1) is open; that is, it (also) is a neighbourhood of p . Let $p < a$. Then $p \in \Phi_a$. We will show that $\Phi_a \subseteq \{p\} \cup \Phi_p$. Indeed, if $q \notin \{p\} \cup \Phi_p$ then $p < q$, hence by 2.1.1, $a \leq q$, and hence $q \notin \Phi_a$. \square

3.3.3. Furthermore, the correspondence $L \mapsto \Phi L$ extends to a functor

$$\Phi : \mathbf{Frm}_D \rightarrow \mathbf{Top}_D$$

amounting to the restriction of the spectrum functor Σ . That is, $\Phi h(p) = h_*(p)$ and one has

$$(\Phi h)^{-1}[\Phi_a] = \Phi_{h(a)}$$

so that Φh is continuous.

3.4. For any frame L , define a mapping (as in the case of the standard spectrum)

$$\delta_L : L \rightarrow \mathfrak{D}\Phi L$$

by setting $\delta_L(a) = \Phi_a$. By (*) in 3.3, δ_L is a frame homomorphism.

3.4.1. Lemma. *For each completely prime $p \in L$ we have*

$$(\delta_L)_*(\Phi_p) = p.$$

Proof. We have $\Phi_a \subseteq \Phi_p$ iff $a \leq p$: $a \leq p$ implies $\Phi_a \subseteq \Phi_p$, and if $a \not\leq p$ then $p \in \Phi_a \setminus \Phi_p$ and hence $\Phi_a \not\subseteq \Phi_p$.

Now $a \leq (\delta_L)_*(\Phi_p)$ iff $\Phi_a \subseteq \Phi_p$. Hence for any a , $a \leq (\delta_L)_*(\Phi_p)$ iff $a \leq p$, and the statement follows. \square

3.4.2. Lemma. *The δ_L constitute a natural transformation $\delta : \text{Id} \rightarrow \mathfrak{D}\Phi$.*

Proof. By 3.3.1 and 3.4.1 each δ_L is a D -homomorphism. As in the case of the usual spectrum we have $\mathfrak{D}\Phi(h)(\delta_L(a)) = (h_*)^{-1}[\Phi_a] = \Phi_{h(a)} = \delta_M(h(a))$. \square

3.5. For a T_D -space X (where the $X \setminus \overline{\{x\}}$ are completely prime) define

$$\pi_X : X \rightarrow \Phi\mathfrak{D}X, \quad x \mapsto X \setminus \overline{\{x\}}.$$

Proposition. *The mappings π_X constitute a natural equivalence $\pi : \text{Id} \rightarrow \Phi\mathfrak{D}$.*

Proof. First, π_X is onto and one-one since for a T_D -space, $X \setminus \overline{\{x\}}$ are precisely the completely primes, and if $x \neq y$ then $\overline{\{x\}} \neq \overline{\{y\}}$ (our spaces are T_0).

Next, $\pi_X^{-1}[\Phi_U] = \{x \mid U \not\subseteq X \setminus \overline{\{x\}}\} = \{x \mid x \in U\} = U$, proving that π_X is a homeomorphism.

Finally, if $f : X \rightarrow Y$ is a continuous map we have

$$\Phi \mathfrak{D}f(\pi_X(x)) = (\mathfrak{D}f)^*(X \setminus \overline{\{x\}}) = Y \setminus \overline{\{f(x)\}} = \pi_Y(f(x)),$$

that is, π is natural. \square

3.5.1. Proposition. *The contravariant functors $\Phi : \mathbf{Frm}_D \rightarrow \mathbf{Top}_D$ and $\mathfrak{D} : \mathbf{Top}_D \rightarrow \mathbf{Frm}_D$ are adjoint on the right with the adjunction maps $\delta_L : L \rightarrow \mathfrak{D}\Phi L$ and $\pi_X : X \rightarrow \Phi \mathfrak{D}X$ where the latter is a homeomorphism for each X .*

Proof. As in the case of the standard spectrum, consider the composition

$$\mathfrak{D}X \xrightarrow{\delta_{\mathfrak{D}X}} \mathfrak{D}\Phi \mathfrak{D}X \xrightarrow{\mathfrak{D}\pi_X} \mathfrak{D}X.$$

We have

$$\begin{aligned} \mathfrak{D}\pi_X(\delta_{\mathfrak{D}X}(U)) &= \pi_X^{-1}(\Phi_U) = \{x \mid X \setminus \overline{\{x\}} \in \Phi_U\} = \\ &= \{x \mid U \not\subseteq X \setminus \overline{\{x\}}\} = \{x \mid x \in U\} = U. \end{aligned}$$

Similarly for

$$\Phi L \xrightarrow{\pi_{\Phi L}} \Phi \mathfrak{D}\Phi L \xrightarrow{\Phi \delta_L} \Phi L.$$

We have

$$\Phi(\delta_L)(\pi_{\Phi L}(p)) = (\delta_L)_*(\Phi L \setminus \overline{\{p\}}) = (\delta_L)_*(\Phi_p) = p$$

by 3.3.1 and 3.4.1. \square

3.5.2. In particular, since π is a natural equivalence,

Φ and \mathfrak{D} induce a dual equivalence between \mathbf{Top}_D and the subcategory of \mathbf{Frm}_D given by the T_D -spatial frames (recall 1.7).

3.5.3. The following fact is related to a result from Bruns, [5].

Proposition. *A frame L is T_D -spatial if and only if any proper interval $\{x \mid a \leq x \leq b\}$ in L contains a covered pair.*

Proof. Let $L = \mathfrak{D}X$ for some T_D -space X and $U \subsetneq Y$ in $\mathfrak{D}X$. Then, for $x \in V \setminus U$, $(V \setminus \overline{\{x\}}) \cup \{x\}$ is open since $(X \setminus \overline{\{x\}}) \cup \{x\}$ is open by 2.2.1; further, $U \subseteq X \setminus \overline{\{x\}}$ as $x \notin U$ and hence

$$U \subseteq V \cap (X \setminus \overline{\{x\}}) = (V \setminus \overline{\{x\}}) \ll (V \setminus \overline{\{x\}}) \cup \{x\} \subseteq V,$$

exhibiting a covered pair between U and V .

Conversely, given the condition in question, we have to show that δ_L is an isomorphism, and for this it is enough to prove $\Phi_a \not\leq \Phi_b$ whenever $a < b$. Now, by our hypothesis, we have $u, v \in L$ such that $a \leq u < v \leq b$, hence the homomorphism

$$h : L \rightarrow \{u, v\} \cong \mathbf{2}, \quad s \mapsto (s \vee u) \wedge v,$$

and therefore the completely prime filter

$$F = \{s \in L \mid h(s) = v\} = \{s \in L \mid s \vee u \geq v\} = u \uparrow v$$

(recall 2.5). Finally, F is a slicing filter by 2.6.2, and since $a \notin F$ and $b \in F$ this shows for the (completely!) prime p associated with F that $p \notin \Phi_a$ and $p \in \Phi_b$. \square

3.6. Consider the inverse $j_X : \Phi\mathfrak{D}X \rightarrow X$ to the adjunction homeomorphism π_X , given by the formula

$$j_X(X \setminus \overline{\{x\}}) = x.$$

Now let us extend this definition to a general space X (recall that our spaces are T_0 and hence j_X is well-defined).

For an open set U in X we have

$$j_X^{-1}[U] = \{X \setminus \overline{\{x\}} \mid x \in U\} = \{X \setminus \overline{\{x\}} \mid U \not\subseteq X \setminus \overline{\{x\}}\} = \Phi_U$$

so that the $j_X^{-1}[U]$ are precisely the open subsets of $\Phi\mathfrak{D}X$. Thus

j_X is a homeomorphic embedding of (the T_D -space) $\Phi\mathfrak{D}X$ onto the subspace of X constituted by all the T_D -points.

This suggests something like a T_D -coreflection. But \mathbf{Top}_D is not coreflective in \mathbf{Top} : it is not closed under quotients, not even if we restrict ourselves to the T_0 -case, as is seen in the following

3.6.1. Example. Let \mathbb{N} be the set of natural numbers, $\omega \notin \mathbb{N}$. Consider

$$X = (\mathbb{N} \times \{0, 1\}) \cup \{\omega\}$$

endowed by the following topology:

$$U \subseteq X \text{ is open if } \begin{cases} (\exists n, (n, 1) \in U) \Rightarrow \omega \in U, \text{ and} \\ \omega \in U \Rightarrow \exists k (n \geq k \Rightarrow (n, 0) \in U). \end{cases}$$

Thus in particular $\mathbb{N} \times \{0\}$, $(\mathbb{N} \times \{0\}) \cup \{\omega\}$ and each $(\mathbb{N} \times \{0\}) \cup \{\omega\} \cup \{(n, 1)\}$ are open, and we see that X is T_D .

Now define $q : X \rightarrow Y = \mathbb{N} \times \{\omega\}$ by setting $q(n, i) = n$ and $q(\omega) = \omega$. In the corresponding quotient topology on Y , U is open iff either it is \emptyset or contains ω and all $n \geq k$ for some k . Thus, Y is still T_0 , but it is not T_D : ω is not a T_D -point.

3.7. Let \mathcal{A} be a general category and let \mathcal{C} be a full subcategory of \mathcal{A} . Let us have to each object $A \in \mathcal{A}$ assigned an $A^\circ \in \mathcal{C}$ and a monomorphism

$$\iota_A : A^\circ \rightarrow A$$

such that

$$\text{for } C \in \mathcal{C}, C^\circ = C \text{ and } \iota_C = \text{id}_C, \text{ in particular } A^{\circ\circ} = A^\circ.$$

Define a subcategory \mathcal{A}° of \mathcal{A} as follows.

- the class of objects of \mathcal{A}° is the same as that of \mathcal{A} , and
- $f : A \rightarrow B$ from \mathcal{A} is a morphism in \mathcal{A}° iff there exists an $f^\circ : A^\circ \rightarrow B^\circ$ such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \iota_A \uparrow & & \uparrow \iota_B \\ A^\circ & \xrightarrow{f^\circ} & B^\circ \end{array}$$

commutes.

We have the immediate

Observation. 1. *The correspondence $f \mapsto f^\circ$ constitutes a functor $\mathcal{A}^\circ \rightarrow \mathcal{A}$, and $\iota = (\iota_A)_A$ is a natural transformation.*

2. \mathcal{C} is a full subcategory of \mathcal{A}° .

3. \mathcal{C} is a mono-coreflective subcategory of \mathcal{A}° with coreflection maps $\iota_A : \mathcal{A}^\circ \rightarrow \mathcal{A}$

3.7.1. This applies to the situation in 3.6. Let us specify more concretely the subcategory of **Top** appearing as the \mathcal{A}° above.

Lemma. *There exists a continuous $f^\circ : \Phi\mathfrak{D}X \rightarrow \Phi\mathfrak{D}Y$ such that*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ j_X \uparrow & & \uparrow j_Y \\ \Phi\mathfrak{D}X & \xrightarrow{f^\circ} & \Phi\mathfrak{D}Y \end{array}$$

commutes iff $\mathfrak{D}f$ is a D -homomorphism.

Proof. $(\mathfrak{D}f)_*(X \setminus \overline{\{x\}}) = Y \setminus \overline{\{f(x)\}}$. Thus, by 2.3.2, the completely prime elements are sent by $(\mathfrak{D}f)_*$ to completely prime ones iff the image of a T_D -point is a T_D -point, that is, iff f can be restricted to the subspaces of T_D points. \square

3.7.2. Conclusion. Define

${}_D\mathbf{Top}$

as the category of all (T_0) spaces and the continuous maps f such that $\mathfrak{D}f$ is a D -homomorphism. Then

\mathbf{Top}_D is a (mono-)coreflective subcategory of ${}_D\mathbf{Top}$.

4 T_D versus sobriety

4.1. Lemma. *Let j be the identical embedding $X \setminus \{x\} \rightarrow X$. Then $\mathfrak{D}j$ is an isomorphism iff x is not a T_D -point.*

Proof. We have $\mathfrak{D}j(U) = U \setminus \{x\}$. $\mathfrak{D}j$ is trivially onto, and it is one-one if and only if $U \setminus \{x\}$ is never open unless $x \notin U$. \square

4.2. Observation. *For $Z \subseteq Y \subseteq X$ let $k : Z \rightarrow Y$ and $j : Y \rightarrow X$ be the identical embedding and put $l = j \cdot k$. If $\mathfrak{D}l$ is an isomorphism then both $\mathfrak{D}j$ and $\mathfrak{D}k$ are isomorphisms.*

($\mathfrak{D}j \cdot \mathfrak{D}k$ is one-one, hence $\mathfrak{D}k$ is, which makes it an isomorphism; then $\mathfrak{D}j = \mathfrak{D}l(\mathfrak{D}k)^{-1}$.)

4.3. For any spaces X and Y consider the relation

$R(X, Y) : X$ is a proper subspace of Y such that the identical embedding $X \rightarrow Y$ induces an isomorphism $\mathfrak{D}Y \rightarrow \mathfrak{D}X$.

Then, sobriety and T_D have the following parallel characterizations.

Proposition. X is $\begin{cases} \text{sober} \\ T_D \end{cases}$ iff $\neg(\exists Y) \begin{cases} R(X, Y) \\ R(Y, X) \end{cases}$.

Proof. (1) If X is not sober then $R(X, Y)$ for the sobrification $Y \supsetneq X$. Conversely, given Y such that $R(X, Y)$, take any $y \in Y \setminus X$ and let

$$\mathcal{F} = \{U \cap X \mid y \in U \in \mathfrak{D}Y\}.$$

Then \mathcal{F} is a completely prime filter in $\mathfrak{D}X$, given $R(X, Y)$, which is not fixed in X (since all our spaces are T_0), showing X is not sober.

(2) If X is not T_D then there exists a non- T_D -point $x \in X$, and by 4.1 we have $R(X, Y)$ for $Y = X \setminus \{x\}$. Conversely, given Y such that $R(X, Y)$, take any $x \in X \setminus Y$ and consider the identical embeddings

$$Y \rightarrow X \setminus \{x\} \rightarrow X.$$

Then $R(Y, X)$ implies $R(X \setminus \{x\}, X)$ by 4.2, and by 4.1 x is not a T_D -point, showing X is not T_D . \square

4.4.1. Corollary. If $X \subsetneq Y$ and $\mathfrak{D}j_{XY}$ is an isomorphism then X is not sober and Y is not T_D .

In particular we obtain the well known fact that

a non-trivial sobrification is never T_D .

4.5. In [4], Berger proved (a.o.) that hereditarily sober spaces are T_D . In fact we have more.

Corollary. If for each $x \in X$ the space $X \setminus \{x\}$ is sober then X is T_D .

(Indeed, for any a non- T_D -point $x \in X$ we have $R(X \setminus \{x\}, X)$, contradicting the sobriety of $X \setminus \{x\}$ by 4.3.)

4.6. The non-fixed completely prime filters are (up to isomorphism) precisely the individual points y that one can *add* and keep the $\mathfrak{D}(X \cup \{y\})$ isomorphic with $\mathfrak{D}X$.

The non- T_D -points are precisely the individual points x that one can *subtract* and keep the $\mathfrak{D}(X \setminus \{x\})$ isomorphic with $\mathfrak{D}X$.

The symmetry between sobriety and the T_D property (which has been expressed already in 4.3) cannot be stretched much further, though: while we can add all the completely non-fixed prime filters to X to obtain the *maximal* Y with $R(X, Y)$, namely the sobrification, we do not generally have a *minimal* $X \subseteq Y$ with $R(X, Y)$. We would have to remove, first, all the non- T_D -points, which could result in the void subspace.

4.6.3. Example. Let X be any order-dense subset of the set \mathbb{R} of reals, endowed with the topology consisting of \emptyset and all

$$> x = \{u \mid x < y\}.$$

In X , obviously, none of the points is T_D (removing any point from $> x$ destroys the openness).

Since T_D -spatiality implies the existence of covered elements (recall 3.5.3) our $\mathfrak{D}X$ is also an example of a spatial frame that is not T_D -spatial.

4.7. In point-free topology one uses the concept of *subfitness* (if $a < b$ then $a \vee c < 1 = b \vee c$ for some c) as a weaker counterpart of T_1 . The question naturally arises whether there is some relation between the subfitness of $\mathfrak{D}X$ and the T_D -property of X . There is not. We have

Proposition. *For T_0 -spaces X , $\mathfrak{D}X$ subfit neither implies nor is implied by X being T_D .*

Proof. (\Rightarrow) Let Y be any T_1 -space and $X \supseteq Y$ its sobrification. Then $\mathfrak{D}X \cong \mathfrak{D}Y$, and since $\mathfrak{D}Y$ is trivially subfit the same holds for $\mathfrak{D}X$. Suppose now that X is T_D . Then every neighbourhood filter $\mathcal{U}(x)$, $x \in X$, is a slicing filter of $\mathfrak{D}X$, and by the relation between X and Y it follows that every completely prime filter of $\mathfrak{D}Y$ is slicing. That, however, implies that every completely prime filter of $\mathfrak{D}Y$ is fixed (2.6.1), saying Y is sober – which is known not to hold for every Y .

(\Leftarrow) Let X be the unit interval $[0, 1]$ with the usual downsets as open sets. Then X is T_D but it is not subfit since for any $0 \leq \delta < \mu < 1$ and for any downset $U \subseteq [0, 1]$ such that $(\downarrow \mu) \cup U = [0, 1]$ we also have $(\downarrow \delta) \cup U = [0, 1]$.

□

References

- [1] C.E. Aull and W.J. Thron, *Separation axioms between T_0 and T_1* , Indag. Math. **24** (1962), 26-37.
- [2] R.N. Ball, *Distributive Cauchy lattices*, Algebra Universalis **18** (1984), 134-174.
- [3] R.N. Ball, *Convergence and Cauchy structures in lattice ordered groups*, Trans.Amer.Math.Soc. **259** (1980), 357-392.
- [4] S.F. Berger, *Hereditary sobriety and the cardinality of T_1 topologies on countable sets*, Quaestiones Mathematicae **20** (1997), 109-126.
- [5] G. Bruns, *Darstellungen und Erweiterungen geordneter Mengen II*, Journal für Mathematik **210** (1962), 1-23.
- [6] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique, tome I: le langage des schémas*, I.H.E.S. Publ.Math., no.4.
- [7] P.T. Johnstone, *Stone spaces*, Cambridge studies in advanced math. No 3, Cambridge University Press, Cambridge, 1983.
- [8] S. Mac Lane, *Categories for the Working Mathematician*, Springer-Verlag, New York 1971.
- [9] A. Pultr, *Frames*, Chapter in: Handbook of Algebra, Vol.3, (Ed. by M. Hazewinkel), Elsevier 2003, 791-858.
- [10] A. Pultr and A. Tozzi, *A note on reconstruction of spaces and maps from lattice data*, Quaestiones Mathematicae **24** (2001), 55-63.
- [11] W.J. Thron, *Lattice-equivalence of topological spaces*, Duke Math. J. **29** (1962), 671-679.