

# Non-three-colorable common graphs exist \*

Hamed Hatami <sup>†</sup>    Jan Hladký <sup>‡</sup>    Daniel Král' <sup>§</sup>

Serguei Norine <sup>¶</sup>    Alexander Razborov <sup>||</sup>

May 22, 2011

## Abstract

A graph  $H$  is called *common* if the total number of copies of  $H$  in every graph and its complement asymptotically minimizes for random graphs. A former conjecture of Burr and Rosta, extending a conjecture of Erdős asserted that every graph is common. Thomason disproved both conjectures by showing that  $K_4$  is not common. It is now known that in fact the common graphs are very rare. Answering a question of Sidorenko and of Jagger, Štoviček and Thomason from 1996 we show that the 5-wheel is common. This provides the first example of a common graph that is not three-colorable.

---

\*HH was partially supported by an NSERC Discovery Grant. JH was supported by EPSRC award EP/D063191/1. DK: The work leading to this invention has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 259385. SN was supported in part by NSF under Grant No. DMS-0803214. Part of AR's work was done while the author was at Steklov Mathematical Institute, supported by the Russian Foundation for Basic Research, and at Toyota Technological Institute, Chicago.

<sup>†</sup>School of Computer Science, McGill University, Montreal, Canada., email: [hatami@cs.mcgill.ca](mailto:hatami@cs.mcgill.ca)

<sup>‡</sup>Department of Applied Mathematics, Faculty of Mathematics and Physics, Charles University, Malostranské náměstí 25, 118 00 Prague, Czech Republic, and DIMAP, Department of Computer Science, University of Warwick, Coventry CV4 7AL, United Kingdom, email: [honzahladky@gmail.com](mailto:honzahladky@gmail.com)

<sup>§</sup>Institute for Theoretical Computer Science, Faculty of Mathematics and Physics, Charles University, Malostranské náměstí 25, 118 00 Prague, Czech Republic., email: [kral@kam.mff.cuni.cz](mailto:kral@kam.mff.cuni.cz)

<sup>¶</sup>Department of Mathematics, Princeton University, Princeton, NJ, USA., email: [snorin@math.princeton.edu](mailto:snorin@math.princeton.edu)

<sup>||</sup>Department of Computer Science, University of Chicago, IL, USA., email: [razborov@cs.uchicago.edu](mailto:razborov@cs.uchicago.edu)

# 1 Introduction

A natural question in extremal graph theory is how many monochromatic subgraphs isomorphic to a graph  $H$  must be contained in any two-coloring of the edges of the complete graph  $K_n$ . Equivalently, how many subgraphs isomorphic to a graph  $H$  must be contained in a graph and its complement?

Goodman [Goo59] showed that for  $H = K_3$ , the optimum solution is essentially obtained by a typical random graph. The graphs  $H$  that satisfy this property are called *common*. Erdős [Erd62] conjectured that all complete graphs are common. Later, this conjecture was extended to all graphs by Burr and Rosta [BR80]. Sidorenko [Sid89] disproved Burr and Rosta's conjecture by showing that a triangle with a pendant edge is not common. Later Thomason [Tho89] disproved Erdős's conjecture by showing that for  $p \geq 4$ , the complete graphs  $K_p$  are not common. It is now known that in fact the common graphs are very rare. For example, Jagger, Šťovíček and Thomason [JŠT96] showed that every graph that contains  $K_4$  as a subgraph is not common. If we work with  $k$ -edge-colorings of  $K_n$  rather than 2-edge-colorings we get the notion of a  $k$ -*common* graph. Cummings and Young [CY] recently proved that no graph containing the triangle  $K_3$  is 3-common, a counterpart of the result of Jagger, Šťovíček and Thomason above.

There are some classes of graphs that are known to be common. Sidorenko [Sid89] showed that cycles are common. A conjecture due to Erdős and Simonovits [ES84] and Sidorenko [Sid91, Sid93] asserts that for every bipartite graph  $H$ , among graphs of given density random graphs essentially contain the least number of subgraphs isomorphic to  $H$ . It is not hard to see that every graph  $H$  with the latter property is common, therefore this conjecture would imply that all bipartite graphs are common. The Erdős-Simonovits-Sidorenko conjecture has been verified for a handful of graphs [Sid93, Sid96, Hat10, CFS10], and hence there are various classes of bipartite graphs that are known to be common. In [JŠT96] and [Sid96] some graph operations are introduced that can be used to "glue" common graphs in order to construct new common graphs. However none of these operations can increase the chromatic number to a number larger than three, and as a result, all of the known common graphs are of chromatic number at most 3. With these considerations Jagger, Šťovíček and Thomason [JŠT96] state "We regard the determination of the commonality of  $W_5$  [the wheel with 5 spokes] as the most interesting open problem in the area."

We will prove in Theorem 3.1 that  $W_5$  (see Figure 1) is common. This




Figure 1: The 5-wheel.

will also answer a question of Sidorenko [Sid96]. He showed [Sid96, Theorem 8] that every graph that is obtained by adding a vertex of full degree to a bipartite graph of average degree at least one satisfying the Erdős-Simonovits-Sidorenko conjecture is common. Sidorenko further asked whether in this theorem both conditions of being bipartite and having average degree at least one are essential in order to obtain a common graph. Our result answers his question in the negative, as  $W_5$  is obtained by adding a vertex of full degree to a non-bipartite graph.

The proof of Theorem 3.1 is a rather standard Cauchy-Schwarz calculation in flag algebras [Raz07], and is generated with the aid of a computer using semi-definite programming. A similar approach was successfully applied for example in [Raz10, HKN09, BT11, Grz11, HHK<sup>+</sup>11].

## 2 Preliminaries

We write vectors with bold font, e.g.  $\mathbf{a} = (\mathbf{a}(1), \mathbf{a}(2), \mathbf{a}(3))$  is a vector with three coordinates. For every positive integer  $k$ ,  $[k]$  denotes the set  $\{1, \dots, k\}$ .

All graphs in this paper are finite and simple (that is, loops and multiple edges are not allowed). For every natural number  $n$ , let  $\mathcal{M}_n$  denote the set of all simple graphs on  $n$  vertices up to an isomorphism. For a graph  $G$ , let  $V(G)$  and  $E(G)$ , respectively denote the set of the vertices and the edges of  $G$ . The complement of  $G$  is denoted by  $G^*$ .

The *homomorphism density* of a graph  $H$  in a graph  $G$ , denoted by  $t(H; G)$ , is the probability that a random map from the vertices of  $H$  to the vertices of  $G$  is a *graph homomorphism*, that is it maps every edge of  $H$  to an edge of  $G$ . If  $H \in \mathcal{M}_\ell$ ,  $G \in \mathcal{M}_n$ , and  $\ell \leq n$ , then  $t_0(H; G)$  denotes the probability that a random **injective** map from  $V(H)$  to  $V(G)$  is a graph homomorphism, and  $p(H, G)$  denotes the probability that a random set of  $\ell$  vertices of  $G$  induces a graph isomorphic to  $H$ .

We have the following chain rule (cf. [Raz07, Lemma 2.2]):

$$t_0(H; G) = \sum_{F \in \mathcal{M}_\ell} t_0(H; F) p(F, G), \quad (2.1)$$

where  $|V(H)| \leq \ell \leq |V(G)|$ .

**Definition 2.1.** *A graph  $H$  is called common if*

$$\liminf_{n \rightarrow \infty} \min_{G \in \mathcal{M}_n} (t(H; G) + t(H; G^*)) \geq 2^{1-|E(H)|}. \quad (2.2)$$

It is easy to see that as  $n \rightarrow \infty$ , for a random graph  $G$  on  $n$  vertices, we have, with high probability,  $t(H; G) + t(H; G^*) = 2^{1-|E(H)|} \pm o(1)$ . Thus,  $H$  is common if the total number of copies of  $H$  in every graph and its complement asymptotically minimizes for random graphs. Note also that since  $t(H; G)$  and  $t_0(H; G)$  are asymptotically equal (again, as  $n \rightarrow \infty$ ), one could use  $t_0(H; G)$  in place of  $t(H; G)$  in (2.2), and this is what we will do in our proof.

## 2.1 Flag algebras

We assume certain familiarity with the theory of flag algebras from [Raz07]. However, for the proof of the central Theorem 3.1 only the most basic notions are required. Thus, instead of trying to duplicate definitions, we occasionally give pointers to relevant places in [Raz07].

In our application of the flag algebras calculus we work exclusively with the theory of simple graphs (cf. [Raz07, §2]). As in [Raz07], flags of type  $\sigma$  and size  $k$  are denoted by  $\mathcal{F}_k^\sigma$ . The flag algebra generated by all flags of type  $\sigma$  is denoted by  $\mathcal{A}^\sigma$  (cf. [Raz07, §2]). Apart from already defined model  $W_5 \in \mathcal{M}_6$  we need to introduce the following models, types, and flags.

We shall work with five types  $\sigma_0, \sigma_1, \dots, \sigma_4$  of size four which are illustrated in Figure 2. For a type  $\sigma$  of size  $k$  and a set of vertices  $V \subseteq [k]$  in  $\sigma$ , let  $F_V^\sigma$  denote the flag  $(G, \theta) \in \mathcal{F}_{k+1}^\sigma$  in which the only unlabeled vertex  $v$  is connected to the set  $\{\theta(i) : i \in V\}$ . We further define  $f_V^\sigma \in \mathcal{A}^\sigma$  by

$$f_V^\sigma \stackrel{\text{def}}{=} F_\emptyset^\sigma - \frac{1}{|\text{Aut}(\sigma)|} \cdot \sum_{\eta \in \text{Aut}(\sigma)} F_{\eta(V)}^\sigma.$$

These elements form a basis (for  $V \neq \emptyset$  and with repetitions) in the space spanned by those  $f \in \mathcal{A}_{k+1}^\sigma$  that are both  $\text{Aut}(\sigma)$ -invariant and asymptotically vanish on random graphs; other than that, our particular choice of elements with this property is more or less arbitrary.




Figure 2: Types.

Recall that in [Raz07, §2.2] a certain “averaging operator”  $[\cdot]$  was introduced. This operator plays a central role in the flag algebra calculus.

Let  $*$   $\in$   $\text{Aut}(\mathcal{A}^0)$  be the involution that corresponds to taking the complementary graph. That is, we extend  $*$  linearly from  $\bigcup_n \mathcal{M}_n$  to  $\mathcal{A}^0$ .

### 3 Main result

We can now state the main result of the paper.

**Theorem 3.1.** *The 5-wheel  $W_5$  is common.*

*Proof.* Let  $\widehat{W}_5 \in \mathcal{A}^0$  be the element that counts the injective homomorphism density of the 5-wheel, that is

$$\widehat{W}_5 \stackrel{\text{def}}{=} \sum_{F \in \mathcal{M}_6} t_0(W_5, F) F.$$

We shall prove that

$$\widehat{W}_5 + \widehat{W}_5^* \geq 2^{-9}, \tag{3.1}$$

where the inequality  $\leq$  in the algebra  $\mathcal{A}^0$  is defined in [Raz07, Definition 6]. An alternate interpretation of this inequality [Raz07, Corollary 3.4] is that

$$\liminf_{n \rightarrow \infty} \min_{G \in \mathcal{M}_n} (p(\widehat{W}_5, G) + p(\widehat{W}_5^*, G)) \geq 2^{-9}.$$

Since  $p(\widehat{W}_5, G) = \sum_{F \in \mathcal{M}_6} t_0(\widehat{W}_5; F)p(F; G) = t_0(\widehat{W}_5; G)$  by (2.1), and, likewise,  $p(\widehat{W}_5^*, G) = p(\widehat{W}_5, G^*) = t_0(\widehat{W}_5; G^*)$ , (3.1) implies Theorem 3.1.

We now give a proof of (3.1). To this end we work with suitable quadratic forms  $Q_{\sigma_i}^{+/-}$  defined by symmetric matrices  $M_{\sigma_i}^{+/-}$  and vectors  $\mathbf{g}_i^{+/-}$  in the algebras  $\mathcal{A}^{\sigma_i}$ . The numerical values of the matrices  $M_{\sigma_i}^{+/-}$  and vectors  $\mathbf{g}_i^{+/-}$  are given in the appendix. It is essential that all the matrices  $M_{\sigma_i}^{+/-}$  are positive definite which can be verified using any general mathematical software. Next we define

$$R := \left( \sum_{i=0}^4 \llbracket Q_{\sigma_i}^+(\mathbf{g}_i^+) \rrbracket_{\sigma_i} \right) + \llbracket Q_{\sigma_1}^-(\mathbf{g}_1^-) \rrbracket_{\sigma_1} + \llbracket Q_{\sigma_4}^-(\mathbf{g}_4^-) \rrbracket_{\sigma_4}.$$

We claim that

$$\widehat{W}_5 + \widehat{W}_5^* = 2^{-9} + R + R^*. \quad (3.2)$$

All the terms in (3.2) can be expressed as linear combinations of graphs from  $\mathcal{M}_6$  and thus checking (3.2) amounts to checking the coefficients of the 156 flags from  $\mathcal{M}_6$ . We offer a C-code available at <http://kam.mff.cuni.cz/~kral/wheel> that verifies the equality (3.2).

By [Raz07, Theorem 3.14], we have

$$\left( \sum_{i=0}^4 \llbracket Q_{\sigma_i}^+(\mathbf{g}_i^+) \rrbracket_{\sigma_i} \right) + \llbracket Q_{\sigma_1}^-(\mathbf{g}_1^-) \rrbracket_{\sigma_1} + \llbracket Q_{\sigma_4}^-(\mathbf{g}_4^-) \rrbracket_{\sigma_4} \geq 0.$$

Therefore, (3.2) implies (3.1).  $\square$

Theorem 3.1 shows that a typical random graph  $G = G_{n, \frac{1}{2}}$  asymptotically minimizes the quantity  $t(W_5; G) + t(W_5; G^*)$ . Extending our method, we convinced ourselves that  $G_{n, \frac{1}{2}}$  is essentially the only minimizer of  $t(W_5; G) + t(W_5; G^*)$ . In terms of flag algebras this means that the homomorphism  $\phi \in \text{Hom}^+(\mathcal{A}^0, \mathbb{R})$  (see [Raz07, Definition 5]) satisfying  $\phi(\widehat{W}_5 + \widehat{W}_5^*) = 2^{-9}$  is unique.

The outline of the argument is as follows. Let  $\rho \in \mathcal{M}_2$  denote a graph consisting of a single edge, let  $C_4 \in \mathcal{M}_4$  denote the cycle of length 4, and, as before, let

$$\widehat{C}_4 \stackrel{\text{def}}{=} \sum_{F \in \mathcal{M}_4} t_0(C_4; F)F.$$

The Erdős-Simonovits-Sidorenko conjecture is known for  $C_4$  [Sid91], and it implies that  $\widehat{C}_4 \geq \rho^4$  and  $\widehat{C}_4^* \geq (1 - \rho)^4$  in  $\mathcal{A}^0$ . Therefore,  $C_4 + C_4^* \geq 1/8$  (i.e.,  $C_4$  is common), and, moreover, every  $\phi \in \text{Hom}^+(\mathcal{A}^0, \mathbb{R})$  attaining equality must satisfy  $\phi(\rho) = 1/2$  and  $\phi(\widehat{C}_4) = 1/16$ .

On the other hand, it is shown in [CGW89] that the density of edges and the density cycles of length 4 characterize quasi-random graphs, implying that the homomorphism  $\phi$  satisfying  $\phi(\widehat{C}_4 + \widehat{C}_4^*) = 1/8$  is unique (and corresponds to quasi-random graphs). Therefore, to verify the uniqueness of the homomorphism  $\phi$  satisfying  $\phi(\widehat{W}_5 + \widehat{W}_5^*) = 2^{-9}$  it suffices to show that

$$\widehat{W}_5 + \widehat{W}_5^* \geq 2^{-9} + \frac{1}{100} (\widehat{C}_4 + \widehat{C}_4^* - 1/8). \quad (3.3)$$

We have used a computer program to verify (3.3), and it is telling us that this inequality holds with quite a convincing level of accuracy  $10^{-10}$ . But we have not converted the floating point computations into a rigorous proof.

## 4 Conclusion

In this paper we have exhibited the first example of a common graph that is not three-colorable. This naturally gives rise to the following interesting question: do there exist common graphs with arbitrarily large chromatic number?

## References

- [BR80] Stefan A. Burr and Vera Rosta. On the Ramsey multiplicities of graphs—problems and recent results. *J. Graph Theory*, 4(4):347–361, 1980.
- [BT11] Rahil Baber and John Talbot. Hypergraphs do jump. *Combin. Probab. Comput.*, 20(2):161–171, 2011.
- [CFS10] David Conlon, Jacob Fox, and Benny Sudakov. An approximate version of Sidorenko’s conjecture. *Geom. Funct. Anal.*, 20(6):1354–1366, 2010.
- [CGW89] Fan-Rong K. Chung, Ronald L. Graham, and Richard M. Wilson. Quasi-random graphs. *Combinatorica*, 9(4):345–362, 1989.

- [CY] James Cummings and Michael Young. Graphs containing triangles are not 3-common. *J. Combin.*, to appear.
- [Erd62] Paul Erdős. On the number of complete subgraphs contained in certain graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 7:459–464, 1962.
- [ES84] Paul Erdős and Miklós Simonovits. Cube-supersaturated graphs and related problems. In *Progress in graph theory (Waterloo, Ont., 1982)*, pages 203–218. Academic Press, Toronto, ON, 1984.
- [Goo59] Al W. Goodman. On sets of acquaintances and strangers at any party. *Amer. Math. Monthly*, 66:778–783, 1959.
- [Grz11] Andrzej Grzesik. On the maximum number of  $C_5$ 's in a triangle-free graph. arXiv:1102.0962, 2011.
- [Hat10] Hamed Hatami. Graph norms and Sidorenko's conjecture. *Israel J. Math.*, 175(1):125–150, 2010.
- [HHK<sup>+</sup>11] Hamed Hatami, Jan Hladký, Daniel Král', Serguei Norine, and Alexander Razborov. On the number of pentagons in triangle-free graphs. arXiv:1102.1634, 2011.
- [HKN09] Jan Hladký, Daniel Král', and Serguei Norine. Counting flags in triangle-free digraphs. arXiv:0908.2791, 2009.
- [JŠT96] Chris Jagger, Pavel Šťovíček, and Andrew Thomason. Multiplicities of subgraphs. *Combinatorica*, 16(1):123–141, 1996.
- [Raz07] Alexander A. Razborov. Flag algebras. *J. Symbolic Logic*, 72(4):1239–1282, 2007.
- [Raz10] Alexander A. Razborov. On 3-hypergraphs with forbidden 4-vertex configurations. *SIAM J. Discrete Math.*, 24(3):946–963, 2010.
- [Sid89] Alexander Sidorenko. Cycles in graphs and functional inequalities. *Mat. Zametki*, 46(5):72–79, 104, 1989.
- [Sid91] Alexander Sidorenko. Inequalities for functionals generated by bipartite graphs. *Diskret. Mat.*, 3(3):50–65, 1991.

- [Sid93] Alexander Sidorenko. A correlation inequality for bipartite graphs. *Graphs Combin.*, 9(2):201–204, 1993.
- [Sid96] Alexander Sidorenko. Randomness friendly graphs. *Random Structures Algorithms*, 8(3):229–241, 1996.
- [Tho89] Andrew Thomason. A disproof of a conjecture of Erdős in Ramsey theory. *J. London Math. Soc. (2)*, 39(2):246–255, 1989.

## A The matrices $M_i^{+/-}$ and the vectors $\mathbf{g}_i^{+/-}$

Here, we list the numerical values of the matrices  $M_i^{+/-}$  and the vectors  $\mathbf{g}_i^{+/-}$ .

The vectors  $\mathbf{g}_i^+$  are given by the tuples

$$\begin{aligned} \mathbf{g}_0^+ &\stackrel{\text{def}}{=} (f_{\{1\}}^{\sigma_0}, f_{\{1,2\}}^{\sigma_0}, f_{\{1,2,3\}}^{\sigma_0}, f_{\{1,2,3,4\}}^{\sigma_0}) \\ \mathbf{g}_1^+ &\stackrel{\text{def}}{=} (f_{\{1\}}^{\sigma_1}, f_{\{3\}}^{\sigma_1}, f_{\{1,2\}}^{\sigma_1}, f_{\{1,3\}}^{\sigma_1}, f_{\{3,4\}}^{\sigma_1}, f_{\{1,2,3\}}^{\sigma_1}, f_{\{1,3,4\}}^{\sigma_1}, f_{\{1,2,3,4\}}^{\sigma_1}) \\ \mathbf{g}_2^+ &\stackrel{\text{def}}{=} (f_{\{1\}}^{\sigma_2}, f_{\{2\}}^{\sigma_2}, f_{\{4\}}^{\sigma_2}, f_{\{1,2\}}^{\sigma_2}, f_{\{1,4\}}^{\sigma_2}, f_{\{2,3\}}^{\sigma_2}, f_{\{2,4\}}^{\sigma_2}, f_{\{1,2,3\}}^{\sigma_2}, \\ &\quad f_{\{1,2,4\}}^{\sigma_2}, f_{\{2,3,4\}}^{\sigma_2}, f_{\{1,2,3,4\}}^{\sigma_2}) \\ \mathbf{g}_3^+ &\stackrel{\text{def}}{=} (f_{\{1\}}^{\sigma_3}, f_{\{2\}}^{\sigma_3}, f_{\{1,2\}}^{\sigma_3}, f_{\{2,3\}}^{\sigma_3}, f_{\{1,2,3\}}^{\sigma_3}, f_{\{2,3,4\}}^{\sigma_3}, f_{\{1,2,3,4\}}^{\sigma_3}) \\ \mathbf{g}_4^+ &\stackrel{\text{def}}{=} (f_{\{1\}}^{\sigma_4}, f_{\{1,2\}}^{\sigma_4}, f_{\{1,3\}}^{\sigma_4}, f_{\{1,2,3\}}^{\sigma_4}, f_{\{1,2,3,4\}}^{\sigma_4}), \end{aligned}$$

and the vectors  $\mathbf{g}_i^-$  are given by

$$\begin{aligned} \mathbf{g}_1^- &\stackrel{\text{def}}{=} (F_{\{3\}}^{\sigma_1} - F_{\{4\}}^{\sigma_1}, F_{\{1,3,4\}}^{\sigma_1} - F_{\{2,3,4\}}^{\sigma_1}, F_{\{1,3\}}^{\sigma_1} - F_{\{2,3\}}^{\sigma_1}, F_{\{1,3\}}^{\sigma_1} - F_{\{2,4\}}^{\sigma_1}, \\ &\quad F_{\{1,3\}}^{\sigma_1} - F_{\{3,4\}}^{\sigma_1}) \\ \mathbf{g}_4^- &\stackrel{\text{def}}{=} (F_{\{1,2\}}^{\sigma_4} - F_{\{3,4\}}^{\sigma_4}, F_{\{1,3\}}^{\sigma_4} - F_{\{2,3\}}^{\sigma_4}, F_{\{1,3\}}^{\sigma_4} - F_{\{2,4\}}^{\sigma_4}, F_{\{1,3\}}^{\sigma_4} - F_{\{3,4\}}^{\sigma_4}). \end{aligned}$$

The matrices  $M_i^{+/-}$  are listed on the next three pages.

$$M_0^+ \stackrel{\text{def}}{=} \frac{1}{2 \cdot 10^8} \times \begin{pmatrix} 104133330 & -67645847 & -126443014 & -53041562 \\ -67645847 & 58559244 & 68999274 & 28961030 \\ -126443014 & 68999274 & 166581934 & 69653308 \\ -53041562 & 28961030 & 69653308 & 29368489 \end{pmatrix}$$

$$\begin{aligned}
M_3^+ &\stackrel{\text{def}}{=} \frac{1}{24 \cdot 10^8} \times \begin{pmatrix} 1770465360 & -40788068 & 770354664 & -280179622 & -1109635560 & -593033461 & -1434435065 \\ -40788068 & 503182008 & -377074674 & -65682192 & -316936632 & 337167432 & -405260664 \\ 770354664 & -377074674 & 942288720 & -5442408 & -584215338 & -635915808 & -299584920 \\ -280179622 & -65682192 & -5442408 & 90869472 & 187091280 & -48623352 & 356458176 \\ -1109635560 & -316936632 & -584215338 & 187091280 & 1325422128 & 196268064 & 1280101992 \\ -593033461 & 337167432 & -635915808 & -48623352 & 196268064 & 706802676 & -31363774 \\ -1434435065 & -405260664 & -299584920 & 356458176 & 1280101992 & -31363774 & 1763018404 \end{pmatrix} \\
M_4^+ &\stackrel{\text{def}}{=} \frac{1}{12 \cdot 10^8} \times \begin{pmatrix} 6589068 & -137160 & 60408 & -3635796 & -5354976 \\ -137160 & 3975070 & -399180 & -720636 & -1388043 \\ 60408 & -399180 & 3506988 & -1778640 & -3413616 \\ -3635796 & -720636 & -1778640 & 5107716 & 3969708 \\ -5354976 & -1388043 & -3413616 & 3969708 & 12276592 \end{pmatrix} \\
M_1^- &\stackrel{\text{def}}{=} \frac{1}{48 \cdot 10^8} \times \begin{pmatrix} 1871684759 & 828164352 & 153135600 & 2205677647 & 32494800 \\ 828164352 & 647325323 & 122226960 & 1702274830 & 23569680 \\ 153135600 & 122226960 & 32894794 & 317036160 & 988560 \\ 2205677647 & 1702274830 & 317036160 & 4533494520 & 62236800 \\ 32494800 & 23569680 & 988560 & 62236800 & 7445060 \end{pmatrix} \\
M_4^- &\stackrel{\text{def}}{=} \frac{1}{24 \cdot 10^8} \times \begin{pmatrix} 371929992 & -665160 & 31885344 & 6896381 \\ -665160 & 4952616 & 15347271 & -425892 \\ 31885344 & 15347271 & 420643536 & 5244336 \\ 6896381 & -425892 & 5244336 & 1704738 \end{pmatrix}
\end{aligned}$$

$$M_2^+ \stackrel{\text{def}}{=} \frac{1}{24 \cdot 10^8} \times$$

4114457904	-2123660510	578302533	2402100408	1609339896	-4979381511
-2123660510	4697332052	-146727648	-2893487330	-831349224	5132020824
578302533	-146727648	2842930424	-2377739616	2453284752	-1134538157
2402100408	-2893487330	-2377739616	5029589784	-1305679056	-3694198620
1609339896	-831349224	2453284752	-1305679056	2899169976	-3008866416
-4979381511	5132020824	-1134538157	-3694198620	-3008866416	9045922946
-1073916061	1140828192	949692648	-1628657160	227603736	1585531176
-711542544	-2533278088	-2122945241	2987352093	-2158976640	-492543642
-108075291	-3120849612	799767696	-17138568	1272333144	-2720802624
-311854200	586989168	-646840455	174993936	-824389152	1167719184
-1172726832	-2130186959	-1452441435	1346820763	-1468496784	119548200
-1073916061	-711542544	-108075291	-311854200	-1172726832	
1140828192	-2533278088	-3120849612	586989168	-2130186959	
949692648	-2122945241	799767696	-646840455	-1452441435	
-1628657160	2987352093	-17138568	174993936	1346820763	
227603736	-2158976640	1272333144	-824389152	-1468496784	
1585531176	-492543642	-2720802624	1167719184	119548200	
1198933584	-594013398	-787158072	-14360286	-864511462	
-594013398	4445640792	1152146526	408353664	3139157376	
-787158072	1152146526	4353119928	-778415544	2410765872	
-14360286	408353664	-778415544	430490652	217228440	
-864511462	3139157376	2410765872	217228440	3407087808	

$$M_1^+ \stackrel{\text{def}}{=} \frac{1}{24 \cdot 10^8} \times \begin{pmatrix} 3376427096 & -550659377 & 1175122309 & -274818336 & -1951510989 & 133242698 & -2978772360 & -1118255328 \\ -550659377 & 3579306230 & -2818779263 & 254758382 & 1853810147 & -3593215008 & 1149060744 & -2243131164 \\ 1175122309 & -2818779263 & 2446135762 & -153160723 & -1883990616 & 2571244464 & -1644918408 & 1392930672 \\ -274818336 & 254758382 & -153160723 & 259013952 & 207245488 & -524428416 & 59129384 & -87439632 \\ -1951510989 & 1853810147 & -1883990616 & 207245488 & 2026568566 & -1339529064 & 2075124696 & -196178016 \\ 133242698 & -3593215008 & 2571244464 & -524428416 & -1339529064 & 4383894552 & -474279456 & 2753404296 \\ -2978772360 & 1149060744 & -1644918408 & 59129384 & 2075124696 & -474279456 & 2987175794 & 578705400 \\ -1118255328 & -2243131164 & 1392930672 & -87439632 & -196178016 & 2753404296 & 578705400 & 2302497768 \end{pmatrix}$$