

# Exercises for Combinatorial and Computational Geometry II

## Series 2 — $k$ -holes, halving lines and graph drawings

deadline 7. 4. 2020

1. Let  $X$  be a finite non-empty set of points in the plane in general position. Prove that the following identities hold for the numbers of  $k$ -holes in  $X$ :

$$(a) \sum_{k=1}^{|X|} (-1)^k \cdot \#k\text{-holes} = -1, \quad [2]$$

- (b) If  $|X| \geq 2$ , then

$$\sum_{k=1}^{|X|} (-1)^k \cdot k \cdot \#k\text{-holes} = -\#\text{points inside } \text{conv}(X). \quad [2]$$

Hint: move points continuously along curves into a suitable configuration.

2. Let  $X$  be a set of  $n$  points in the plane in general position. Prove that the number of 4-holes is at least quadratic in  $n$ . [2]
3. Let  $P$  be a finite set of points in the plane that is not necessarily in general position and that contains no 5-hole. Prove that every convex 5-gon  $Q$  determined by points of  $P$  contains at least 1 point of  $P$  in the closed „inner“ 5-gon that is determined by the diagonals of  $Q$ . [2]
4. For  $n$  even, let  $P$  be a set of  $n$  points in the plane in general position. Furthermore, let  $k \leq n/2$  and let  $h$  be a line that does not intersect  $P$  and splits the plane into two halfplanes such that one of them contains exactly  $k$  points of  $P$ . Show that  $h$  intersects exactly  $k$  halving lines of  $P$ . [2]
5. Let  $P$  be a set of  $n$  points in the plane in general position. A pair of points in  $P$  is a  $k$ -edge if the line determined by these points separates exactly  $k$  points from  $P$  in one of the open hyperplanes. Let  $E_k(P)$  be the number of  $k$ -edges in  $P$  and furthermore let  $\overline{c\tau}(P)$  be the number of unordered 4-tuples of points in  $P$  in convex position.

Prove the following identity:

$$\overline{c\tau}(P) = 3 \binom{n}{4} - \sum_{k=0}^{\lfloor \frac{n-2}{2} \rfloor} E_k(P) \cdot k \cdot (n - k - 2).$$

Hint: count in two ways the number of ordered triples  $(a, \overline{bc}, d)$  such that  $a, b, c, d$  are pairwise different points of  $P$ ,  $a$  lies to the left of the line  $\overline{bc}$  and  $d$  lies to the right of the line of  $\overline{bc}$ . [2]

6. Two edges in a graph are *independent* if they do not share a vertex. In a *drawing* of a graph we assume that edges have only finitely many points in common, and that every common point of two edges is either their common endpoint or a crossing.

Determine for which values of  $m$  and  $n$ , in every drawing of the graph  $K_{m,n}$  the total number of crossings of independent pairs of edges is odd. Hint: what happens during a continuous deformation of the edges? [3]