

# Exercises for Combinatorial and Computational Geometry II

## Series 1 — Erdős–Szekeres theorem

hints 10. 3. 2020, deadline 17. 3. 2020

**Please choose a nickname that will be used in the list of scores on the webpage of the exercises. If you have already chosen a nickname, you can just sign your solutions with either your name or your nickname.**

1. (a) Show that for every  $k \in \mathbb{N}$  there is an  $n(k) \in \mathbb{N}$  such that every set of  $n(k)$  points in the plane contains  $k$  points in *general position* or  $k$  points on a common line. [2]  
(b) Prove the previous statement with  $n(k)$  at most polynomial in  $k$ . [1]
2. (a) Prove the Erdős–Szekeres theorem in  $\mathbb{R}^d$ : for every  $d \geq 3$  and  $k \in \mathbb{N}$  there is an  $n = n_d(k) \in \mathbb{N}$  such that every set of  $n$  points in  $\mathbb{R}^d$  in *general position* (no  $m + 2$  points in an affine subspace of dimension  $m$ , for  $m = 1, 2, \dots, d - 1$ ) contains a subset of  $k$  points in convex position. [2]  
(b) Show that every sufficiently large point set in  $\mathbb{R}^3$  in general position contains a 7-hole. [2]
3. Let  $P$  be a set of  $3n - 1$  points in the plane in convex position. Every closed segment between two points in  $P$  is colored either red or blue. Prove that there exist  $n$  pairwise disjoint red segments or  $n$  pairwise disjoint blue segments. [3]
4. Prove that there is a sufficiently large constant  $C$  such that the  $n \times n$  grid  $\{(i, j); i = 1, 2, \dots, n; j = 1, 2, \dots, n\}$  has no subset in convex position with more than  $Cn^{2/3}$  points. [4, hint]
5. Prove that for  $h \geq 1$ , the Horton set with  $2^h$  points has no subset in convex position with more than  $4h$  points. [2]

---

web: <http://kam.mff.cuni.cz/kvg/eng.html>