

Vector space

Definition: A *vector space* over a field $(\mathbb{K}, +, \cdot)$ is a set V together with a binary operation $+$ on V and an operation $\cdot : \mathbb{K} \times V \rightarrow V$, where:

- ▶ $(V, +)$ is an Abelian group
- ▶ $\forall a, b \in \mathbb{K}, \forall \mathbf{v} \in V : (a \cdot b) \cdot \mathbf{v} = a \cdot (b \cdot \mathbf{v})$
- ▶ $\forall \mathbf{v} \in V : 1 \cdot \mathbf{v} = \mathbf{v}$
- ▶ $\forall a, b \in \mathbb{K}, \forall \mathbf{v} \in V : (a + b) \cdot \mathbf{v} = (a \cdot \mathbf{v}) + (b \cdot \mathbf{v})$
- ▶ $\forall a \in \mathbb{K}, \forall \mathbf{u}, \mathbf{v} \in V : a \cdot (\mathbf{u} + \mathbf{v}) = (a \cdot \mathbf{u}) + (a \cdot \mathbf{v})$

Elements of \mathbb{K} are called *scalars*, elements of V are called *vectors*.

Vector space

Definition: A *vector space* over a field $(\mathbb{K}, +, \cdot)$ is a set V together with a binary operation $+$ on V and an operation $\cdot : \mathbb{K} \times V \rightarrow V$, where:

- ▶ $(V, +)$ is an Abelian group
- ▶ $\forall a, b \in \mathbb{K}, \forall v \in V : (a \cdot b) \cdot v = a \cdot (b \cdot v)$
- ▶ $\forall v \in V : 1 \cdot v = v$
- ▶ $\forall a, b \in \mathbb{K}, \forall v \in V : (a + b) \cdot v = (a \cdot v) + (b \cdot v)$
- ▶ $\forall a \in \mathbb{K}, \forall u, v \in V : a \cdot (u + v) = (a \cdot u) + (a \cdot v)$

Elements of \mathbb{K} are called *scalars*, elements of V are called *vectors*.

Distinguish the zero scalar $0 \in \mathbb{K}$ and the zero vector $\mathbf{0} \in V$;
also the opposite scalar $-a \in \mathbb{K}$ and the opposite vector $-v \in V$.

There is the inverse scalar $a^{-1} \in \mathbb{K}$ but **no** inverse vector!

There is **no** product of form $v \cdot a$!

The product symbols \cdot are often omitted and have priority to $+$.


Examples

- ▶ The *arithmetic* vector space \mathbb{K}^n of dimension n over \mathbb{K} .
Vectors are ordered n -tuples of elements of \mathbb{K} .
Additions and scalar multiplications are done by components.
Any field \mathbb{K} yields the vector space \mathbb{K}^1 of the same cardinality.
- ▶ $\mathbb{K}^{m \times n}$... matrices of order $m \times n$ over \mathbb{K} .
- ▶ $V = \{\mathbf{0}\}$ the *trivial* vector space over any field \mathbb{K} .
- ▶ Polynomials with coefficients in \mathbb{K} .
- ▶ Polynomials of bounded degree.


The vector space of real functions over the field \mathbb{R}

Vectors f, g are the real functions of real variable

Addition



A scalar multiple




Set systems as vector spaces

Let \mathcal{X} be a system of subsets of a set X that is closed under taking symmetric difference Δ .

Then $(\mathcal{X}, \Delta, \cdot)$ is a vector space over \mathbb{Z}_2 , where \cdot is defined as: $0 \cdot A = \emptyset$, the neutral element of Δ , and $1 \cdot A = A$ for all $A \in \mathcal{X}$.

Observe that $\forall A, B, C \in \mathcal{X} : (A \Delta B) \Delta C = A \Delta (B \Delta C)$,

since the result contains those elements of X that belong to the odd number of sets A, B and C .



The diagram consists of two rows of Venn diagrams. Each Venn diagram has three overlapping circles labeled A, B, and C. The top row shows the expression $(A \Delta B) \Delta C = A \Delta (B \Delta C)$. The first part, $(A \Delta B) \Delta C$, is represented by three circles where the regions with an odd number of colored circles (cyan, purple, yellow) are shaded. The second part, $A \Delta (B \Delta C)$, is represented by three circles where the regions with an odd number of colored circles (cyan, blue, yellow) are shaded. The final result is a Venn diagram where the regions with an odd number of colored circles (cyan, blue, yellow) are shaded. The bottom row shows the same expression with different colors: $(A \Delta B) \Delta C = A \Delta (B \Delta C)$ where A is cyan, B is red, and C is grey. The first part shows the regions with an odd number of colored circles (cyan, purple, yellow) shaded. The second part shows the regions with an odd number of colored circles (cyan, red, yellow) shaded. The final result is a Venn diagram where the regions with an odd number of colored circles (cyan, red, yellow) are shaded.


We may represent any $A \in \mathcal{X}$ by its characteristic function

$$\chi_A : X \rightarrow \mathbb{Z}_2 \text{ defined as } \chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$


The set $A \Delta B$ has the characteristic function $\chi_{A \Delta B} = \chi_A + \chi_B$, since $1 + 1 = 0$ in \mathbb{Z}_2 .

A vector space on graphs

Let $X = E_G$ and \mathcal{X} contain those edge sets A such that every vertex of G is incident with an even number of edges in A . Such sets A induce the so called *even subgraphs* of G .



Observe that Δ preserves the even degree, since the symmetric difference of two sets of *even* cardinality, namely the *edges* incident to a *vertex*, has also an *even* cardinality.



Properties

Observation: For any $\mathbf{v} \in V$ and $a \in \mathbb{K}$: $0\mathbf{v} = a\mathbf{0} = \mathbf{0}$

Proof:

$$0\mathbf{v} = 0\mathbf{v} + \mathbf{0} = 0\mathbf{v} + 0\mathbf{v} - 0\mathbf{v} = (0 + 0)\mathbf{v} - 0\mathbf{v} = 0\mathbf{v} - 0\mathbf{v} = \mathbf{0}$$

$$a\mathbf{0} = a\mathbf{0} + \mathbf{0} = a\mathbf{0} + a\mathbf{0} - a\mathbf{0} = a(\mathbf{0} + \mathbf{0}) - a\mathbf{0} = a\mathbf{0} - a\mathbf{0} = \mathbf{0}$$

Observation: For any $\mathbf{v} \in V$: $(-1)\mathbf{v} = -\mathbf{v}$

$$\text{Proof: } (-1)\mathbf{v} + \mathbf{v} = (-1)\mathbf{v} + 1\mathbf{v} = (-1 + 1)\mathbf{v} = 0\mathbf{v} = \mathbf{0}$$

Observation: If $a\mathbf{v} = \mathbf{0}$ then $a = 0$ or $\mathbf{v} = \mathbf{0}$.

$$\text{Proof: If } a \neq 0 \text{ then } \mathbf{v} = 1\mathbf{v} = a^{-1}a\mathbf{v} = a^{-1}\mathbf{0} = \mathbf{0}.$$

Subspace

Definition: Let V be a vector space over \mathbb{K} then a *subspace* U is a nonempty subset of V satisfying:

$$\blacktriangleright \forall \mathbf{u}, \mathbf{v} \in U : \mathbf{u} + \mathbf{v} \in U$$

$$\blacktriangleright \forall \mathbf{v} \in U, \forall a \in \mathbb{K} : a\mathbf{v} \in U$$

Observation: Any subspace is also a vector space because $\mathbf{0} = 0\mathbf{v} \in U$ and $-\mathbf{v} = (-1)\mathbf{v} \in U$ and other axioms hold already on V .


Example: For $V = \mathbb{R}^3$, and a line U in \mathbb{R}^3 through the origin, the line U is a subspace of V .

If we also consider a plane W in \mathbb{R}^3 containing the line U , then U is a subspace of W that is a subspace of V .

Intersection of subspaces

Theorem: Let $(U_i, i \in I)$ be any system of subspaces of a space V . The intersection of this system $\bigcap_{i \in I} U_i$ is also a subspace of V .

Example: Planes π and ρ are subspaces of the vector space \mathbb{R}^3 . Their intersection is the line p . It is also a subspace in \mathbb{R}^3 .



Intersection of subspaces

Theorem: Let $(U_i, i \in I)$ be any system of subspaces of a space V . The intersection of this system $\bigcap_{i \in I} U_i$ is also a subspace of V .

Proof: Let $W = \bigcap_{i \in I} U_i$. We show that W is closed on $+$ and \cdot .

$\forall \mathbf{u}, \mathbf{v} \in W :$

$$\mathbf{u}, \mathbf{v} \in W \Rightarrow \forall i \in I : \mathbf{u}, \mathbf{v} \in U_i \Rightarrow \forall i \in I : \mathbf{u} + \mathbf{v} \in U_i \Rightarrow \mathbf{u} + \mathbf{v} \in W$$

$\forall a \in \mathbb{K}, \mathbf{v} \in W :$

$$\mathbf{v} \in W \Rightarrow \forall i \in I : \mathbf{v} \in U_i \Rightarrow \forall i \in I : a\mathbf{v} \in U_i \Rightarrow a\mathbf{v} \in W$$

Note that when $I = \emptyset$, then the empty intersection

$$W = \bigcap_{i \in \emptyset} U_i = V \text{ is a subspace of } V \text{ itself.}$$

Linear combination, linear hull

Definition: A *linear combination* of vectors $\mathbf{v}_1, \dots, \mathbf{v}_k \in V$ over \mathbb{K} is any vector $\mathbf{u} = a_1 \mathbf{v}_1 + \dots + a_k \mathbf{v}_k$ where $a_1, \dots, a_k \in \mathbb{K}$.

Definition: The *linear hull* $\mathcal{L}(X)$ of a subset X of a vector space V is the intersection of all subspaces U of V that contain X .

Formally $\mathcal{L}(X) = \bigcap \{U : X \subseteq U, U \text{ is a subspace of } V\}$

$\mathcal{L}(X)$ is also called *the subspace of V generated by X* or the *linear span*. The linear hull is also denoted by $\text{span}(X)$.

Examples: For $V = \mathbb{R}^3$, $\mathcal{L}(\{(2, 2, 2)^T\}) = \{(a, a, a)^T, a \in \mathbb{R}\}$
... the line containing points having all three coordinates identical

$\mathcal{L}(\{(1, 0, 0)^T, (0, 1, 0)^T\}) = \{(a, b, 0)^T, a, b \in \mathbb{R}\}$
... the plane determined by the first two axes

Linear combination, linear hull

Definition: A *linear combination* of vectors $\mathbf{v}_1, \dots, \mathbf{v}_k \in V$ over \mathbb{K} is any vector $\mathbf{u} = a_1 \mathbf{v}_1 + \dots + a_k \mathbf{v}_k$ where $a_1, \dots, a_k \in \mathbb{K}$.

Definition: The *linear hull* $\mathcal{L}(X)$ of a subset X of a vector space V is the intersection of all subspaces U of V that contain X .

Formally $\mathcal{L}(X) = \bigcap \{U : X \subseteq U, U \text{ is a subspace of } V\}$

$\mathcal{L}(X)$ is also called *the subspace of V generated by X* or the *linear span*. The linear hull is also denoted by $\text{span}(X)$.

Theorem: Let V be a vector space over \mathbb{K} and X be a subset of V . Then $\mathcal{L}(X)$ is the set of all linear combinations of vectors from X .

Example for $W = \{\mathbf{x} : \mathbf{Ax} = \mathbf{0}\}$

$$\text{For } \mathbf{A} = \begin{pmatrix} 1 & -2 & 4 & 0 & 3 \\ -2 & 4 & -6 & 6 & 0 \\ 4 & -8 & 13 & -9 & -4 \\ 3 & -6 & 9 & -9 & 1 \end{pmatrix} \sim \sim \begin{pmatrix} 1 & -2 & 4 & 0 & 3 \\ 0 & 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

we solve $W = \{p_1(2, 1, 0, 0, 0)^T + p_2(12, 0, -3, 1, 0)^T : p_1, p_2 \in \mathbb{R}\}$

1. W could be viewed the *intersection* of the following subspaces:

$$U_1 = \{\mathbf{x} : (1, -2, 4, 0, 3) \cdot \mathbf{x} = 0\}$$

$$U_2 = \{\mathbf{x} : (-2, 4, -6, 6, 0) \cdot \mathbf{x} = 0\}$$

$$U_3 = \{\mathbf{x} : (4, -8, 13, -9, -4) \cdot \mathbf{x} = 0\}$$


$$U_4 = \{\mathbf{x} : (3, -6, 9, -9, 1) \cdot \mathbf{x} = 0\}$$

Each U_i corresponds to one equation of the system.

(Each is a *hyperplane* in \mathbb{R}^5 .)

2. The same subspace W was already described as a *set of linear combinations*
 $W = \mathcal{L}((2, 1, 0, 0, 0)^T, (12, 0, -3, 1, 0)^T)$

Correctness of Gaussian elimination and backward substitution yields that both ways describe the same space.



Proof of the theorem

Theorem: Let V be a vector space over \mathbb{K} and X be a subset of V . Then $\mathcal{L}(X)$ is the set of all linear combinations of vectors from X .

For $W_1 = \bigcap_{X \subseteq U_i \subseteq V} U_i$, $W_2 = \left\{ \sum_{i=1}^k a_i \mathbf{v}_i : k \in \mathbb{N}, a_i \in \mathbb{K}, \mathbf{v}_i \in X \right\}$.

we want to show $W_1 = \mathcal{L}(X) = W_2$.

W_2 is a subspace, because it is closed on scalar multiples

$$\mathbf{u} \in W_2 \Rightarrow \mathbf{u} = \sum_{i=1}^k a_i \mathbf{v}_i \Rightarrow \alpha \mathbf{u} = \alpha \sum_{i=1}^k a_i \mathbf{v}_i = \sum_{i=1}^k (\alpha a_i) \mathbf{v}_i \Rightarrow \alpha \mathbf{u} \in W_2$$

and analogously also on addition $\mathbf{u}, \mathbf{u}' \in W_2 \Rightarrow \dots \Rightarrow \mathbf{u} + \mathbf{u}' \in W_2$

Since $X \subseteq W_2$, we have W_2 among the intersecting subspaces U_i .

Hence $W_1 \subseteq W_2$.

Every U_i contains X and is closed on addition and scalar multiples.

Thus every U_i contains all linear combinations of vectors of X .

Hence $\forall U_i : W_2 \subseteq U_i \Rightarrow W_2 \subseteq W_1$.

Spaces determined by a matrix

Definition: The *kernel* of $\mathbf{A} \in \mathbb{K}^{m \times n}$ is $\ker(\mathbf{A}) = \{\mathbf{x} \in \mathbb{K}^n : \mathbf{A}\mathbf{x} = \mathbf{0}\}$,
the *row space* $\mathcal{R}(\mathbf{A}) \subseteq \mathbb{K}^n$ is the linear hull of the rows of \mathbf{A} ,
the *column space* $\mathcal{C}(\mathbf{A}) \subseteq \mathbb{K}^m$ is the linear hull of the columns of \mathbf{A} .

Example: Given $\mathbf{A} = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 2 & 0 & 2 & 1 \\ 1 & 1 & 2 & 0 \end{pmatrix} \in \mathbb{Z}_3^{3 \times 4}$

The row space:

$$\mathcal{R}(\mathbf{A}) = \left\{ \begin{array}{lll} (0, 0, 0, 0)^T, & (1, 2, 0, 1)^T, & (2, 1, 0, 2)^T, \\ (2, 0, 2, 1)^T, & (0, 2, 2, 2)^T, & (1, 1, 2, 0)^T, \\ (1, 0, 1, 2)^T, & (2, 2, 1, 0)^T, & (0, 1, 1, 1)^T \end{array} \right\} \subseteq \mathbb{Z}_3^4$$

The column space:

$$\mathcal{C}(\mathbf{A}) = \left\{ \begin{array}{lll} (0, 0, 0)^T, & (1, 2, 1)^T, & (2, 1, 2)^T, \\ (2, 0, 1)^T, & (0, 2, 2)^T, & (1, 1, 0)^T, \\ (1, 0, 2)^T, & (2, 2, 0)^T, & (0, 1, 1)^T \end{array} \right\} \subseteq \mathbb{Z}_3^3$$

Spaces determined by a matrix

Definition: The *kernel* of $\mathbf{A} \in \mathbb{K}^{m \times n}$ is $\ker(\mathbf{A}) = \{\mathbf{x} \in \mathbb{K}^n : \mathbf{A}\mathbf{x} = \mathbf{0}\}$, the *row space* $\mathcal{R}(\mathbf{A}) \subseteq \mathbb{K}^n$ is the linear hull of the rows of \mathbf{A} , the *column space* $\mathcal{C}(\mathbf{A}) \subseteq \mathbb{K}^m$ is the linear hull of the columns of \mathbf{A} .

Formally:

$$\mathcal{C}(\mathbf{A}) = \{\mathbf{u} \in \mathbb{K}^m : \mathbf{u} = \mathbf{A}\mathbf{x}, \mathbf{x} \in \mathbb{K}^n\}$$

$$\mathcal{R}(\mathbf{A}) = \{\mathbf{v} \in \mathbb{K}^n : \mathbf{v} = \mathbf{A}^T \mathbf{y}, \mathbf{y} \in \mathbb{K}^m\}$$

Observation: The kernel $\ker(\mathbf{A})$ is a subspace of \mathbb{K}^n .

Observation: Elementary transforms do not alter $\mathcal{R}(\mathbf{A})$ nor $\ker(\mathbf{A})$.

Observation: $\forall \mathbf{v} \in \mathcal{R}(\mathbf{A}), \forall \mathbf{x} \in \ker(\mathbf{A}) : \mathbf{v}^T \mathbf{x} = 0$.

Proof: $\mathbf{v}^T \mathbf{x} = (\mathbf{A}^T \mathbf{y})^T \mathbf{x} = \mathbf{y}^T \mathbf{A}\mathbf{x} = \mathbf{y}^T \mathbf{0} = 0$ for some $\mathbf{y} \in \mathbb{K}^m$.