

Spaces determined by a matrix

The *row space* $\mathcal{R}(\mathbf{A}) \subseteq \mathbb{K}^n$ is the linear hull of the rows of \mathbf{A} ,
the *column space* $\mathcal{C}(\mathbf{A}) \subseteq \mathbb{K}^m$ is the linear hull of the columns of \mathbf{A} .

Example: Given $\mathbf{A} = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 2 & 0 & 2 & 1 \\ 1 & 1 & 2 & 0 \end{pmatrix} \in \mathbb{Z}_3^{3 \times 4}$

The row space:

$$\mathcal{R}(\mathbf{A}) = \left\{ \begin{array}{lll} (0, 0, 0, 0)^T, & (\mathbf{1}, \mathbf{2}, \mathbf{0}, \mathbf{1})^T, & (2, 1, 0, 2)^T, \\ (\mathbf{2}, \mathbf{0}, \mathbf{2}, \mathbf{1})^T, & (0, 2, 2, 2)^T, & (\mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{0})^T, \\ (1, 0, 1, 2)^T, & (2, 2, 1, 0)^T, & (0, 1, 1, 1)^T \end{array} \right\} \subseteq \mathbb{Z}_3^4$$

The column space:

$$\mathcal{C}(\mathbf{A}) = \left\{ \begin{array}{lll} (0, 0, 0)^T, & (\mathbf{1}, \mathbf{2}, \mathbf{1})^T, & (2, 1, 2)^T, \\ (\mathbf{2}, \mathbf{0}, \mathbf{1})^T, & (\mathbf{0}, \mathbf{2}, \mathbf{2})^T, & (\mathbf{1}, \mathbf{1}, \mathbf{0})^T, \\ (1, 0, 2)^T, & (2, 2, 0)^T, & (0, 1, 1)^T \end{array} \right\} \subseteq \mathbb{Z}_3^3$$

Lemma: If $\mathbf{A}' = \mathbf{BA}$ then $\dim(\mathcal{C}(\mathbf{A}')) \leq \dim(\mathcal{C}(\mathbf{A}))$

Proof: Denote by $\mathbf{u}_1, \dots, \mathbf{u}_n$ the columns of \mathbf{A} .

The columns $\mathbf{u}'_1, \dots, \mathbf{u}'_n$ of \mathbf{A}' satisfy $\mathbf{u}'_i = \mathbf{B}\mathbf{u}_i$.

When $\mathbf{w}' \in \mathcal{C}(\mathbf{A}')$ then $\mathbf{w}' = \sum_{i=1}^n a_i \mathbf{u}'_i$ for some $a_1, \dots, a_n \in \mathbb{K}$.

$$\mathbf{w}' = \sum_{i=1}^n a_i \mathbf{B}\mathbf{u}_i = \mathbf{B} \sum_{i=1}^n a_i \mathbf{u}_i = \mathbf{B}\mathbf{w} \text{ for } \mathbf{w} = \sum_{i=1}^n a_i \mathbf{u}_i \in \mathcal{C}(\mathbf{A})$$

W.l.o.g. $\mathbf{u}_1, \dots, \mathbf{u}_d$ form a basis of $\mathcal{C}(\mathbf{A})$, $d = \dim(\mathcal{C}(\mathbf{A}))$

thus $\mathbf{w} = \sum_{i=1}^d b_i \mathbf{u}_i$ for some $b_1, \dots, b_d \in \mathbb{K}$.

$$\text{Now } \mathbf{w}' = \mathbf{B}\mathbf{w} = \mathbf{B} \sum_{i=1}^d b_i \mathbf{u}_i = \sum_{i=1}^d b_i \mathbf{B}\mathbf{u}_i = \sum_{i=1}^d b_i \mathbf{u}'_i,$$

i.e. $\mathbf{u}'_1, \dots, \mathbf{u}'_d$ generate $\mathcal{C}(\mathbf{A}')$. Thus

$$\dim(\mathcal{C}(\mathbf{A}')) \leq d = \dim(\mathcal{C}(\mathbf{A})).$$

Both spaces have the same dimension



Theorem: Any $\mathbf{A} \in \mathbb{K}^{m \times n}$ satisfies: $\dim(\mathcal{R}(\mathbf{A})) = \dim(\mathcal{C}(\mathbf{A}))$.

Proof: Let $\mathbf{A} \sim \mathbf{A}'$ in REF, i.e. there is a regular \mathbf{R} s.t. $\mathbf{A}' = \mathbf{R}\mathbf{A}$.
By the lemma $\dim(\mathcal{C}(\mathbf{A}')) \leq \dim(\mathcal{C}(\mathbf{A}))$.

From $\mathbf{A} = \mathbf{R}^{-1}\mathbf{A}'$ we get $\dim(\mathcal{C}(\mathbf{A}')) \geq \dim(\mathcal{C}(\mathbf{A}))$ and indeed $=$.

For matrices \mathbf{A}' in REF the theorem holds directly:

$\dim(\mathcal{R}(\mathbf{A}')) = \# \text{ of pivots} = \text{rank}(\mathbf{A}') = \dim(\mathcal{C}(\mathbf{A}'))$



Since $\mathcal{R}(\mathbf{A}) = \mathcal{R}(\mathbf{A}')$, we get

$\dim(\mathcal{R}(\mathbf{A})) = \dim(\mathcal{R}(\mathbf{A}')) = \dim(\mathcal{C}(\mathbf{A}')) = \dim(\mathcal{C}(\mathbf{A}))$.

Example

Use Gauss-Jordan elimination:

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 2 & 0 & 2 & 1 \\ 1 & 1 & 2 & 0 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} = \mathbf{A}'$$

$\mathcal{R}(\mathbf{A}) = \mathcal{R}(\mathbf{A}')$ yields $\dim(\mathcal{R}(\mathbf{A})) = \dim(\mathcal{R}(\mathbf{A}')) = \text{rank}(\mathbf{A}) = 2$.

From $\mathbf{A}' = \begin{pmatrix} 0 & 2 & 0 \\ 2 & 2 & 0 \\ 1 & 2 & 1 \end{pmatrix} \mathbf{A}$ we get $\dim(\mathcal{C}(\mathbf{A}')) \leq \dim(\mathcal{C}(\mathbf{A}))$.

The matrix of the transformation is regular — it has an inverse, so

from $\mathbf{A} = \begin{pmatrix} 1 & 2 & 0 \\ 2 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix} \mathbf{A}'$ we get $\dim(\mathcal{C}(\mathbf{A})) \leq \dim(\mathcal{C}(\mathbf{A}'))$.

The space $\mathcal{C}(\mathbf{A}')$ is generated by the **columns with pivots**, here by the first two vectors of the standard basis. Hence $\dim(\mathcal{C}(\mathbf{A}')) = 2$. In particular $\mathcal{C}(\mathbf{A}') = \{(p_1, p_2, 0)^T, p_1, p_2 \in \mathbb{Z}_3\}$.

Consequences

- ▶ $\text{rank}(\mathbf{A}^T) = \text{rank}(\mathbf{A})$
- ▶ When \mathbf{R} and \mathbf{R}' are regular of order $m \times m$ and $n \times n$, resp. then $\text{rank}(\mathbf{RA}) = \text{rank}(\mathbf{AR}') = \text{rank}(\mathbf{A})$
- ▶ $\mathcal{R}(\mathbf{BA}) \subseteq \mathcal{R}(\mathbf{A})$, $\mathcal{C}(\mathbf{BA}) \subseteq \mathcal{C}(\mathbf{B})$
- ▶ $\text{rank}(\mathbf{BA}) \leq \min\{\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B})\}$

Theorem: For any $\mathbf{A} \in \mathbb{K}^{m \times n}$: $\dim(\ker(\mathbf{A})) + \text{rank}(\mathbf{A}) = n$

Proof: Let $d = n - \text{rank}(\mathbf{A})$ is the number of free variables and $\mathbf{x}_1, \dots, \mathbf{x}_d$ the suitable solutions of $\mathbf{Ax} = \mathbf{0}$.

These solutions are linearly independent as each \mathbf{x}_i is the only one which has a nonzero coefficient by the i -th free variable.

Hence $\mathbf{x}_1, \dots, \mathbf{x}_d$ form a basis of $\ker(\mathbf{A})$ and $\dim(\ker(\mathbf{A})) = d = n - \text{rank}(\mathbf{A})$