

MATHEMATICAL ANALYSIS 3 (NMAI056)

summer term 2025/26

lecturer: Martin Klazar

**LECTURE 2 (February 23, 2026) OSTROWSKI'S
THEOREM. COMPACT METRIC SPACES.**

• *Ostrowski's theorem.* On any field F we have the *trivial norm*. It is a function $\|\cdot\|$ with $\|0_F\| = 0$ and $\|x\| = 1$ for $x \neq 0_F$.

Exercise 1 *Prove that a trivial norm is a norm.*

From the usual absolute value $|\cdot|$ on \mathbb{Q} , \mathbb{R} and \mathbb{C} , we get many other norms by exponentiation.

Exercise 2 *Prove that for any $c > 0$, $|\cdot|^c$ is a norm (on \mathbb{Q} , \mathbb{R} , and \mathbb{C}) if and only if $c \leq 1$. We will call this norm the *modified absolute value*.*

For $\alpha \in \mathbb{Q}$ and a prime p , the *canonical p -adic norm* $\|\cdot\|_p$ is defined by

$$\|\alpha\|_p = p^{-\text{ord}_p(\alpha)}$$

– in the general p -adic norm $|\cdot|_p$ we set $c = 1/p$.

Exercise 3 *Let $M = \{2, 3, 5, 7, 11, \dots\} \cup \{\infty\}$ and $\|\cdot\|_\infty = |\cdot|$ (ordinary absolute value). Prove that for every nonzero number $\alpha \in \mathbb{Q}$ we have the product formula*

$$\prod_{p \in M} \|\alpha\|_p = 1.$$

Exercise 4 *Let $\|\cdot\|$ be a nontrivial norm on the field \mathbb{Q} . Prove that $\exists n \in \mathbb{N}$ ($n \geq 2 \wedge \|n\| \neq 1$).*

Exercise 5 Prove that for every two coprime numbers $a, b \in \mathbb{Z}$ there exist numbers $c, d \in \mathbb{Z}$ such that

$$ac + db = 1 .$$

Theorem 6 (A. Ostrowski, 1916) Let $\|\cdot\|$ be a norm on the field \mathbb{Q} . Then (exactly) one of three cases occurs.

1. $\|\cdot\|$ is a trivial norm.
2. There exists a number $c \in (0, 1]$ such that $\|x\| = |x|^c$.
3. There exists a number $c \in (0, 1)$ and a prime number p such that $\|x\| = |x|_p = c^{\text{ord}_p(x)}$.

Thus modified absolute values and p -adic norms are the only non-trivial norms on the field of fractions.

Proof. Let $\|\cdot\|$ be a nontrivial norm. By Exercise 4 there exists an $n \in \mathbb{N} \setminus \{1\}$ such that $\|n\| \neq 1$. Two cases occur.

1. There is an $n \in \mathbb{N}$ such that $\|n\| > 1$. Let n_0 be the smallest such n . Clearly, $n_0 \geq 2$ a

$$1 \leq m < n_0 \Rightarrow \|m\| \leq 1 . \tag{1}$$

We define a (unique) real number $c > 0$ by

$$\|n_0\| = n_0^c . \tag{2}$$

Every number $n \in \mathbb{N}$ has a (unique) base n_0 expansion:

$$n = a_0 + a_1 n_0 + a_2 n_0^2 + \cdots + a_s n_0^s \text{ where} \\ a_i, s \in \mathbb{N}_0, 0 \leq a_i < n_0 \text{ and } a_s \neq 0 .$$

For $n_0 = 10$ this is the usual decadic notation: for example, $2024 = 2 \cdot 10^3 + 0 \cdot 10^2 + 2 \cdot 10^1 + 4 \cdot 10^0$. We have

$$\begin{aligned}
\|n\| &= \|a_0 + a_1 n_0 + a_2 n_0^2 + \cdots + a_s n_0^s\| \\
&\stackrel{\Delta\text{-ineq. and multipl. of } \|\cdot\|}{\leq} \sum_{j=0}^s \|a_j\| \cdot \|n_0\|^j \\
&\stackrel{\text{eq. (1) and (2)}}{\leq} \sum_{j=0}^s n_0^{jc} \leq n_0^{sc} \sum_{i=0}^{\infty} (1/n_0^c)^i \\
&\stackrel{n_0^s \leq n}{\leq} n^c C \text{ where } C = \sum_{i=0}^{\infty} (1/n_0^c)^i .
\end{aligned}$$

Hence

$$\forall n \in \mathbb{N}_0 : \|n\| \leq C n^c . \quad (3)$$

This in fact holds with $C = 1$. Let $m, n \in \mathbb{N}$. By the multiplicativity of the norm and inequality (3),

$$\|n\|^m = \|n^m\| \leq C (n^m)^c = C (n^c)^m .$$

We take the m -th root and get $\|n\| \leq C^{1/m} n^c$. For $m \rightarrow \infty$ we have $C^{1/m} \rightarrow 1$. Thus

$$\forall n \in \mathbb{N}_0 (\|n\| \leq n^c) . \quad (4)$$

We derive the reverse inequality $\|n\| \geq n^c$, $n \in \mathbb{N}_0$. For every $n \in \mathbb{N}$, the above base n_0 expansion of n yields

$$n_0^{s+1} > n \geq n_0^s .$$

By the Δ -inequality,

$$\|n_0\|^{s+1} = \|n_0^{s+1}\| \leq \|n\| + \|n_0^{s+1} - n\| .$$

Hence

$$\begin{aligned}
\|n\| &\geq \|n_0\|^{s+1} - \|n_0^{s+1} - n\| \stackrel{(2) \text{ and } (4)}{\geq} n_0^{(s+1)c} - (n_0^{s+1} - n)^c \\
&\stackrel{n \geq n_0^s}{\geq} n_0^{(s+1)c} - (n_0^{s+1} - n_0^s)^c = n_0^{(s+1)c} \left(1 - \left(1 - \frac{1}{n_0}\right)^c\right) \\
&\stackrel{n_0^{s+1} > n}{\geq} n^c C' \text{ where } C' = 1 - \left(1 - \frac{1}{n_0}\right)^c > 0.
\end{aligned}$$

The trick with the m -th root gives again

$$\forall n \in \mathbb{N}_0 \left(\|n\| \geq n^c \right).$$

Hence

$$\forall n \in \mathbb{N}_0 \left(\|n\| = n^c \right).$$

From the multiplicativity of the norm $\|\cdot\|$, we get that $\|x\| = |x|^c$ for any $x \in \mathbb{Q}$. By Exercise 2, $c \in (0, 1]$. We get case 2 of Ostrowski's theorem.

2. $\forall n \in \mathbb{N} : \|n\| \leq 1$ and $\exists n \in \mathbb{N} : \|n\| < 1$. Let n_0 be minimum such n ; again $n_0 \geq 2$. We claim that $n_0 = p$ is a prime number. Indeed, if we could express n_0 as $n_0 = n_1 n_2$ with $n_i \in \mathbb{Z}$ and $1 < n_1, n_2 < n_0$, we get the contradiction

$$1 > \|n_0\| = \|n_1 n_2\| = \|n_1\| \cdot \|n_2\| = 1 \cdot 1 = 1$$

— we used the multiplicativity of norms and that $\|m\| = 1$ for any $m \in \mathbb{N}$ with $1 \leq m < n_0$. We show that every prime number q different from p has the norm $\|q\| = 1$. For the contrary, let $q \neq p$ be another prime number with $\|q\| < 1$. We take a large $m \in \mathbb{N}$ such that $\|p\|^m, \|q\|^m < \frac{1}{2}$. By Exercise 5 there exist integers a and

b such that $aq^m + bp^m = 1$. Then

$$\begin{aligned} 1 &= \|1\| = \|aq^m + bp^m\| \leq \|a\| \cdot \|q\|^m + \|b\| \cdot \|p\|^m \\ &< 1 \cdot \frac{1}{2} + 1 \cdot \frac{1}{2} = 1 \end{aligned}$$

which is a contradiction; we used the triangle inequality, multiplicativity of norms, and the fact that now $\|a\| \leq 1$ for every $a \in \mathbb{Z}$.

Thus $\|q\| = 1$ for every prime number $q \neq p$. From this, multiplicativity of norms and prime factorization of any non-zero fraction x we get the expression

$$\begin{aligned} \|x\| &= \left\| \prod_{q=2,3,5,\dots} q^{\text{ord}_q(x)} \right\| = \prod_{q=2,3,5,\dots} \|q\|^{\text{ord}_q(x)} = \|p\|^{\text{ord}_p(x)} \\ &= c^{\text{ord}_p(x)}, \text{ where } c = \|p\| \in (0, 1). \end{aligned}$$

Also $\|0\| = c^{\text{ord}_p(0)} = c^\infty = 0$. Thus we get case 3 of Ostrowski's theorem. \square

The preceding proof is taken from the book by Neal Koblitz,

p-adic Numbers, p-adic Analysis, and Zeta-Functions, Springer-Verlag, New York, 1984.

This book contains lot of information about the p -adic norm $\|\cdot\|_p$ and related p -adic analysis.

- *Compact sets in metric spaces.* We review limits of sequences in metric spaces (MS). Let (M, d) be a MS, $(a_n) \subset M$ be a sequence of points in it and $a \in M$ be a point. We say that (a_n) has the *limit* a in the MS (M, d) if for every ε there is an n_0 such that

$$n \geq n_0 \Rightarrow d(a_n, a) < \varepsilon.$$

From now on $\varepsilon > 0$ is a real number and $n_0, n \in \mathbb{N}$. We write that $\lim a_n = a$ or $\lim_{n \rightarrow \infty} a_n = a$. If the sequence (a_n) has a limit, we say that it is *convergent*, otherwise it is *divergent*.

Let (M, d) be a MS and $X \subset M$. We say that the set X is *compact* if every sequence $(a_n) \subset X$ has a subsequence (a_{m_n}) such that

$$\lim_{n \rightarrow \infty} a_{m_n} = a \in X.$$

The MS (M, d) is *compact* when the set M is compact.

The Bolzano–Weierstrass theorem states that in the MS

$$(\mathbb{R}, |x - y|)$$

every closed and bounded interval $X = [a, b]$ is a compact set. We give a few examples of compact sets and compact MSs.

Exercise 7 *In every MS every finite set is compact.*

Exercise 8 *Is the real axis $(\mathbb{R}, |x - y|)$ a compact MS?*

Exercise 9 *Which other intervals on the real axis besides $[a, b]$ are compact sets?*

Exercise 10 *Let $X = [a, b] \times [c, d]$ be a rectangle in the plane, that is, in the Euclidean space (\mathbb{R}^2, e_2) . Prove that X is compact.*

Exercise 11 *Let (M, d) be MS and $A, B \subset M$. Which of the following implications holds?*

A and B are compact $\Rightarrow A \cup B$ is compact

A and B are compact $\Rightarrow A \cap B$ is compact

$A \subset B$ and B is compact $\Rightarrow A$ is compact

A and B are compact $\Rightarrow A \setminus B$ is compact

• *Extrema and compact sets.* We begin with continuous maps between MSs. Let (M, d) and (N, e) be MS. A map $f: M \rightarrow N$ is *continuous at the point* $a \in M$ if for every ε there is a δ such that

$$x \in M \wedge d(x, a) < \delta \Rightarrow e(f(x), f(a)) < \varepsilon$$

($\delta > 0$ is a real number). A map is *continuous* if it is continuous at every point of the space.

Exercise 12 *Let $f: M \rightarrow N$ be a map between metric spaces and $a \in M$. Prove Heine's definition of continuity at a point:*

$$\begin{aligned} f \text{ is continuous at } a &\iff \\ \iff \forall (a_n) \subset M : \lim a_n = a &\Rightarrow \lim f(a_n) = f(a). \end{aligned}$$

Theorem 13 (extremal values) *Let (M, d) be MS, $f: M \rightarrow \mathbb{R}$ be a continuous function, and $X \subset M$ be a nonempty compact set. Then there exist $a, b \in X$ such that*

$$\forall x \in X : f(a) \leq f(x) \leq f(b).$$

Thus f attains on the set X both the smallest value $f(a)$ and the largest value $f(b)$.

Proof. We show that $f[X] = \{f(x) : x \in X\}$ is a bounded subset of \mathbb{R} . If $f[X]$ were not bounded from above, we could take a sequence $(a_n) \subset X$ with $\lim f(a_n) = +\infty$. By the assumption, (a_n) has a convergent subsequence (a_{m_n}) with

$$\lim a_{m_n} = a \in X.$$

By the continuity of f at a and Exercise 12, $\lim f(a_{m_n}) = f(a) \in \mathbb{R}$. This is a contradiction because $\lim f(a_{m_n}) = +\infty$. Boundedness of $f[X]$ from below follows in a similar way.

So we can define the real numbers $A = \inf(f[X])$ and $B = \sup(f[X])$. By the definition of infima, there is a sequence $(a_n) \subset X$ such that $\lim f(a_n) = A$. By the assumption, (a_n) has a convergent subsequence (a_{m_n}) with $\lim a_{m_n} = a \in X$. By the continuity of f at a and Exercise 12, $\lim f(a_{m_n}) = f(a)$. Subsequences preserve limits and $\lim f(a_{m_n}) = A$. Thus $f(a) = A$ and for every $x \in X$,

$$f(a) = A \leq f(x)$$

because $A = \inf(f[X])$. The other element $b \in X$ is obtained similarly. \square

• *Products of metric spaces.* For metric spaces (M, d) and (N, e) , we define their *product* $(M \times N, d \times e)$ as follows. $M \times N$ is the Cartesian product of the sets M and N and the $d \times e$ metric on it is given by

$$(d \times e)((a_1, a_2), (b_1, b_2)) = \sqrt{d(a_1, b_1)^2 + e(a_2, b_2)^2}.$$

Exercise 14 *Prove that the product of two MSs is a MS.*

Exercise 15 *Prove that the product of two Euclidean MSs*

$$(\mathbb{R}^m, e_m) \text{ and } (\mathbb{R}^n, e_n)$$

is (except for a formality in notation) the Euclidean MS

$$(\mathbb{R}^{m+n}, e_{m+n}).$$

What is the “formality”?

• *Characterization of compact sets in Euclidean MSs.* We defined the ball $B(a, r)$ in a MS last time. A set $X \subset M$ in a MS (M, d) is *open* if

$$\forall a \in X \exists r : B(a, r) \subset X.$$

Here $r > 0$ is a real number, the radius of the ball $B(a, r)$. X is *closed* if $M \setminus X$ is open. X is *bounded* if

$$\exists a \in M \exists r : X \subset B(a, r).$$

The *diameter* of the set X is, with $V = \{d(a, b) : a, b \in X\}$ ($\subset [0, +\infty)$), defined as

$$\text{diam}(X) := \begin{cases} \sup(V) & \dots \text{ the set } V \text{ is bounded from above and} \\ +\infty & \dots \text{ the set } V \text{ is unbounded from above.} \end{cases}$$

Exercise 16 *Prove that any set X is bounded if and only if $\text{diam}(X) < +\infty$.*

Exercise 17 *Prove that for every unbounded set X there exists a sequence $(a_n) \subset X$ such that $m < n \Rightarrow d(a_m, a_n) \geq 1$.*

In the following two exercises we review basic properties of open and closed sets in a MS.

Exercise 18 *Let (M, d) be a MS. Then the following holds.*

1. *The sets \emptyset and M are both open and closed.*
2. *Any finite intersection of open subsets of M is an open set and any finite union of closed subsets of M is a closed set.*
3. *Any union of open subsets of M is an open set and any intersection of closed subsets of M is a closed set.*

Exercise 19 *Let (M, d) be a MS and $X \subset M$. Then*

$$\begin{aligned} & \text{the set } X \text{ is closed} \iff \\ & \iff \forall (a_n) \subset X \forall a \in M \left(\lim a_n = a \Rightarrow a \in X \right). \end{aligned}$$

Theorem 20 *Let (M, d) and (N, e) be MS. The following holds.*

1. *If $X \subset M$ is a compact set, then X is closed and bounded. The opposite implication does not hold, see Exercise 22.*
2. *If (M, d) and (N, e) are compact, then the product $(M \times N, d \times e)$ is compact too.*

Proof. 1. If X is not closed, then by Exercise 19 there exists a convergent sequence $(a_n) \subset X$ such that $\lim a_n = a \in M \setminus X$. This sequence does not have a convergent subsequence with limit in X , since each subsequence has limit a . When X is not bounded, we easily construct a sequence $(a_n) \subset X$ such that $m < n \Rightarrow d(a_m, a_n) > 1$ (Exercise 17). This sequence clearly has no convergent subsequence.

2. Let $(a_n) = ((a_{n,1}, a_{n,2}))$ be a sequence in the product MS. We choose a subsequence (b_n) such that $(b_{n,1})$ has a limit $b \in M$ in (M, d) . From (b_n) we select a subsequence (c_n) such that $(c_{n,2})$ has a limit $c \in N$ in (N, e) . It is not difficult to see that (c_n) is a subsequence of the sequence (a_n) and that it has in the product MS the limit

$$\lim c_n = (b, c) \in M \times N .$$

□

Exercise 21 *Let (M, d) be a compact MS and $X \subset M$ be a closed set. Prove that X is compact.*

Exercise 22 *Let M be an infinite set and the metric d on it is given as $d(a, b) = 1$ for $a \neq b$ and $d(a, a) = 0$. Show that (M, d) is a MS that is bounded and closed but not compact.*

Theorem 23 (compact sets in \mathbb{R}^n) *In every Euclidean MS (\mathbb{R}^n, e_n) , a set $X \subset \mathbb{R}^n$ is compact if and only if it is bounded and closed.*

Proof. By the first part of the previous theorem, it suffices to prove that every bounded and closed set $X \subset \mathbb{R}^n$ is compact. From its boundedness it follows that for a real number $a > 0$,

$$X \subset K = [-a, a]^n = [-a, a] \times [-a, a] \times \cdots \times [-a, a] \subset \mathbb{R}^n .$$

The Euclidean MS (K, e_n) is compact by the Bolzano–Weierstrass theorem, part 2 of the previous theorem, and Exercise 15. Clearly, X is also closed in (K, e_n) (problem 24), so according to Exercise 21, X is compact in (K, e_n) and therefore in (\mathbb{R}^n, e_n) (Exercise 25). \square

Exercise 24 *Let (M, d) be a MS, $A \subset B \subset M$ and A be a closed set in $(M, d) \Rightarrow A$ is closed also in the subspace (B, d) .*

Exercise 25 *Let (M, d) be a MS and $A \subset B \subset M$. Then A is compact in $(M, d) \iff A$ is compact in the subspace (B, d) .*

THANK YOU FOR YOUR ATTENTION

Homework Exercises. Please send to me (klazar@kam.mff.cuni.cz) by the end of the coming Sunday solutions to the Exercises 5, 9, 11, 17 and 22.