Exercises for Combinatorial and Computational Geometry 2 Series 2 - k-holes, halving lines and graph drawings

deadline 1. 4. 2025

1. Let X be a finite non-empty set of points in the plane in general position. Prove that the following identities hold for the numbers of k-holes in X:

(a)
$$\sum_{k=1}^{|X|} (-1)^k \cdot \#k\text{-holes} = -1,$$
(b) If $|X| \ge 2$, then
$$\sum_{k=1}^{|X|} (-1)^k \cdot k \cdot \#k\text{-holes} = -\#\text{points inside conv}(X).$$
[2]

Hint: move points continuously along curves into a suitable configuration.

- 2. Let X be a set of n points in the plane in general position. Prove that the number of 4-holes is at least quadratic in n. [2]
- 3. Let P be a finite set of points in the plane that is not necessarily in general position and that contains no 5-hole. Prove that every convex 5-gon Q determined by points of P contains at least 1 point of P in the closed "inner" 5-gon that is determined by the diagonals of Q. [2]
- 4. For *n* even, let *P* be a set of *n* points in the plane in general position. Furthermore, let $k \leq n/2$ and let *h* be a line that does not intersect *P* and splits the plane into two halfplanes such that one of them contains exactly *k* points of *P*. Show that *h* intersects exactly *k* halving segments of *P*. [2]
- 5. Let P be a set of n points in the plane in general position. A pair of points in P is a k-edge if the line determined by these points separates exactly k points from P in one of the open hyperplanes. Let $E_k(P)$ be the number of k-edges in P and furthermore let $\overline{cr}(P)$ be the number of unordered 4-tuples of points in P in convex position.

Prove the following identity:

$$\overline{cr}(P) = 3\binom{n}{4} - \sum_{k=0}^{\lfloor \frac{n-2}{2} \rfloor} E_k(P) \cdot k \cdot (n-k-2).$$

Hint: count in two ways the number of ordered triples (a, \overline{bc}, d) such that a, b, c, d are pairwise different points of P, a lies to the left of the line \overline{bc} and d lies to the right of the line of \overline{bc} . [2]

6. Two edges in a graph are *independent* if they do not share a vertex. In a *drawing* of a graph we assume that edges have only finitely many points in common, and that every common point of two edges is either their common endpoint or a crossing.

Determine for which values of m and n, in every drawing of the graph $K_{m,n}$ the total number of crossings of independent pairs of edges is odd. Hint: what happens during a continuous deformation of the edges? [3]

web: https://kam.mff.cuni.cz/~kvgweb/kvg/eng.html