

Using Dynamical Systems to Construct Infinitely Many Primes	483
Recent Results	487
Recent Progress in Additive Combinatorics	497
Recent Progress in Algebraic Combinatorics	505
Recent Progress in Algebraic Combinatorics	513
Recent Progress in Algebraic Combinatorics	521
Recent Progress in Algebraic Combinatorics	529
Recent Progress in Algebraic Combinatorics	537
Recent Progress in Algebraic Combinatorics	545
Recent Progress in Algebraic Combinatorics	553
Recent Progress in Algebraic Combinatorics	561
Recent Progress in Algebraic Combinatorics	569
Recent Progress in Algebraic Combinatorics	577
Recent Progress in Algebraic Combinatorics	585
Recent Progress in Algebraic Combinatorics	593
Recent Progress in Algebraic Combinatorics	601
Recent Progress in Algebraic Combinatorics	609
Recent Progress in Algebraic Combinatorics	617
Recent Progress in Algebraic Combinatorics	625
Recent Progress in Algebraic Combinatorics	633
Recent Progress in Algebraic Combinatorics	641
Recent Progress in Algebraic Combinatorics	649
Recent Progress in Algebraic Combinatorics	657
Recent Progress in Algebraic Combinatorics	665
Recent Progress in Algebraic Combinatorics	673
Recent Progress in Algebraic Combinatorics	681
Recent Progress in Algebraic Combinatorics	689
Recent Progress in Algebraic Combinatorics	697
Recent Progress in Algebraic Combinatorics	705
Recent Progress in Algebraic Combinatorics	713
Recent Progress in Algebraic Combinatorics	721
Recent Progress in Algebraic Combinatorics	729
Recent Progress in Algebraic Combinatorics	737
Recent Progress in Algebraic Combinatorics	745
Recent Progress in Algebraic Combinatorics	753
Recent Progress in Algebraic Combinatorics	761
Recent Progress in Algebraic Combinatorics	769
Recent Progress in Algebraic Combinatorics	777
Recent Progress in Algebraic Combinatorics	785
Recent Progress in Algebraic Combinatorics	793
Recent Progress in Algebraic Combinatorics	801
Recent Progress in Algebraic Combinatorics	809
Recent Progress in Algebraic Combinatorics	817
Recent Progress in Algebraic Combinatorics	825
Recent Progress in Algebraic Combinatorics	833
Recent Progress in Algebraic Combinatorics	841
Recent Progress in Algebraic Combinatorics	849
Recent Progress in Algebraic Combinatorics	857
Recent Progress in Algebraic Combinatorics	865
Recent Progress in Algebraic Combinatorics	873
Recent Progress in Algebraic Combinatorics	881
Recent Progress in Algebraic Combinatorics	889
Recent Progress in Algebraic Combinatorics	897
Recent Progress in Algebraic Combinatorics	905
Recent Progress in Algebraic Combinatorics	913
Recent Progress in Algebraic Combinatorics	921
Recent Progress in Algebraic Combinatorics	929
Recent Progress in Algebraic Combinatorics	937
Recent Progress in Algebraic Combinatorics	945
Recent Progress in Algebraic Combinatorics	953
Recent Progress in Algebraic Combinatorics	961
Recent Progress in Algebraic Combinatorics	969
Recent Progress in Algebraic Combinatorics	977
Recent Progress in Algebraic Combinatorics	985
Recent Progress in Algebraic Combinatorics	993
Recent Progress in Algebraic Combinatorics	1001

The American Mathematical Monthly

ISSN: 0002-9890 (Print) 1930-0972 (Online) Journal homepage: <http://maa.tandfonline.com/loi/uamm20>

Using Dynamical Systems to Construct Infinitely Many Primes

Andrew Granville

To cite this article: Andrew Granville (2018) Using Dynamical Systems to Construct Infinitely Many Primes, The American Mathematical Monthly, 125:6, 483-496, DOI: 10.1080/00029890.2018.1447732

To link to this article: <https://doi.org/10.1080/00029890.2018.1447732>



Published online: 24 May 2018.



Submit your article to this journal [↗](#)



Article views: 866



View Crossmark data [↗](#)

Using Dynamical Systems to Construct Infinitely Many Primes

Andrew Granville 

Abstract. Euclid's proof can be reworked to construct infinitely many primes, in many different ways, using ideas from arithmetic dynamics.

1. CONSTRUCTING INFINITELY MANY PRIMES. The first known proof that there are infinitely many primes, which appears in Euclid's *Elements*, is a proof by contradiction. This proof does not indicate how to find primes, despite establishing that there are infinitely many of them. However a minor variant does do so. The key to either proof is the theorem that

Every integer $q > 1$ has a prime factor.

(This was first formally proved by Euclid on the way to establishing the fundamental theorem of arithmetic.) The idea is to

Construct an infinite sequence of distinct, pairwise coprime, integers a_0, a_1, \dots ,

that is, a sequence for which $\gcd(a_m, a_n) = 1$ whenever $m \neq n$. We then obtain infinitely many primes, the prime factors of the a_n , as proved in the following result.

Proposition 1. *Suppose that a_0, a_1, \dots is an infinite sequence of distinct, pairwise coprime, integers. Let p_n be a prime divisor of a_n whenever $|a_n| > 1$. Then the p_n form an infinite sequence of distinct primes.*

Proof. The p_n are distinct for if not, then $p_m = p_n$ for some $m \neq n$ and so

$$p_m = \gcd(p_m, p_n) \text{ divides } \gcd(a_m, a_n) = 1,$$

a contradiction. As the a_n are distinct, any given value (in particular, $-1, 0$, and 1) can be attained at most once, and therefore all but at most three of the a_n have absolute value > 1 , and so have a prime factor p_n . ■

To construct such a sequence by modifying Euclid's proof, let $E_0 = 2$ and

$$E_n = E_0 E_1 \cdots E_{n-1} + 1 \text{ for each } n \geq 1.$$

If $m < n$, then E_m divides $E_0 E_1 \cdots E_{n-1} = E_n - 1$ (as E_m is one of the terms in the product) and so $\gcd(E_m, E_n)$ divides $\gcd(E_n - 1, E_n) = 1$, which implies that $\gcd(E_m, E_n) = 1$. Therefore, if p_n is a prime divisor of E_n for each $n \geq 0$, then p_0, p_1, \dots is an infinite sequence of distinct primes.

The *Fermat numbers*, F_0, F_1, \dots , defined by $F_n = 2^{2^n} + 1$ for each $n \geq 0$, are a more familiar sequence of pairwise coprime integers. Fermat had actually conjectured that the F_n are all primes. His claim starts off correct: $3, 5, 17, 257, 65537$ are all prime, but is false for $F_5 = 641 \times 6700417$, as Euler famously noted. It is an open

doi.org/10.1080/00029890.2018.1447732

MSC: Primary 11N05, Secondary 37P05

question as to whether there are infinitely many primes of the form F_n .¹ Nonetheless we can prove that the F_n are pairwise coprime in a similar way to the E_n , using the identity

$$F_n = F_0 F_1 \cdots F_{n-1} + 2 \text{ for each } n \geq 1$$

and the fact that the F_n are all odd. We deduce from Proposition 1 that if p_n is a prime divisor of F_n for each $n \geq 0$, then p_0, p_1, \dots is an infinite sequence of distinct primes.²

For any sequence of integers $(a_n)_{n \geq 0}$, we call p_n a *private prime factor* of a_n if it divides a_n and no other a_m . The primes p_n constructed in Proposition 1 are therefore each private prime factors of the a_n .

2. A SIMPLER FORMULATION. In Section 1, the sequences $(E_n)_{n \geq 0}$ and $(F_n)_{n \geq 0}$ were constructed in a similar way, multiplying all of the terms of the sequence so far together and adding a constant. We would like a simpler, unified way to view these two sequences, with an eye to generalization. This is not difficult because the recurrence for the E_n can be rewritten as

$$E_{n+1} = E_0 E_1 \cdots E_{n-1} \cdot E_n + 1 = (E_n - 1)E_n + 1 = f(E_n),$$

where $f(x) = x^2 - x + 1$. Similarly, the F_n -values can be determined by

$$F_{n+1} = (2^{2^n} + 1)(2^{2^n} - 1) + 2 = F_n(F_n - 2) + 1 = f(F_n),$$

where $f(x) = x^2 - 2x + 2$. So they are both examples of sequences $(x_n)_{n \geq 0}$ for which

$$x_{n+1} = f(x_n)$$

for some polynomial $f(x) \in \mathbb{Z}[x]$. The terms of the sequence are all given by the recursive formula, so that

$$x_n = \underbrace{f(f(\dots f(x_0)))}_{n \text{ times}} = f^n(x_0),$$

where the notation f^n denotes the polynomial obtained by composing f with itself n times (which is definitely *not* the n th power of f). Any such sequence $(x_n)_{n \geq 0}$ is called the *orbit* of x_0 under the map f , since the sequence is completely determined once one knows x_0 and f . We sometimes write the orbit as $x_0 \rightarrow x_1 \rightarrow x_2 \rightarrow \dots$.

The key to the proof that the E_n 's are pairwise coprime is that $E_n \equiv 1 \pmod{E_m}$ whenever $n > m \geq 0$; and the key to the proof that the F_n 's are pairwise coprime is that $F_n \equiv 2 \pmod{F_m}$ whenever $n > m \geq 0$. These congruences are not difficult to deduce using the following lemma from elementary number theory.

¹ The only Fermat numbers known to be primes have $n \leq 4$. We know that the F_n are composite for $5 \leq n \leq 30$ and for many other n besides. It is always a significant moment when a Fermat number is factored for the first time. It could be that all F_n with $n > 4$ are composite, or they might all be prime from some sufficiently large n onward, or some might be prime and some composite. Currently, we have no way of knowing which is true.

² Goldbach was the first to use the prime divisors of the Fermat numbers to prove that there are infinitely many primes, in a letter to Euler in late July 1730.

Lemma 1. *Let $g(x)$ be a polynomial with integer coefficients. For any integers $a \neq b$, their difference $a - b$ divides $g(a) - g(b)$. Moreover, if m is an integer for which $a \equiv b \pmod{m}$, then $g(a) \equiv g(b) \pmod{m}$.*

For $f(x) = x^2 - x + 1$, the orbit of 0 under the map f is $0 \rightarrow 1 \rightarrow 1 \rightarrow \dots$. Therefore if $n > m \geq 0$, then we have

$$E_n = f^{n-m}(E_m) \equiv f^{n-m}(0) = 1 \pmod{E_m},$$

and so $(E_n, E_m) = (1, E_m) = 1$. (For clarity, we have used an “=” sign when two numbers are equal despite working with a congruence. Typically we use the notation (a, b) rather than $\gcd(a, b)$.) Similarly if $f(x) = x^2 - 2x + 2$, then the orbit of 0 under the map f is $0 \rightarrow 2 \rightarrow 2 \rightarrow \dots$ and so $F_n = f^{n-m}(F_m) \equiv f^{n-m}(0) = 2 \pmod{F_m}$, which implies that if $n > m$, then $(F_n, F_m) = (2, F_m) = 1$.

This reformulation of two of the best-known proofs of the infinitude of primes hints at the possibility of a more general approach.

3. DIFFERENT STARTING POINTS. We just saw that the orbit of 2 under the map $x \rightarrow x^2 - x + 1$ is an infinite sequence of pairwise coprime integers, $2 \rightarrow 3 \rightarrow 7 \rightarrow 43 \rightarrow 1807 \rightarrow \dots$. What about other orbits? The orbit of 4 is also an infinite sequence of pairwise coprime integers beginning $4 \rightarrow 13 \rightarrow 157 \rightarrow 24493 \rightarrow \dots$, as is the orbit of 5 which begins $5 \rightarrow 21 \rightarrow 421 \rightarrow \dots$. The same proof as before yields that no two integers in a given orbit have a common factor.

The orbit of 3 under the map $x \rightarrow x^2 - 2x + 2$ yielded the Fermat numbers $3 \rightarrow 5 \rightarrow 17 \rightarrow 257 \rightarrow \dots$, but starting at 4 we get $4 \rightarrow 10 \rightarrow 82 \rightarrow 6562 \rightarrow \dots$. These are obviously not pairwise prime as every number in the orbit is even, but if we divide through by 2, then we get

$$2 \rightarrow 5 \rightarrow 41 \rightarrow 3281 \rightarrow \dots$$

which are pairwise coprime. To prove this, note that $x_0 = 4$ and $x_{n+1} = x_n^2 - 2x_n + 2$ for all $n \geq 0$, and so the above proof yields that if $m < n$ then $(x_m, x_n) = (x_m, 2) = 2$. Therefore, taking $a_n = x_n/2$ for every n we deduce that $(a_m, a_n) = (x_m/2, x_n/2) = (x_m/2, 1) = 1$. This same idea works for every orbit under this map: we get an infinite sequence of pairwise coprime integers by dividing through by 1 or 2, depending on whether x_0 is odd or even.

However, things can be more complicated: Consider the orbit of $x_0 = 3$ under the map $x \rightarrow x^2 - 6x - 1$. We have

$$3 \rightarrow -10 \rightarrow 159 \rightarrow 24326 \rightarrow 591608319 \rightarrow \dots$$

Here x_n is divisible by 3 if n is even, and is divisible by 2 if n is odd. If we let $a_n = x_n/3$ when n is even, and $a_n = x_n/2$ when n is odd, then one can show the terms of the resulting sequence,

$$1 \rightarrow -5 \rightarrow 53 \rightarrow 12163 \rightarrow 197202773 \rightarrow \dots,$$

are indeed pairwise coprime.

Another surprising example is given by the orbit of 6 under the map $x \rightarrow 7 + x^5(x - 1)(x - 7)$. Reducing the elements of the orbit mod 7 we find that

$$6 \rightarrow 5 \rightarrow 4 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 0 \rightarrow 0 \rightarrow \dots \pmod{7},$$

as $x^5(x-1)(x-7) \equiv x^6(x-1) \pmod{7}$, which is $\equiv x-1 \pmod{7}$ if $x \not\equiv 0 \pmod{7}$ by Fermat's little theorem. So x_n is divisible by 7 for every $n \geq 6$, but for no smaller n , and to obtain the pairwise coprime a_n we let $a_n = x_n$ for $n \leq 5$, and $a_n = x_n/7$ once $n \geq 6$.

It starts to look as though it might become complicated to formulate how to define a sequence $(a_n)_{n \geq 0}$ of pairwise coprime integers in general; certainly a case-by-case description is unappealing. However, there is a simpler way to obtain the a_n : In these last two examples, we have $a_n = x_n/(x_n, 6)$ and then $a_n = x_n/(x_n, 7)$, respectively, for all $n \geq 0$, a description that will generalize well.

4. DYNAMICAL SYSTEMS AND THE INFINITUDE OF PRIMES. One models evolution by determining the future development of the object of study from its current state (more sophisticated models incorporate the possibility of random mutations). This gives rise to *dynamical systems*, a rich and bountiful area of study. One simple model is that the state of the object at time n is denoted by x_n , and given an initial state x_0 , one can find subsequent states via a map $x_n \rightarrow f(x_n) = x_{n+1}$ for some given function $f(\cdot)$. Orbits of linear polynomials $f(\cdot)$ are easy to understand,³ but quadratic polynomials can give rise to evolution that is very far from what one might naively guess (the reader might look into the extraordinary Mandelbrot set).

This is the set-up that we had above! For us, $f(x)$ is a polynomial with integer coefficients and x_0 an integer. It will be useful to use dynamical systems terminology.

If $f^n(\alpha) = \alpha$ for some integer $n \geq 1$, then (the orbit of) α is *periodic*, and the smallest such n is the *exact period* length for α . The orbit begins with the *cycle*

$$\alpha, f(\alpha), \dots, f^{n-1}(\alpha)$$

of distinct values, and repeats itself, so that $f^n(\alpha) = \alpha$, $f^{n+1}(\alpha) = f(\alpha)$, \dots , and, in general $f^{n+k}(\alpha) = f^k(\alpha)$ for all $k \geq 0$.

The number α is *preperiodic* if $f^m(\alpha)$ is periodic for some $m \geq 0$, and is *strictly preperiodic* if α is preperiodic but not itself periodic. In all of our examples so far, 0 has been strictly preperiodic. In fact, if any two elements of the orbit of α are equal, say $f^{m+n}(\alpha) = f^m(\alpha)$, then $f^{k+n}(\alpha) = f^{k-m}(f^{m+n}(\alpha)) = f^{k-m}(f^m(\alpha)) = f^k(\alpha)$ for all $k \geq m$, so that α is preperiodic.

Finally, α has a *wandering orbit* if it is not preperiodic, that is, if its orbit never repeats itself so that the $\{f^m(\alpha)\}_{m \geq 0}$ are all distinct. Therefore, we wish to start only with integers x_0 that have wandering orbits.

We now state our general result for constructing infinitely many primes from orbits of a polynomial map.

Theorem 1. Suppose that $f(x) \in \mathbb{Z}[x]$, and that 0 is a strictly preperiodic point of the map $x \rightarrow f(x)$. Let $\ell(f) = \text{lcm}[f(0), f^2(0)]$. For any integer x_0 that has a wandering orbit, $(x_n)_{n \geq 0}$, let

$$a_n = \frac{x_n}{\gcd(x_n, \ell(f))} \text{ for all } n \geq 0.$$

The $(a_n)_{n \geq 0}$ are an infinite sequence of pairwise coprime integers, and if $n \geq 3$ then a_n has a private prime factor.

For example, $\ell(f) = 1, 2, 6$, and 7, respectively, for the four polynomials $f(x)$ in the examples of Section 3.

³ One can verify: if $f(t) = at + b$, then $x_n = x_0 + nb$ if $a = 1$, and $x_n = a^n x_0 + b/(1-a)$ if $a \neq 1$.

Proof. Suppose that $k = n - m > 0$. Then, by [Lemma 1](#),

$$x_n = f^k(x_m) \equiv f^k(0) \pmod{x_m},$$

and so $\gcd(x_m, x_n)$ divides $\gcd(x_m, f^k(0))$, which divides $f^k(0)$. But this divides

$$L(f) := \text{lcm}[f^k(0) : k \geq 1],$$

which is the lcm of a finite number of nonzero integers, as 0 is preperiodic. Therefore, (x_m, x_n) divides $L(f)$, and so (x_m, x_n) divides both $(x_m, L(f))$ and $(x_n, L(f))$. This implies that $A_m := x_m / (x_m, L(f))$ divides $x_m / (x_m, x_n)$, and A_n divides $x_n / (x_m, x_n)$. But $x_m / (x_m, x_n)$ and $x_n / (x_m, x_n)$ are pairwise coprime which therefore implies that $(A_m, A_n) = 1$.

We will show below that $L(f) = \ell(f)$, that is, the lcm of all the elements of the orbit of 0 is the same as the lcm of the first two terms. This then implies that $a_n = A_n$ for all n .

One can also prove that $|a_n| > 1$ for all $n \geq 3$ (see the discussion of one case in [Section 9](#)), and all such n must have a private prime factor. The result follows. ■

The example $f(x) = 3 - x(x - 3)^2$ with $0 \rightarrow 3 \rightarrow 3$ has a wandering orbit $2 \rightarrow 1 \rightarrow -1 \rightarrow \dots$, so that if $x_0 = 2$ then $x_1 = 1$ and $x_2 = -1$. Therefore, we cannot in general improve the lower bound, $n \geq 3$, in [Theorem 1](#).

We now determine all polynomials f that satisfy the hypothesis of [Theorem 1](#).

5. POLYNOMIAL MAPS FOR WHICH 0 IS STRICTLY PREPERIODIC.

There are severe restrictions on the possible exact periods.

If 0 is strictly preperiodic, then its exact period length is either one or two. We have already seen the examples $x^2 - x + 1$ for which $0 \rightarrow 1 \rightarrow 1$, and $x^2 - 6x - 1$ for which $0 \rightarrow -1 \rightarrow 6 \rightarrow -1$, where 0 is preperiodic with period length one and two, respectively. In fact exact periods cannot be any larger.

Lemma 2. *Let $f(x) \in \mathbb{Z}[x]$. If the orbit of a_0 is periodic, then its exact period length is either one or two.*

Proof. Let N be the exact period length so that $a_N = a_0$, and then $a_{N+k} = f^k(a_N) = f^k(a_0) = a_k$ for all $k \geq 1$. Now assume that $N > 1$ so that $a_1 \neq a_0$. [Lemma 1](#) implies that $a_{n+1} - a_n$ divides $f(a_{n+1}) - f(a_n) = a_{n+2} - a_{n+1}$ for all $n \geq 0$. Therefore,

$$\begin{aligned} a_1 - a_0 &\text{ divides } a_2 - a_1, \text{ which divides } a_3 - a_2, \dots, \text{ which divides} \\ a_N - a_{N-1} &= a_0 - a_{N-1}; \text{ and this divides } a_1 - a_N = a_1 - a_0, \end{aligned}$$

the nonzero number we started with. We deduce that

$$|a_1 - a_0| \leq |a_2 - a_1| \leq \dots \leq |a_{j+1} - a_j| \leq \dots \leq |a_1 - a_0|,$$

and so these are all equal. Therefore there must be some $j \geq 1$ for which $a_{j+1} - a_j = -(a_j - a_{j-1})$, else each $a_{j+1} - a_j = a_j - a_{j-1} = \dots = a_1 - a_0$ and so

$$0 = a_N - a_0 = \sum_{j=0}^{N-1} (a_{j+1} - a_j) = \sum_{j=0}^{N-1} (a_1 - a_0) = N(a_1 - a_0) \neq 0,$$

a contradiction. Therefore $a_{j+1} = a_{j-1}$, and we deduce that

$$a_2 = a_{N+2} = f^{N+1-j}(a_{j+1}) = f^{N+1-j}(a_{j-1}) = a_N = a_0,$$

as desired. ■

To determine which polynomials can be used in [Theorem 1](#), we need to understand the possible orbits of 0.

A classification of when 0 is a strictly preperiodic point. There are just four possibilities for the period length and the preperiod length of the orbit of 0. We now give examples of each type. For some nonzero integer a , the orbit of 0 is one of:

$0 \rightarrow a \rightarrow a \rightarrow \cdots$	for $x \rightarrow x^2 - ax + a$ (e.g., $(a_n)_{n \geq 0}$ and $(F_n)_{n \geq 0}$); or
$0 \rightarrow -a \rightarrow a \rightarrow a \rightarrow \cdots$	for $a = 1$ or 2 , with $x \rightarrow 2x^2 - 1$ or $x^2 - 2$, respectively; or
$0 \rightarrow -1 \rightarrow a \rightarrow -1 \rightarrow \cdots$	with $f(x) = x^2 - ax - 1$; or
$0 \rightarrow 1 \rightarrow 2 \rightarrow -1 \rightarrow 2 \rightarrow -1 \rightarrow \cdots$	with $f(x) = 1 + x + x^2 - x^3$; and

all other possible orbits are obtained from these, by using the observation that if $0 \rightarrow a \rightarrow b \rightarrow \cdots$ is an orbit for $x \rightarrow f(x)$, then $0 \rightarrow -a \rightarrow -b \rightarrow \cdots$ is an orbit for $x \rightarrow g(x)$ where $g(x) = -f(-x)$.

It is not difficult to determine all polynomials f for which 0 has one of these orbits. For example, in the first case above we simply need to find all f for which $f(0) = f(a) = a$. One example is $f_0(x) = a$ so that f is another example if and only if $f - f_0$ has roots at 0 and a . That is, $f(x) - f_0(x)$ equals $x(x - a)g(x)$ for some $g(x) \in \mathbb{Z}[x]$. The analogous approach works in all of the other cases.

Proving these are the only possible orbits is not difficult using [Lemma 1](#), but a little tedious. To give an example, let us determine the possible orbits of 0 with two elements in the preperiod and period length 1, that is, of the shape $0 \rightarrow b \rightarrow a \rightarrow a \rightarrow \cdots$ for distinct nonzero integers a and b . Now $b = b - 0$ divides $f(b) - f(0) = a - b$ and so b divides a . Moreover, $a = a - 0$ divides $f(a) - f(0) = a - b$ and so a divides b . Therefore $|b| = |a|$, and so $b = -a$ as a and b are distinct.

We make several deductions from analyzing the four cases of the classification:

1. 0 is strictly preperiodic if and only if $f^2(0) = f^4(0) \neq 0$;
2. $L(f) = \text{lcm}[f(0), f^2(0)] =: \ell(f)$, as claimed in the proof of [Theorem 1](#);
3. x_0 has a wandering orbit if and only if $x_2 \neq x_4$: To see this, note that if x_0 is periodic, then $x_{n+2} = x_n$ for all $n \geq 0$ as the period length is either one or two. Moreover, if x_0 is strictly preperiodic, then 0 is strictly preperiodic for the map $x \rightarrow f(x + x_0) - x_0$, with orbit $0 \rightarrow b_1 \rightarrow b_2 \rightarrow \cdots$ where $b_n = x_n - x_0$. In all four cases of our classification $b_2 = b_4$, and so $x_2 = x_4$.

6. PRIVATE PRIME FACTORS WHEN 0 IS A PERIODIC POINT. We now turn our attention to maps $x \rightarrow f(x)$ for which 0 is not strictly preperiodic, beginning

with when 0 is periodic with f of degree > 1 . We have seen that the period must have length one or two.

When 0 is periodic with period length 1. We must have $f(x) = x^r g(x)$ for some $g(x) \in \mathbb{Z}[x]$ with $g(0) \neq 0$, and $r \geq 1$. We will assume that $g(x)$ has degree ≥ 1 , else the integers in the orbit of x_0 contain only the prime factors of x_0 and of $g(0)$, so do not help us to construct infinitely many primes.

Theorem 2. *Suppose that $f(x) = x^r g(x)$ for some nonconstant $g(x) \in \mathbb{Z}[x]$ with $g(0) \neq 0$, and $r \geq 1$. For any given wandering orbit $(x_n)_{n \geq 0}$ with $x_0 \in \mathbb{Z}$, define*

$$a_{n+1} := \frac{g(x_n)}{\gcd(g(x_n), g(0))} \text{ for all } n \geq 0.$$

The $(a_n)_{n \geq 0}$ are an infinite sequence of pairwise coprime integers and, once n is sufficiently large, each a_n has a private prime factor.

Proof. Given a wandering orbit $(x_n)_{n \geq 0}$ we see that x_m divides $x_{m+1} = x_m^r g(x_m)$ for all m ; and so x_m divides x_n whenever $m \leq n$. Now a_{n+1} is a divisor of x_{n+1} for all $n \geq 0$. If $m < n$, then

$$(g(x_m), g(x_n)) \text{ divides } (x_{m+1}, g(x_n)), \text{ which divides } (x_n, g(x_n)), \\ \text{which divides } (x_n, g(0)), \text{ which divides } g(0).$$

Therefore $(a_{m+1}, a_{n+1}) = 1$, that is, the $(a_n)_{n \geq 1}$ are pairwise coprime, and so a_n has a private prime factor provided $|a_n| > 1$. Now $a_n \neq 0$, else $x_{n+1} = 0$, which would contradict the assumption that the orbit of x_0 is wandering. If $|a_n| = 1$, then $g(x_n)$ is a divisor d , of $g(0)$, that is, x_n must be a root of the polynomial

$$\prod_{d \text{ is a divisor of } g(0)} (g(x) - d).$$

But this has finitely many roots so, as the orbit of x_0 is wandering, there can only be finitely many n for which $|a_n| = 1$. Therefore, a_n must have a private prime factor for all sufficiently large n . ■

When 0 is periodic with period length 2. We rule out the polynomials $f(x) = a - x$ for any constant a , else there are no wandering orbits.

Theorem 3. *Suppose that $f(x) \in \mathbb{Z}[x]$ with $f(x) + x$ nonconstant, such that 0 is periodic under the map $x \rightarrow f(x)$, with period length 2. We write $f^2(x) = x^r G(x)$ with $G(0) \neq 0$ and $r \geq 1$. For any given wandering orbit $(x_n)_{n \geq 0}$ with $x_0 \in \mathbb{Z}$, define*

$$a_{n+2} := \frac{G(x_n)}{\gcd(G(x_n), G(0)f(0))} \text{ for all } n \geq 0.$$

The $(a_n)_{n \geq 0}$ are an infinite sequence of pairwise coprime integers and, once n is sufficiently large, each a_n has a private prime factor.

Proof. We begin by showing that $G(x)$ cannot be a constant, c . If so, then $f^2(x) = cx^r$ but $f(0) \neq 0$. Therefore, $f(x)$ has just one root, as $f^2(x)$ has at least as many distinct roots as f . Writing $f(x) = c(x - a)^d$ with $a \neq 0$, the roots of $f^2(x)$ are the roots of $c(x - a)^d - a$, each repeated d times. This implies that $d = 1$ as the roots of

$c(x - a)^d - a$ are distinct, and so $f^2(c) = c^2x - ac(c + 1)$ and so $c = -1$, that is, $f(x) = a - x$, which we have already ruled out.

Now, by definition, a_{n+2} is a divisor of x_{n+2} for all $n \geq 0$. Suppose that $n > m \geq 0$. If $n - m$ is even, then $(G(x_m), G(x_n))$ divides $G(0)$, by the argument of the previous subsection, and so $(a_{m+2}, a_{n+2}) = 1$. If $n - m$ is odd, then $G(x_m)$ divides x_{m+2} and so divides x_{n+1} . On the other hand, $G(x_n)$ divides $x_{n+2} = f(x_{n+1})$, and so $(G(x_m), G(x_n))$ divides $(x_{n+1}, f(x_{n+1})) = (x_{n+1}, f(0))$ which divides $f(0)$, and so $(a_{m+2}, a_{n+2}) = 1$.

Therefore, the $(a_n)_{n \geq 2}$ are pairwise coprime, and so a_n has a private prime factor provided $|a_n| > 1$. We then show, just as in the previous subsection, that $a_n \neq 0$, and there are only finitely many n for which $|a_n| = 1$. ■

We can combine the first three theorems to obtain the following result.

Corollary 1. *Suppose that $f(x) \in \mathbb{Z}[x]$ is not of the form cx^d or $a - x$, for any constants a or c , and that 0 is preperiodic for the map $x \rightarrow f(x)$. Then, for any given wandering orbit $(x_n)_{n \geq 0}$ with $x_0 \in \mathbb{Z}$, each x_n has a private prime factor for all sufficiently large n .*

We now turn our attention to the more difficult case of those f for which 0 itself has a wandering orbit.

7. POLYNOMIAL MAPS WHERE 0 HAS A WANDERING ORBIT. Henceforth we will suppose that 0 has a wandering orbit under the map $x \rightarrow f(x)$, for some polynomial $f(x) \in \mathbb{Z}[x]$. We will investigate whether the terms of wandering integer orbits all have private prime factors from some point onward. Where better to begin than with the orbit that begins with 0?

Orbits that begin with 0 are divisibility sequences. If $n = m + k$, then $x_n = f^k(x_m) \equiv f^k(0) = x_k \pmod{x_m}$, and so $(x_n, x_m) = (x_k, x_m)$. Subtracting m several times from n , we deduce that if $n = mq + r$ then $(x_n, x_m) = (x_r, x_m)$, which mirrors that $(n, m) = (r, m)$, the key step in the Euclidean algorithm. Therefore, by applying the same steps as in the Euclidean algorithm we deduce that

$$(x_m, x_n) = x_{(m,n)}.$$

Therefore $(x_n)_{n \geq 1}$ is a *strong divisibility sequence*; for example, $x_n = 2^n - 1$, the orbit of 0 under the map $x \rightarrow f(x) = 2x + 1$. Other examples of strong divisibility sequences include the Fibonacci numbers and any second-order linear recurrence sequence starting with 0, and the denominators of multiples of a given point on an elliptic curve.

A divisibility sequence cannot have private prime factors, because if prime p divides x_m then it divides x_n for every n divisible by m . Instead one looks for *primitive prime factors*, that is, primes p_n that divide x_n but do not divide x_m for any m , $1 \leq m < n$. It is known that for any second-order linear recurrence sequence beginning with a 0, and for any elliptic divisibility sequence, x_n has a primitive prime factor for all sufficiently large n . The two key ideas in the proof are to show that the x_n are fast-growing and that if p is prime factor of x_n , but not a primitive prime factor, then p but not p^2 divides $x_n/x_{n/p}$. In particular, one deduces that x_n is too large for all its prime power factors to be powers of imprimitive primes.

In the example $x_n = 2^n - 3$ (the orbit of -2 under the map $x \rightarrow 2x + 3$), and for general wandering orbits, one cannot be so precise about the power to which imprimitive primes p divide x_n , and so it is an open question as to whether the x_n all have

primitive prime factors for sufficiently large n . Nonetheless, we expect that this is true and, in the next subsection, we will consider this question following roughly the same outline.

The growth of the x_n and (x_m, x_n) . The x_n grow very fast, and we can gain a quite precise understanding: If $f(x) \in \mathbb{Z}[x]$ has degree $d > 1$ and if x_0 is an integer whose orbit is wandering, then there exist real numbers $1 \geq \alpha > 0$ and β , which depend only on f , for which

$$|x_n| \text{ is the integer nearest to } \alpha \tau^{d^n} + \beta, \text{ when } n \text{ is sufficiently large,} \quad (1)$$

where $\tau > 1$ is a constant that depends on both f and x_0 .⁴ (We will sketch a proof of (1) in Section 8, as well as find an easily determined sequence of approximations that converge rapidly to τ .) In the most interesting case, the orbit of 0, we write $\tau = \tau_0$.

Since $\alpha \leq 1$, (1) implies that $|x_r| \leq 2\tau^{d^r}$ and $|f^r(0)| \leq 2\tau_0^{d^r}$ if r is sufficiently large. Suppose that $0 \leq m \leq n-1$. Since (x_m, x_n) divides $f^{n-m}(0)$ as well as $|x_m|$, we deduce that

$$(x_m, x_n) \leq \min\{|x_m|, f^{n-m}(0)\} \leq 2\tau_*^{d^{n/2}} \quad (2)$$

if n is sufficiently large, where $\tau_* = \max\{\tau, \tau_0\}$.

We can therefore deduce that the x_n grow so fast that the product of all of the $\gcd(x_m, x_n)$ with $m < n$ is far smaller than x_n , so it seems very likely that x_n has primitive prime factors. However, this argument cannot be made into a proof since some of the primes that divide the x_m with $m < n$ might divide x_n to an extraordinarily high power.

Ruling out too many large prime power divisors. The *abc* conjecture can be used to bound solutions to equations in which all of the variables are divisible by surprisingly high powers of primes. The idea is to study pairwise coprime integer solutions to the equation

$$a + b = c.$$

The *abc*-conjecture states, in a quantitative form, that a , b , and c cannot all be divisible only by large powers of primes.⁵ For example, if we have a putative solution to Fermat's last theorem, like $x^n + y^n = z^n$ we can take $a = x^n$, $b = y^n$, and $c = z^n$, and the *abc*-conjecture should rule out any possible solution, at least once the numbers involved are sufficiently large. One does so by bounding the size of a , b , and c in terms of the product of the distinct primes that divide a , b , and c . To put this in quantitative form requires some ϵ 's and that sort of thing.

Fix $\epsilon > 0$ (which should be thought of as being small). There exists a constant $\kappa_\epsilon > 0$ (which depends on ϵ , and chosen so it works for all examples), such that if a and b

⁴ More generally, the *height*, $h(r)$, of a rational number $r = p/q$ is given by $\max\{|p|, |q|\}$; this equals $|r|$ if r is a nonzero integer. For any orbit $\{x_n\}_{n \geq 0}$ of a rational function $\phi(t) = f(t)/g(t)$, with $f(t), g(t) \in \mathbb{Z}[t]$ of degree d , there exists a constant $\tau > 0$ (the exponential of the *dynamical canonical height*, as in [1] or Section 3.4 of [8]), and constants $c_1, c_2 > 0$ such that $c_1 \tau^{d^n} < h(x_n) < c_2 \tau^{d^n}$.

⁵ Mochizuki has recently claimed to have proved the *abc*-conjecture, but the experts have as yet struggled to agree, definitively, that Mochizuki's extraordinary ideas are all correct.

are coprime positive integers with $c = a + b$, then

$$c \leq \kappa_\epsilon \left(\prod_{\substack{p \text{ prime} \\ p \text{ divides } abc}} p \right)^{1+\epsilon}.$$

This is the *abc-conjecture*, one of the great questions of modern mathematics. In the special case that $b = 1$, we have $c = a + 1$, and so

$$a \leq \kappa_\epsilon \left(\prod_{\substack{p \text{ prime} \\ p \text{ divides } a(a+1)}} p \right)^{1+\epsilon}.$$

The product here is over all primes p that divide the polynomial $f(x) = x^2 + x$ evaluated at $x = a$. Langevin [5] proved that a far-reaching generalization, which applies to the prime divisors of any given polynomial, is actually a consequence of the *abc-conjecture*.

Consequence of the *abc-conjecture*: Suppose that $f(x) \in \mathbb{Z}[x]$ has at least d distinct roots, for some $d \geq 2$. For any fixed $\epsilon > 0$, there exists a constant $\kappa_{\epsilon, f} > 0$, which depends on both ϵ and f , such that for any integer a we have

$$|a| \leq \kappa_{\epsilon, f} \left(\prod_{\substack{p \text{ prime} \\ p \text{ divides } f(a)}} p \right)^{1/(d-1)+\epsilon}.$$

We will apply this to our polynomial f with $\epsilon = 1$, writing κ for $\kappa_{1, f}$ for $x_n = f(x_{n-1})$. Therefore, if f has at least two distinct roots, then the *abc-conjecture* implies that

$$|x_{n-1}| \leq \kappa \left(\prod_{\substack{p \text{ prime} \\ p \text{ divides } x_n}} p \right)^2. \quad (3)$$

The product of the primes if x_n has no new prime factors. If every prime factor of x_n already divides a previous term in the sequence, then we have that

$$\prod_{\substack{p \text{ prime} \\ p \text{ divides } x_n}} p \text{ divides } \prod_{m=0}^{n-1} \prod_{\substack{p \text{ prime} \\ p \text{ divides } (x_m, x_n)}} p, \text{ which divides } \prod_{m=0}^{n-1} (x_m, x_n).$$

Taking the product of (2) over all m , $0 \leq m \leq n - 1$, we deduce that

$$\prod_{\substack{p \text{ prime} \\ p \text{ divides } x_n}} p \leq \prod_{m=0}^{n-1} (x_m, x_n) \leq 2^n \tau_*^{nd^{n/2}}$$

once n is sufficiently large. Substituting this into (3) gives⁶

$$|x_{n-1}| \leq \kappa 2^{2n} \tau_*^{2nd^{n/2}}.$$

Finally, comparing this to (1), taking logarithms, and dividing through by $\log \tau$, we see that there exists a constant $\gamma > 0$ (which can be given explicitly in terms of τ and τ_*) such that, for large enough n ,

$$d^n \leq \gamma n d^{n/2}.$$

This is false once n is sufficiently large, since d^n grows much faster than $nd^{n/2}$ as n increases. Therefore, we have proved the following conditional result.

Theorem 4. *Assume the abc-conjecture. Suppose that $f(x) \in \mathbb{Z}[x]$ has at least two distinct roots, and that 0 and x_0 both have wandering orbits under the map $x \rightarrow f(x)$. Then x_n has a primitive prime factor for all sufficiently large n .*

Note that this includes the possibility that $x_0 = 0$; and so $f^n(0)$ has a primitive prime factor for all sufficiently large n , assuming the *abc*-conjecture.

Primitive prime factors of rational functions. Now suppose that $\phi(t) = f(t)/g(t)$, with $f(t), g(t) \in \mathbb{Z}[t]$. Beginning with a rational number r_0 , we define $r_{n+1} = \phi(r_n)$ for all $n \geq 0$, and write $r_n = a_n/b_n$ where $\gcd(a_n, b_n) = 1$ and $b_n \geq 1$.

Suppose that $f(0) = 0$ but $f(t)$ is not of the form at^d . Ingram and Silverman [4] showed that if the orbit of r_0 is not eventually periodic, then a_n has a primitive prime factor for all sufficiently large n . It is believed that some result of this sort should hold whether or not $f(0) = 0$, and indeed this has been confirmed by Gratton, Nguyen, and Tucker ([3]) under the assumption of the *abc*-conjecture.

8. ASYMPTOTIC GROWTH OF WANDERING ORBITS. Fix an integer $d > 1$. Let $(y_n)_{n \geq 0}$ be any sequence of positive integers for which

$$y_n^d < y_{n+1} < (y_n + 1)^d \text{ for all } n \geq 0.$$

If $\ell_n = y_n^{1/d^n}$ and $u_n = (y_n + 1)^{1/d^n}$, then $\ell_n < \ell_{n+1} < \dots < u_{n+1} \leq u_n$. The ℓ_n are therefore an increasing bounded sequence, so must tend to a limit, call it τ . Then $\ell_n < \tau \leq u_n$ for all n and so $y_n < \tau^{d^n} \leq y_n + 1$, with equality in the upper bound if and only if $y_n = \tau^{d^n} - 1$ for all n . Therefore y_n is the largest integer $< \tau^{d^n}$. In fact $y_n = \lfloor \tau^{d^n} \rfloor$, other than in the special case in which τ is an integer and $y_n = \tau^{d^n} - 1$ for all n .

Mills [6] observed, in 1947, that advances in our understanding of the distribution of primes imply that if N is sufficiently large, then there is a prime between N^3 and $(N + 1)^3$. So taking $p_0 = y_0$ to be any prime greater than N , and then selecting p_{n+1} to be any prime in $(p_n^3, (p_n + 1)^3)$ for each $n \geq 0$, one obtains a *Mills' constant*, τ , such that

$$\lfloor \tau^{3^n} \rfloor \text{ is the prime } p_n \text{ for every integer } n \geq 0.$$

This proof yields a lot of flexibility in constructing τ with this property (as we expect many primes in any interval of the form $(N^3, (N + 1)^3)$), so it seems unlikely that these numbers τ are anything special beyond the beautiful property identified by Mills.

⁶ An analogous argument is used to bound the imprimitive prime factors of recurrence sequences, and in a related context in [4].

Sketch of a modification to prove (1). For any polynomial $f(x) = ax^d + bx^{d-1} + \dots$ with $a > 0$, the polynomial

$$\alpha^{-1}(f(x) - \beta) - (\alpha^{-1}(x - \beta))^d,$$

where $\alpha = a^{-1/(d-1)}$ and $\beta = -b/(da)$, has degree $\leq d - 2$. Given an integer orbit $(x_n)_{n \geq 0}$, let $y_n = \alpha^{-1}(x_n - \beta)$, so that $y_{n+1} - y_n^d$ can be written as a polynomial in y_n of degree $\leq d - 2$ (where the coefficients of the polynomial are independent of n). This implies that, for any $\epsilon > 0$, if n is sufficiently large, then

$$(y_n - \epsilon)^d + \epsilon < y_{n+1} < (y_n + \epsilon)^d - \epsilon.$$

One can then modify the proof above to show that $(y_n - \epsilon)^{1/d^n}$ tends to a limit τ and that $y_n - \epsilon < \tau^{d^n} < y_n + \epsilon$. This implies that

$$x_n = \alpha y_n + \beta \in \left(\alpha \tau^{d^n} + \beta - \frac{1}{2}, \alpha \tau^{d^n} + \beta + \frac{1}{2} \right)$$

taking $\epsilon = 1/2\alpha$, and so x_n is the nearest integer to $\alpha \tau^{d^n} + \beta$ as claimed. ■

There is a power series $P(T) = c_1 T + c_0 + \frac{c_{-1}}{T} + \frac{c_{-2}}{T^2} + \dots$ which satisfies the *functional equation*

$$P(x^d) = f(P(x)).$$

To determine the c_i , we compare the coefficients on both sides of the equation: Comparing the coefficients of T^d implies that $c_1 = \alpha$; comparing the coefficients of T^{d-1} implies that $c_0 = \beta$. For each $j \geq 0$, the coefficients of T^{d-1-j} involve polynomials in the c_i with $i > -j$ as well as a linear term in c_{-j} , which allows us to determine c_{-j} .

Assuming that $P(T)$ absolutely converges once $|T|$ is sufficiently large, if x_m is large enough there is a unique positive real t for which $P(t) = x_m$. Then

$$P(t^d) = f(P(t)) = f(x_m) = x_{m+1},$$

and $P(t^{d^2}) = f(P(t^d)) = f(x_{m+1}) = x_{m+2}$, etc., so that $P(t^{d^k}) = x_{m+k}$ for all $k \geq 0$. Taking $\tau = t^{1/d^m}$ we deduce that

$$x_n = P(\tau^{d^n})$$

for all sufficiently large n .

One can sidestep the convergence issue by working with finite length truncations of $P(T)$. Thus if $P_k(t) = \sum_{j=1-k}^1 c_j t^j$, one can prove there exists a constant $C_k > 0$ such that

$$|x_n - P_k(\tau^{d^n})| \leq C_k / \tau^{kd^n}$$

for all $n \geq 0$.

An easily determined sequence of approximations that converge rapidly to τ . In correspondence, Soundararajan noted that the product

$$a^{1/(d-1)x_0} \cdot \left(\frac{x_1}{ax_0^d}\right)^{1/d} \cdot \left(\frac{x_2}{ax_1^d}\right)^{1/d^2} \cdots \left(\frac{x_n}{ax_{n-1}^d}\right)^{1/d^n}$$

telescopes to $(a^{1/(d-1)}x_n)^{1/d^n} = (\alpha^{-1}x_n)^{1/d^n}$, which converges to τ as $n \rightarrow \infty$. Since $x_n = f(x_{n-1}) = ax_{n-1}^d + bx_{n-1}^{d-1} + \cdots$, we see that the quotients x_n/ax_{n-1}^d converge rapidly to 1, and therefore Soundararajan's sequence converges rapidly to τ .

The mysteries of the numbers τ ? Although this last subsection allows one to show the existence of the constant τ , it gives no hint as to what kind of number τ is. The τ corresponding to the Fermat numbers is 2. But what kind of number is τ for Euclid's sequence $(E_n)_{n \geq 0}$? Is it rational? Irrational? Related to π ? I have no idea. Also, in general, is there anything special about the numbers τ_0 , the τ -values that arise out of the wandering orbits of 0? For a given $f(x)$, what does the set of τ -values look like as we vary over all wandering orbits? Is each value isolated, or does the set have limit points? If we also vary over all f of given degree d , we have a countable subset of $\mathbb{R}_{>1}$. Are the τ dense in $\mathbb{R}_{>1}$? Or are there holes in the spectrum? If so, why? Some of these questions are addressed in Section 3.4 of [8], and there are some particularly interesting results and conjectures there on lower bounds for values of τ that are > 1 in terms of the given map and the field of definition. Gaining a better understanding of the numbers τ seems like a subject that is ripe for further and broader study.

9. APPENDIX: PRIVATE PRIME FACTORS FOR ALL $N \geq 3$. One can improve Theorem 1 by noting that if $|a_n| = 1$ (that is, x_n is a divisor of $L(f)$) then $n = 0, 1$, or 2. Our proof is given by a tedious case-by-case analysis. For example, when $0 \rightarrow a \rightarrow a$ with $a > 0$ we proceed as follows.

Suppose that $x_{n+2} = d$ is a divisor of a , for some $n \geq 0$, so that $x_n \rightarrow x_{n+1} \rightarrow d$ where $x_n, x_{n+1}, x_{n+2} = d, a$ are distinct. Now $x(x - a)$ divides $f(x) - a$ and so $x_{n+1}(a - x_{n+1})$ divides $a - x_{n+2} = a - d$, and $x_n(a - x_n)$ divides $x_{n+1} - a$. Moreover, as d divides a and $d \neq a$ we have $1 \leq a - d \leq 2a$.

Let $T = \min\{x_{n+1}, a - x_{n+1}\}$ so that $T(a - T) = x_{n+1}(a - x_{n+1})$ divides $a - d$. Now $T \geq -1$, else $2(a + 2) \leq |T(a - T)| \leq 2a$. If $T = -1$, then $a + 1$ divides $a - d$, so that $d = -1$, and therefore $x_{n+1} = a + 1$ as $x_{n+1} = T$ or $a - T$ and $x_{n+1} \neq d$. Finally, $x_n(a - x_n)$ divides $x_{n+1} - a = 1$ and so $a = 2$ and $x_n = 1$. That is, $a = 2$ and we have $1 \rightarrow 3 \rightarrow -1$.

We may now assume that $T \geq 1$, so that $a - 1 \leq |x_n(a - x_n)| = a - x_{n+1} \leq a - T \leq a - 1$. This implies that $x_{n+1} = T = 1$ and $x_n = a - 1$ as $x_n \neq x_{n+1}$, and $a \geq 2$ as $x_n \neq 0, x_{n+1}$. Moreover, $a - 1$ divides $a - d$ and $d \neq 1$ or a , so $d = 1 + k(1 - a)$ for some $k \geq 1$. If $k \geq 2$, then $|1 + k(1 - a)| > a$ unless $k = 2$ and $a = 3$, which gives the example $2 \rightarrow 1 \rightarrow -3$. For $k = 1$, we have $2 - a$ divides a only if $2 - a$ divides 2 and so $a = 3$ or 4. The $a = 3$ case gives rise to the example $2 \rightarrow 1 \rightarrow -1$. The $a = 4$ case does not work, else $2 = x_n - x_{n+1}$ would divide $f(x_n) - f(x_{n+1}) = x_{n+1} - x_{n+2} = x_{n+1} - d = 3$.

Now n must be 0 in all of these three examples else $x_{n-1} \rightarrow x_n \rightarrow x_{n+1} \rightarrow d$ for some $n \geq 1$. But $a = 3$ with $x_{n-1} \rightarrow 2 \rightarrow 1$ is impossible else $x_{n-1} - 2$ divides $2 - 1 = 1$, so that $x_{n-1} = t$ or $x_{n-1} = a$. And $a = 2$ with $x_{n-1} \rightarrow 1 \rightarrow 3 \rightarrow -1$ is impossible else $x_{n-1} - 1$ divides $3 - 1 = 2$ and $x_{n-1} - 3$ divides $1 - (-1) = 2$, which together imply that $x_{n-1} = 2 = a$. ■

ACKNOWLEDGMENTS. This article was first conceived of in 1997 following invigorating conversations about arithmetic dynamics with Mike Zieve. My understanding of arithmetic dynamics subsequently greatly benefitted from discussions with Jeff Lagarias and Xander Faber. Thanks to them all. I would also like to thank the referees for their many helpful suggestions, and Soundararajan for his observation discussed in Section 8.

ORCID. Andrew Granville  <http://orcid.org/0000-0001-8088-1247>

REFERENCES

- [1] Call, G. S., Silverman, J. H. (1993). Canonical heights on varieties with morphisms. *Compos. Math.* 89: 163–205.
- [2] Edwards, A. W. F. (1964). Infinite coprime sequences. *Math. Gaz.* 48: 416–422.
- [3] Gratton, C., Nguyen, K., Tucker, T. J. (2013). ABC implies primitive prime divisors in arithmetic dynamics. *Bull. Lond. Math. Soc.* 45: 1194–1208.
- [4] Ingram, P., Silverman, J. H. (2009). Primitive divisors in arithmetic dynamics. *Math. Proc. Cambridge Philos. Soc.* 146: 289–302.
- [5] Langevin, M. (1995). Partie sans facteur carré de $F(a, b)$ (modulo la conjecture (abc)). *Séminaire de Théorie des Nombres*, 93–94. Publ. Math. Univ. Caen.
- [6] Mills, W. H. (1947). A prime-representing function. *Bull. Amer. Math. Soc.* 53: 604.
- [7] Rice, B. (2007). Primitive prime divisors in polynomial arithmetic dynamics. *Integers (electronic)*, 7: A26 16 pages.
- [8] Silverman, J. H. (2007). *The Arithmetic of Dynamical Systems*. Graduate Texts in Mathematics, Vol. 241. New York: Springer.

ANDREW GRANVILLE specializes in understanding the distribution of primes. He is co-author of the soon-to-appear graphic novel, *MSI: Anatomy and Permutations* (Princeton University Press).

Département de Mathématiques et de Statistique, Université de Montréal, CP 6128 succ. Centre-Ville, Montréal, QC H3C 3J7, Canada

andrew@dms.umontreal.ca

Department of Mathematics, University College London, Gower Street, London, WC1E 6BT, United Kingdom
a.granville@ucl.ac.uk